ISSN 0021-9673

CUMULATIVE AUTHOR AND SUBJECT INDEXES VOLUMES 326–350

JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

EDITOR, Michael Lederer (Switzerland)
ASSOCIATE EDITOR, K. Macek (Prague)
EDITOR, SYMPOSIUM VOLUMES, E. Heftmann (Orinda, CA)
EDITORIAL BOARD

W. A. Aue (Halifax)
V. G. Berezkin (Moscow)
V. Betina (Bratislava)
A. Bevenue (Belmont, CA)
P. Boček (Brno)
P. Boulanger (Lille)
A. A. Boulton (Saskatoon)
G. P. Cartoni (Rome)
S. Dilli (Kensington, N.S.W.)
L. Fishbein (Jefferson, AR)
R. W. Frei (Amsterdam)
A. Frigerio (Milan)
C. W. Gehrke (Columbia, MO)
E. Gil-Av (Rehovot)
G. Guiochon (Palaiseau)
I. M. Hais (Hradec Králové)
J. K. Haken (Kensington, N.S.W.)
S. Hjertén (Uppsala)
E. C. Horning (Houston, TX)
Cs. Horváth (New Haven, CT)
J. F. K. Huber (Vienna)
A. T. James (Sharnbrook)
J. Janák (Brno)
E. sz. Kováts (Lausanne)
K. A. Kraus (Oak Ridge, TN)
E. Lederer (Gif-sur-Yvette)
A. Liberti (Rome)
H. M. McNair (Blacksburg, VA)
Y. Marcus (Jerusalem)
G. B. Marini-Bettòlo (Rome)
A. J. P. Martin (Lausanne)
Č. Michalec (Prague)
R. Neher (Basel)
G. Nickless (Bristol)
N. A. Parris (Wilmington, DE)
R. L. Patience (Sunbury-on-Thames)
P. G. Righetti (Milan)
O. Samuelson (Göteborg)
R. Schwarzenbach (Dübendorf)
G. Semenza (Zürich)
L. R. Snyder (Yorktown Heights, NY)
A. Zlatkis (Houston, TX)

EDITORS, BIBLIOGRAPHY SECTION
Z. Deyl (Prague), J. Janák (Brno), K. Macek (Prague)

UNIVERSITY OF HAWAII
JUL 18 1986
LIBRARIES

R.
D
71
56

ELSEVIER

JOURNAL OF CHROMATOGRAPHY

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications,* which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis (including the publication of normal values); screening and profiling procedures with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; combinations of chromatographic and electrophoretic methods with other physicochemical techniques such as mass spectrometry. In *Chromatographic Reviews,* reviews on all aspects of chromatography, electrophoresis and related methods are published.

Submission of Papers. Papers in English, French and German may be submitted, in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Review articles are invited or proposed by letter to the Editors and will appear in *Chromatographic Reviews* or *Biomedical Applications.* An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Subscription Orders. Subscription orders should be sent to: Elsevier Science Publishers B.V., P.O. Box 211, 1000 AE Amsterdam, The Netherlands. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications, Chromatographic Reviews* and *Cumulative Author and Subject Indexes, Vols. 326–350*) has 38 volumes in 1986. The subscription prices for 1986 are:

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes, Vols. 326–350*) + *Biomed. Appl.* (Vols. 346–383): Dfl. 6080.00 plus Dfl. 912.00 (postage) (total *ca.* US$ 2742.00)

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes, Vols. 326–350*) only (Vols. 346–373): Dfl. 5040.00 plus Dfl. 672.00 (postage) (total *ca.* US$ 2240.00)

Biomed. Appl. only (Vols. 374–383): Dfl. 1850.00 plus Dfl. 240.00 (postage) (total *ca.* US$ 819.50).

Journals are automatically sent by airmail at no extra costs to Argentina, Australia, Brasil, Canada, China, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan, Thailand and the U.S.A. Back volumes of the *Journal of Chromatography* (Vols. 1 through 345) are available at Dfl. 219.00 (plus postage). Claims for issues not received should be made within three months of publication of the issue. If not, they cannot be honoured free of charge. Customers in the U.S.A. and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Publishing Co. Inc., 52 Vanderbilt Avenue, New York, NY 10017. Tel. (212) 916-1250.

Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal and Science Citation Index.

See page 3 of cover for Publication Schedule, Information for Authors and information on Advertisements.

© ELSEVIER SCIENCE PUBLISHERS B.V. — 1986

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Printed in The Netherlands

JOURNAL *of* CHROMATOGRAPHY

INTERNATIONAL JOURNAL OF CHROMATOGRAPHY,
ELECTROPHORESIS AND RELATED METHODS

Cumulative Author and Subject Indexes Volumes 326–350 (1985)

ELSEVIER
AMSTERDAM — OXFORD — NEW YORK — TOKYO
1986

Electronic data processing:
Büro für Satztechnik W. Meyer KG
Weissensberg, F.R.G.

© ELSEVIER SCIENCE PUBLISHERS B.V. — 1986

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Printed in The Netherlands

AUTHOR INDEX

VOLUMES 326–350

A

Aarons, L., see Adedoyin, A. 345(1985)192
Aarons, L., see Fong, M.H. 333(1985)191
Abdel-Aziz, H., see Chang, C.A. 347(1985)51
Abdel-Hay, M.H., see Malspeis, L. 343(1985)190
Abdullah, M., Davis, R.J.H. and Hill, J.A.
Application of DNA-cellulose chromatography in the isolation of immunoglobulin M and complement component C4b-binding protein from human serum 347(1985)129
Abe, K., see Tatsuhara, T. 339(1985)149
Abe, M., see Komuro, C. 338(1985)209
Abernethy, D.R., Schwartz, J.B. and Todd, E.L.
Diltiazem and desacetyldiltiazem analysis in human plasma using HPLC: improved sensitivity without derivatization 342(1985)216
Abouleish, E., see Shao, McN.J. 345(1985)184
Abraham, E.C., see Bissé, E. 344(1985)81
Abramson, F.P. and Lutz, M.P.
Application of a capillary GC-selected-ion recording MS technique to the analysis of diethylstilbestrol and its phosphorylated precursors in plasma and tissues 339(1985)87
Abreu, R.A. de, see Van Baal, J.M. 336(1985)422
Adair, C.G., Burns, D.T., Crockard, A.D., Desai, Z.R. and Harriott, M.
Modified extraction and chromatography for the measurement of plasma melphalan by ion-pair HPLC 336(1985)429
Adair, C.G., Burns, D.T., Crockard, A.D. and Harriott, M.
Determination of chlorambucil in plasma using reversed-phase HPLC 342(1985)447
Adair, W.L., see Keller, R.K. 343(1985)413
Adams, A.K., see Van Engelen, D.L. 331(1985)77
Adams, J. and Giese, R.W.
Cinnamoylation of the sugar hydroxyls of 3-methylthymidine. Formation of a relatively stable ester derivative 347(1985)99
Adams, P.S. and Haines-Nutt, R.F.
HPLC analysis of methadone hydrochloride in pharmaceuticals 329(1985)438
Adams, R.F., Jones, R.L. and Conway, P.L.
HPLC of microbial acid metabolites 336(1985)125
Adedoyin, A., Aarons, L. and Houston, J.B.
HPLC method for the simultaneous determination of cimetidine and antipyrine in plasma 345(1985)192
Adeney, J.A., see Gerritse, R.G. 347(1985)419
Adlard, M.W., see Rogers, M.E. 326(1985)163
Adzet, T. and Puigmacia, M.
HPLC of caffeoylquinic acid derivatives of *Cynara scolymus* L. leaves 348(1985)447
Agathonos, P., see Katsanos, N.A. 349(1985)369
Ageta, T., Tsutsumi, A., Ishizu, H., Mizoguchi, N. and Kodama, H.
Determination of free choline in human semen using an isotachophoretic analyser 343(1985)186
Agrawal, O.P., see Kharbade, B.V. 347(1985)447
Aguilar, M.-I., Hodder, A.N. and Hearn, M.T.W.
HPLC of amino acids, peptides and proteins. LXV Studies on the optimisation of the reversed-phase gradient elution of polypeptides: evaluation of retention relationships with β-endorphin-related polypeptides 327(1985)115
Aguilar, M.-I., see Hearn, M.T.W. 327(1985)47
Ahlborg, U.G., see Neelakantan, L. 350(1985)307
Ahnoff, M., Ervik, M., Lagerström, P.-O., Persson, B.-A. and Vessman, J.
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73
Akamatsu, N., see Ishii, I. 345(1985)134
Akella, K.S.R., see Husain, S. 348(1985)309
Åkesson, B., see Michelsen, P. 350(1985)417
Akira, K., see Baba, S. 341(1985)251
Akita, M., see Takayanagi, H. 350(1985)75
Akiyama, H., Kudo, S., Odomi, M. and Shimizu, T.
HPLC procedure for the determination of a new antithrombotic and vasodilating agent, cilostazol, in human plasma 338(1985)456

Al Makdessi, S., Andrieu, J.-L., Bacconin, A., Fugier, J.-C., Herilier, H. and Faucon, G.
Assay of lipids in dog myocardium using capillary GC and derivatization with boron trifluoride and methanol 339(1985)25
Alaerts, G., see Termonia, M. 328(1985)367
Alaupovic, P., see McConathy, W.J. 342(1985)47
Albarella, J.P., see Yip, K.F. 326(1985)301
Albert, K., Nieder, M., Bayer, E. and Spraul, M.
Continuous-flow nuclear magnetic resonance 346(1985)17
Albertsson, P.-Å., see Ekberg, B. 333(1985)211
Albin, H.C., see Demotes-Mainard, F.M. 336(1985)438, 342(1985)234
Alexander, I.S., see So, N. 337(1985)81
Alexander, L.R., Justice, J.B. Jr. and Madden, J.
Fatty acid composition of human erythrocyte membranes by capillary GC-MS 342(1985)1
Allen, I.C. and Griffiths, R.
Reversed-phase HPLC method for determination of brain glutamate decarboxylase suitable for use in kinetic studies 336(1985)385
Alley, C.C., see Brooks, J.B. 339(1985)243
Allinquant, B., Musenger, C. and Schuller, E.
Reversed-phase HPLC of nucleotides and oligonucleotides 326(1985)281
Almog, R., see Forward, R. 330(1985)383
Alpert, C.A., Dörschug, M., Saffen, D., Frank, R., Deutscher, J. and Hengstenberg, W.
The bacterial phosphoenolpyruvate-dependent phosphotransferase system. Isolation of active site peptides by reversed-phase HPLC and determination of their primary structure 326(1985)363
Altemura, P., see Rosini, C. 348(1985)79
Alvarado, X., see Ladrón de Guevara, O. 349(1985)91
Amato, A., see Gagliardi, L. 348(1985)321
Ambrus, G., see Szederkényi, F. 333(1985)404
Ames, M.M., Miller, K.J. and Moertel, D.M.
HPLC assay and preclinical pharmacological studies of pibenzimol (bisbenzimidazole) 341(1985)89
Amin, P.R., see Stetson, P.L. 341(1985)217
Anderson, D.J., Taylor, A.J. and Reischer, R.J.
Reversed-phase HPLC of protected oligodeoxynucleotides 350(1985)313
Anderson, D.J. and Walters, R.R.
Affinity chromatographic examination of a retention model for macromolecules 331(1985)1
Anderson, J.M.
Simultaneous determination of abscisic acid and jasmonic acid in plant extracts using HPLC 330(1985)347
Andersson, T., Carlsson, M., Hagel, L., Pernemalm, P.-Å. and Janson, J.-C.
Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33
Ando, T., see Fujimura, K. 350(1985)371
Ando, T., see Otsuka, K. 332(1985)219, 348(1985)39
Ando, T., see Takayanagi, H. 350(1985)63, 350(1985)75
Ando, T., see Terabe, S. 332(1985)211
Andrawes, F. and Deng, P.
Optimization of helium ionization detector 349(1985)405
Andrawes, F., see Deng, P. 349(1985)415
Andreolini, F., Borra, C., Di Corcia, A., Samperi, R. and Raponi, G.
Total oestriol in maternal serum or plasma as measured by LC 337(1985)379
Andrews, A.G.
Estimation of amniotic fluid phospholipids by HPLC 336(1985)139
Andrews, A.T., Taylor, M.D. and Owen, A.J.
Rapid analysis of bovine milk proteins by fast protein LC 348(1985)177
Andrieu, J.-L., see Al Makdessi, S. 339(1985)25
Andronikashvili, T.G., Berezkin, V.G., Laperashvili, L.Y. and Nadiradze, N.A.
On the possibility of increasing the effectiveness of chromatographic columns filled with zeolites 331(1985)402
Ang, K.P., see Gunasingham, H. 341(1985)271
Angelo, H.R., Kranz, T., Strøm, J., Thisted, B. and Sørensen, M.B.
HPLC method for the determination of dextropropoxyphene and nordextropropoxyphene in serum 345(1985)413
Änggård, E., see Blom, Y. 338(1985)89
Angle, C., see Cook, L. 339(1985)293
Ansari, G.A.S. and Hendrix, P.Y.
Rapid and convenient separation of pentachlorophenol from human fat using silica Sep-Pak cartridges 346(1985)435
Ansari, K., see Kaplan, E. 350(1985)435
Antalík, M., see Podhradský, D. 328(1985)402
Antoku, Y., Sakai, T. and Iwashita, H.
Fatty acid analysis in phosphatidylethanolamine subclasses of human erythrocyte membranes by HPLC 342(1985)359
Antonio, S., see Ladrón de Guevara, O. 349(1985)91
Aoki, I., see Tadano, K. 341(1985)228
Aoshima, S., see Oka, K. 345(1985)419
Applegarth, D.A., see Park, C.M. 339(1985)182
Arai, Y., see Hanai, T. 349(1985)323
Arakawa, M., see Kurosawa, N. 343(1985)379
Araki, M., see Tanaka, N. 332(1985)57
Arata, J., see Mikasa, H. 343(1985)179
Aravind, M.K., Miceli, J.N. and Kauffman, R.E.
Measurement of cyclosporin by HPLC following charcoal adsorption from whole blood 344(1985)428
Arendt, G., see Eppert, G. 350(1985)471
Arima, M., see Iwasaki, S. 341(1985)182
Arita, T., see Tadano, K. 341(1985)228
Armstrong, D.W., see Chang, C.A. 347(1985)51
Arnold, F.H., Blanch, H.W. and Wilke, C.R.
LC plate height equations 330(1985)159

Arnts, R.R.
Precolumn sample enrichment device for analysis of ambient volatile organics by GC-MS 329(1985)399
Aronsson, E., see Michelsen, P. 350(1985)417
Arunyanart, M. and Cline Love, L.J.
Determination of drugs in untreated body fluids by micellar chromatography with fluorescence detection 342(1985)293
Asai, H., see Ishii, D. 332(1985)3
Asai, H., see Takeuchi, T. 331(1985)99
Ascalone, V., Catalani, P. and Dal Bo', L.
Determination of progabide and its acid metabolite in biological fluids by HPLC on silica column with UV detection 344(1985)231
Asmus, P.A.
Determination of 2-(4-isobutylphenyl)propionic acid in bulk drug and compressed tablets by reversed-phase HPLC 331(1985)169
Aubert, C., see Coassolo, P. 338(1985)347
Aubert, C., see Tomasini, J.-L. 343(1985)369
Aulsebrook, K.A., see Roberts, L.J. II 338(1985)41
Avery, N.C. and Light, N.
Re-packing reversed-phase HPLC columns as means of regenerating column efficiency and prolonging packing life 328(1985)347
Awadzi, K., see Edwards, G. 343(1985)224

B

Baal, J.M. van, see Van Baal, J.M. 336(1985)422
Baars, A.M., see Vree, T.B. 343(1985)449
Baaske, D.M., see Karnatz, N.N. 330(1985)420
Baba, S., Akira, K., Horie, M. and Mori, Y.
Application of radioisotope tracer techniques to analytical GC: determination of GC peak yield 341(1985)251
Baba, S., Fujioka, M., Shinohara, Y. and Furuta, T.
Determination of testosterone propionate in human plasma by GC-MS 337(1985)205
Baba, S., see Minagawa, K. 343(1985)231
Baba, S., see Shinohara, Y. 338(1985)281
Baba, Y., Yoza, N. and Ohashi, S.
Effect of column temperature on HPLC behaviour of inorganic polyphosphates. I. Isocratic ion-exchange chromatography 348(1985)27
Baba, Y., Yoza, N. and Ohashi, S.
Effect of column temperature on HPLC behaviour of inorganic polyphosphates. II. Gradient ion-exchange chromatography 350(1985)119
Baba, Y., Yoza, N. and Ohashi, S.
Computer-assisted prediction of retention times for inorganic polyphosphates in gradient ion-exchange chromatography 350(1985)461
Bacconin, A., see Al Makdessi, S. 339(1985)25
Bacic, A., see Harris, P.J. 350(1985)304
Bacon, B.R., see Mostow, N.D. 344(1985)137
Baert, E., see Nelis, H.J.C.F. 333(1985)381
Baffi, F., see Mazzucotelli, A. 349(1985)137
Bagchi, S.P., Lutz, T. and Jindal, S.P.
GC-MS determination of nomifensine using a stable isotope-labeled analogue as an internal standard 344(1985)362
Bagnasco, G., see Verzella, G. 349(1985)83
Bailey, C.J., see Richfield-Fratz, N. 331(1985)109
Bailey, J.E. Jr.
Determination of the lower sulfonated subsidiary colors in FD&C Yellow No. 6 by reversed-phase HPLC 347(1985)163
Bailey, J.E. Jr., see Richfield-Fratz, N. 331(1985)109
Baker, I., Shuker, D.E.G. and Tannenbaum, S.R.
A method for the determination of S-methylcysteine by HPLC: application to the study of carcinogenic methylating agents 329(1985)202
Baker, J.O., Tucker, M.P. and Himmel, M.E.
Noble, diatomic and aliphatic gas analysis by aqueous HPLC 346(1985)93
Baker, L.H., see Schilcher, R.B. 337(1985)55
Baker, R.W.R., see Cullen, M.P. 341(1985)420
Baktir, G., Bircher, J., Fisch, H.-U. and Karlaganis, G.
Capillary GLC determination of the benzodiazepine triazolam in plasma using a retention gap 339(1985)192
Balansard, G., Schwadrohn, G., Vidal, E. and Elias, R.
Determination of ophthalmic therapeutic trifluorothymidine and its degradation product by reversed-phase HPLC 348(1985)299
Balansard, G., see Vidal, E. 348(1985)304
Baldwin, R.P., see Halbert, M.K. 345(1985)43
Baldwin, R.P., see Ravichandran, K. 343(1985)99
Ballantyne, D.J., see Perrigo, B.J. 341(1985)81
Ballester, P., see García-Raso, A. 331(1985)406
Ballschmiter, K., see Magg, H. 331(1985)245
Balzer, H.O., see Husseini, H.S. 329(1985)300
Bandi, Z.L. and Reynolds, E.S.
Adsorption and reversed-phase HPLC of *p*-nitrobenzyl esters of monohydroxy fatty acids 329(1985)57
Banes, A.J., Link, G.W. and Snyder, L.R.
Comparison of reversed-phase columns for the separation of tryptic peptides by gradient elution. Correlation of experimental results and model prediction 326(1985)419
Banes, A.J., see Link, G.W. Jr. 331(1985)253
Barbaro, A.M., Guerra, M.C., Biagi, G.L., Pietrogrande, M.C. and Borea, P.A.
Influence of the mobile phase composition

on the reversed-phase TLC behaviour of a series of prostaglandins. Comparison of the extrapolated R_M values 347(1985)209
Barberan, F.A.T., Tomas, F., Hernandez, L. and Ferreres, F.
Reversed-phase HPLC of 5-hydroxyflavones bearing tri- or tetrasubstituted A rings 347(1985)443
Bardalaye, P.C. and Wheeler, W.B.
Simplified method for the clean-up and reversed-phase HPLC determination of benomyl in mangoes 330(1985)403
Bargnoux, H., see Bougerolle, A.M. 345(1985)59
Barkley, R.M., see Sievers, R.E. 349(1985)395
Barkley, R.M., see Wenclawiak, B. 349(1985)469
Barla, F., see Simon, J. 329(1985)389
Barnes, W.N., Ray, A. and Bates, L.J.
Reversed-phase HPLC method for the assay of oxytetracycline 347(1985)173
Barra, D., see Simmaco, M. 349(1985)99
Barreteau, H., see Brouard, A. 345(1985)453
Barreteau, H., see Paillet, M. 343(1985)455
Barrington, K.J., see Torok-Both, G.A. 344(1985)372
Barsett, H. and Paulsen, B.S.
Separation of acidic polysaccharides from *Ulmus glabra* Huds. on Mono P™ 329(1985)315
Barter, P.J., see Ha, Y.C. 341(1985)154
Basili, A., see Gagliardi, L. 348(1985)321
Bassene, E., Laurance, A., Olschwang, D. and Pousset, J.L.
Medicinal African plants. XIX. HPLC determination of vitexin in a crude extract of *Combretum micranthum* G. Don (in French) 346(1985)428
Basta, M.T., see Brooks, J.B. 339(1985)243
Bates, L.J., see Barnes, W.N. 347(1985)173
Battista, H.J., Wehinger, G. and Henn, R.
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77
Baudisch, H., see Lemm, U. 342(1985)393
Baumes, R.L., see Gunata, Y.Z. 331(1985)83
Bayer, E., see Albert, K. 346(1985)17
Bayne, W.F., see Constanzer, M.L. 339(1985)414
Bayne, W.F., see Musson, D.G. 338(1985)357
Bayne, W.F., see Slobodzian, D.K. 338(1985)253, 345(1985)345
Bayne, W.F., see Vincek, W.C. 338(1985)438
Bayonove, C.L., see Gunata, Y.Z. 331(1985)83
Beck, O. and Ryman, T.
Quantification of nitrendipine in plasma by a capillary column GC-MS methods 337(1985)402
Beck, R.H.F., see Praznik, W. 348(1985)187
Becker, C.R., Efcavitch, J.W., Heiner, C.R. and Kaiser, N.F.
Use of a C_4 column for reversed-phase HPLC purification of synthetic oligonucleotides 326(1985)293
Beckman, A.L., see Salzman, S.K. 343(1985)285
Bedford, G.K. and Chiong, M.A.
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides. Reply to E. Harmsen et al 337(1985)203
Bedford, K.R. and White, P.C.
Improved method for the simultaneous determination of morphine, codeine and dihydrocodeine in blood by HPLC with electrochemical detection 347(1985)398
Beecher, G.R., see Khachik, F. 346(1985)237
Beers, M.F., Stern, M., Hurtig, H., Melvin, G. and Scarpa, A.
Simultaneous determination of L-dopa and 3-O-methyldopa in human serum by HPLC 336(1985)380
Beger, J., Merkwitz, H., Szymanowski, J. and Voelkel, A.
Polarity of polyoxyethylene glycol dialkyl ethers and some sulphur analogues measured by GC 333(1985)319
Beger, J., see Szymanowski, J. 330(1985)61
Begert, A., see Boos, R. 328(1985)233
Beijersbergen van Henegouwen, R., see Boeren, E.G. 349(1985)377
Belle, S.J.P. van, see De Smet, M. 345(1985)309
Belle, S.J.P. van, see Van Belle, S.J.P. 337(1985)73
Bendre, S.B., Kadam, A.N. and Ghatge, B.B.
Comparative GC studies of bonded and physically coated PEG 20M and its derivative phases with reference to some high-boiling isomeric solutes 346(1985)359
Benedetti, M. Strolin, see Vajta, S. 343(1985)239
Benfenati, E., see Farina, P. 345(1985)323
Benjamin, L.E., see Fell, D. 345(1985)150
Bennett, H.P.J., see James, S. 326(1985)329
Benovitz, D.E., see Spatola, A.F. 327(1985)165
Benson, B., see Gregory, V.M. 345(1985)140
Bentley, D.P., see Littlewood, T.J. 336(1985)434
Benziger, D.P., see Brown, R.R. 339(1985)440
Berezkin, V.G and Retunsky, V.N.
Retention indices as a tool for the prediction of chromatographic conditions for achieving the separation of mixtures of organic compounds 330(1985)71
Berezkin, V.G., see Andronikashvili, T.G. 331(1985)402
Berezkin, V.G., see Itsikson, L.B. 334(1985)1
Berg, A.J.J. van den, see Van den Berg, A.J.J. 329(1985)311
Berg, G.A. van den, see Van den Berg, G.A. 339(1985)223
Bergamini, C., see Chiavari, G. 346(1985)369
Berger, J.A., see Bougerolle, A.M. 345(1985)59
Berman, S.S., see Siu, K.W.M. 330(1985)87
Bermejo, J., Blanco, C.G. and Guillén, M.D.
Capillary GC of chloro derivatives of 1,4-dimethylbenzene. Separation, identification and prediction of boiling points 331(1985)237
Bernard, N., see Julien, C. 344(1985)51
Bernard, P., see Sabot, J.F. 339(1985)233
Berndt, H. and Krüger, G.
Resolution of enantiomeric amino acid

derivatives by HPLC on chiral stationary phases 348(1985)275
Bernier, J.L., see Lhermitte, M. 342(1985)416
Bernstein, J.R. and Franklin, R.B.
HPLC assay of 2-[2-methoxy-4-(methylsulfinyl)phenyl]-1H-imidazo-[4,5-c]pyridine hydrochloride, a potent cardiotonic agent, and two metabolites in plasma 342(1985)228
Berry, H.K., see Norman, E.J. 337(1985)21
Berthet, D., see Necciari, J. 341(1985)202
Bertrand, O., Cochet, S., Kroviarski, Y., Truskolaski, A. and Boivin, P.
Protein precipitation induced by a textile dye. Precipitation of human plasminogen in the presence of Procion Red HE3B 346(1985)111
Bertsch, W., see Brill, J.H. 349(1985)31, 349(1985)39
Bertucci, C., see Rosini, C. 348(1985)79
Besch, P.K., see Dunaway, H.E. 327(1985)221
Besch, P.K., see Hutchens, T.W. 327(1985)247
Betina, V.
TLC of mycotoxins (Review) 334(1985)211
Betti, A., Coppi, S. and Bighi, C.
Pre-concentration of organic pollutants. Potential interference from the use of styrene copolymer adsorbents 349(1985)181
Betti, A., see Coppi, S. 330(1985)55
Beukhof, J.R., see Koopman, B.J. 337(1985)259
Beuzard, Y., see Mahieu, J.-P. 327(1985)313
Bhargava, V.O., Soine, W.H. and Garrettson, L.K.
HPLC analysis of 1-(β-D-glucopyranosyl)phenobarbital in urine 343(1985)219
Bhat, B.G. and Chandrasekhara, N.
Determination of piperine in biological tissues by TLC and absorption densitometry 338(1985)259
Bhattacharyya, P., see Chowdhury, B.K. 329(1985)178
Bhattacharyya, P.K., see Grant, A. 347(1985)219
Biagi, G.L., see Barbaro, A.M. 347(1985)209
Bialer, M. and Hoch, B.
Rapid GC assay for monitoring valnoctamide in plasma 337(1985)408
Bialer, M., see Yagen, B. 343(1985)67
Bianchi, G. and Landi, M.
Determination of apomorphine in rat plasma and brain by HPLC with electrochemical detection 338(1985)230
Bianchini, J.-P., Gaydou, E.M., Sigoillot, J.-C. and Terrom, G.
Determination of sterol and triterpene alcohol acetates in natural products by reversed-phase LC and GC-MS 329(1985)231
Biancini, G., see Del Principe, D. 342(1985)285
Bibby, M.C., see McDermott, B.J. 338(1985)335
Bicchi, C., D'Amato, A. and Cappelletti, E.
Determination of pyrrolizidine alkaloids in *Senecio inaequidens* D.C. by capillary GC 349(1985)23
Bidleman, T.F., see Foreman, W.T. 330(1985)203
Bidlingmeyer, B., see Colgan, S.T. 333(1985)349
Bidlingmeyer, B.A., Cohen, S.A and Tarvin, T.L.
Rapid analysis of amino acids using pre-column derivatization 336(1985)93
Bietz, J.A., see Huebner, F.R. 327(1985)333
Biggs, W.R., see Fetzer, J.C. 346(1985)81
Bighi, C., see Betti, A. 349(1985)181
Bighi, C., see Pietrogrande, M.C. 349(1985)63
Bigley, F.P. and Grob, R.L.
Determination of phenols in water and wastewater by post-column reaction detection HPLC 350(1985)407
Bihari, M., see Juvancz, Z. 337(1985)121
Binder, S.R. and Sivorinovsky, G.
Measurement of urinary vanilmandelic acid and homovanillic acid by HPLC with electrochemical detection following extraction by ion-exchange and ion-moderated partition 336(1985)173
Binkhuyzen, D., see Thieme, R.E. 337(1985)443
Bircher, J., see Baktir, G. 339(1985)192
Bissé, E. and Abraham, E.C.
New less temperature-sensitive microchromatographic method for the separation and quantitation of glycosylated hemoglobins using a non-cyanide buffer system 344(1985)81
Bitsch, I., see Schulz, A. 328(1985)417
Björkman, S.
Determination of the enantiomers of indoprofen in blood plasma by HPLC after rapid derivatization by means of ethyl chloroformate 339(1985)339
Blair, N., see Selan, F. 338(1985)213
Blair, R., see Misir, R. 331(1985)141
Blaise, P., see Staveris, S. 339(1985)97
Blanch, H.W., see Arnold, F.H. 330(1985)159
Blanco, C.G., see Bermejo, J. 331(1985)237
Blatchford, C., Humphreys, E. and Malcolme-Lawes, D.J.
Electrochemiluminescence as a detection technique for reversed-phase HPLC. II. Low-frequency a.c. electrochemiluminescence 329(1985)281
Blilie, A.L. and Greibrokk, T.
Gradient programming and combined gradient-pressure programming in supercritical fluid chromatography 349(1985)317
Blitterswijk, H. van, see Jansen, E.H.J.M. 347(1985)375, 347(1985)379
Blo, G., see Pietrogrande, M.C. 349(1985)63
Blom, Y., Bondesson, U. and Änggård, E.
Analysis of buprenorphine and its N-dealkylated metabolite in plasma and urine by selected-ion monitoring 338(1985)89
Blondin, A., see Le Hoang, M.D. 346(1985)382
Blum, W. and Grob, K.
Preparation of glass and fused-silica capillary columns coated with immobilized XE-60 346(1985)341

Blumenthal, H.P., see Brooks, M.A. 337(1985)351
Bo', L. dal, see Ascalone, V. 344(1985)231
Boček, P., Gebauer, P., Dolník, V. and Foret, F.
Recent developments in isotachophoresis (Review) 334(1985)157
Boček, P., see Fanali, S. 330(1985)436
Boeren, E.G., Beijersbergen van Henegouwen, R., Bos, I. and Gerner, T.H.
Naphtha analysis. The advantages of a specific olefin trap 349(1985)377
Bogusz, M., Gierz, J., De Zeeuw, R.A. and Franke, J.P.
Influence of the biological matrix on retention behaviour in TLC: evidence of systematic differences between pure and extracted drugs 342(1985)241
Bohov, P., see Krupčík, J. 346(1985)33
Boissel, J.-P., see Riffe, A. 344(1985)332
Boivin, P., see Bertrand, O. 346(1985)111
Bolívar, F., see Ladrón de Guevara, O. 349(1985)91
Bompeix, G., see Saindrenan, P. 347(1985)267
Bonakdar, M., see Wang, J. 344(1985)412
Bondesson, U., see Blom, Y. 338(1985)89
Bondesson, U., see Sjöberg, P. 344(1985)167
Bonn, G.
HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381
Boos, K.-S., see Kemper, K. 336(1985)374
Boos, R., Prey, T. and Begert, A.
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233
Boothe, T., Emran, A.M., Finn, R.D., Kothari, P.J. and Vora, M.M.
Chromatography of radiolabelled anions using reversed-phase LC columns 333(1985)269
Borea, P.A., see Barbaro, A.M. 347(1985)209
Borén, H., Grimvall, A., Palmborg, J., Sävenhed, R. and Wigilius, B.
Optimization of the open stripping system for the analysis of trace organics in water 348(1985)67
Borén, H., see Sävenhed, R. 328(1985)219
Borgia, R., see Pucci, P. 331(1985)425
Borra, C., see Andreolini, F. 337(1985)379
Bory, C., see Boulieu, R. 339(1985)380
Bos, I., see Boeren, E.G. 349(1985)377
Bosin, T.R. and Jarvis, C.A.
Derivatization in aqueous solution, isolation and separation of tetrahydro-β-carbolines and their precursors by LC 341(1985)287
Bosisio, A.B., Rochette, J., Wajcman, H., Gianazza, E. and Righetti, P.G.
Electrophoretic and chromatographic techniques for the differential diagnosis of a haemoglobin abnormality: Hb E heterozygosity 330(1985)299
Bosman, J., see Feitsma, K.G. 333(1985)59
Bossa, F., see Simmaco, M. 349(1985)99
Bossard, A., Kamga, R. and Raulin, F.
Quantitative GC analysis of low-molecular-weight alkylphosphines in the presence of phosphine and hydrocarbons 330(1985)400
Bossche, W. van den, see De Croo, F. 329(1985)422, 349(1985)301
Bossche, W. van den, see De Schutter, J.A. 350(1985)135
Bossle, P.C., Hallowell, S.F., Reutter, D.J. and Sarver, E.W.
Analysis of 2,2'-thiodiethanol in aqueous matrices by LC with electrochemical detection 330(1985)388
Both-Miedema, R., see Jansen, E.H.J.M. 347(1985)375, 347(1985)379
Botha, E., see Labadarios, D. 339(1985)366
Botha, S.A., see Du Preez, J.L. 333(1985)249
Bouchon, B. and Portoukalian, J.
Purification of the peracetylated glycosphingolipids of the Gala series (galactosyl- and galabiosylceramides) 342(1985)385
Bougerolle, A.M., Chabard, J.L., Bargnoux, H., Petit, J., Berger, J.A. and Dordain, G.
Simultaneous determination of metapramine and its dimethylated metabolites in plasma by GC-MS 345(1985)59
Boukouvalas, J., see Jefford, C.W. 347(1985)183
Boulieu, R., Bory, C. and Gonnet, C.
HPLC method for the analysis of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in biological fluids 339(1985)380
Bourgeois, G.L., see Demotes-Mainard, F.M. 342(1985)234
Bousquet, E., Romeo, G. and Giannola, L.I.
HPLC with fluorimetric detection in biological tissues of the 4-bromomethyl-7-methoxycoumarin ester derivative of 5-pyrrolidone-2-carboxylic acid 344(1985)325
Bouvier, E.S.P., see Huang, J.-X. 330(1985)181
Bowden, C.L., see Javors, M.A. 336(1985)259
Bowers, L.D. and Mathews, S.E.
Investigation of the mechanism of peak broadening observed in the HPLC analysis of cyclosporine 333(1985)231
Bowers-Komro, D.M., see Chastain, J.L. 330(1985)153
Boylan, J.M. and Susa, J.B.
Determination of adult sheep plasma catecholamines using [^{3}H]norepinephrine as the internal standard 338(1985)192
Boyle, D.M., Wiehle, R.D., Shahabi, N.A. and Wittliff, J.L.
Rapid, high-resolution procedure for assessment of estrogen receptor heterogeneity in clinical samples 327(1985)369
Brandl, M., see Knoll, R. 345(1985)425
Brandt, D.W., see Hyder, S.M. 327(1985)237
Brandts, P.M., Gelsema, W.J. and De Ligny, C.L.
Influence of additives to the eluent on hydrophobic interaction chromatography of

simple compounds. II. Influence of organic co-solvents on retention at 25°C 333(1985)41
Brantner, H., see Gamerith, G. 328(1985)241
Brassinne, C., Laduron, C., Sculier, J.P. and Coune, A.
HPLC determination of 2-[3'-(methoxycarbonylamino)-phenyl]-3-phenyl-6-methoxycarbonylamino-4-(3H)-quinazolone (NSC-251635) in human serum 339(1985)445
Braumann, T. and Jastorff, B.
Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. I. Cation binding in the mobile phase 329(1985)321
Braumann, T. and Jastorff, B.
Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. II. Quantitative determination of hydrophobicity 350(1985)105
Brauw, M.C. ten Noever de, see Van der Greef, J. 343(1985)397
Brazier, J.L., see Désage, M. 336(1985)285
Breckenridge, A.M., see Edwards, G. 343(1985)224
Breckenridge, A.M., see Mihaly, G.W. 337(1985)166
Bredesen, J.E., see Langseth-Manrique, K. 349(1985)421
Breese, G.R., see Patrick, K.S. 343(1985)329
Breitenbach, S., see Elbert, W. 328(1985)111
Brendel, E., Zschunke, M. and Meineke, I.
HPLC determination of cefonicid in human plasma and urine 339(1985)359
Brennan, S., see Keller, R.K. 343(1985)413
Bres, J., see Bressolle, F. 341(1985)391, 343(1985)443
Breslin, F. and Stephenson, G.P.
Determination of glycerol in pharmaceutical preparations by reversed-phase HPLC using a refractive index detector 329(1985)434
Bressolle, F. and Bres, J.
Quantitative analysis of sulpiride and sultopride by HPLC for pharmacokinetic studies (in French) 341(1985)391
Bressolle, F., Bres, J. and Snoussi, M.
HPLC determination of a new benzamide in biological fluids for use in a human pharmacokinetic study 343(1985)443
Březina, M., see Kříž, J. 347(1985)297
Briggs, M.H., see Thomson, S.B. 338(1985)201
Brill, J.H., Mar, T., Mayfield, H.T. and Bertsch, W.
Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. II. Comparison of the cuticular hydrocarbon patterns between different colonies of *Solenopsis richteri* 349(1985)39
Brill, J.H., Mayfield, H.T., Mar, T. and Bertsch, W.
Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. I. Introduction and characterization of the cuticular hydrocarbon patterns of *Solenopsis invicta* and *S. richteri* 349(1985)31
Brinkman, U.A.T. and Kamminga, D.
Rapid separation of enantiomers by TLC on a chiral stationary phase 330(1985)375
Brinkman, U.A.T., see Maris, F.A. 328(1985)93, 347(1985)75
Brinkman, U.A.T., see Nielen, M.W.F. 330(1985)113
Briza, P., see Winkler, G. 347(1985)83
Brodie, R.R., see Walmsley, L.M. 338(1985)433
Brogard, J.M., see Jehl, F. 339(1985)347
Brondz, I. and Olsen, I.
Differentiation between major species of the *Actinobacillus-Haemophilis-Pasteurella* group by GC of trifluoroacetic acid anhydride derivatives from whole-cell methanolysates 342(1985)13
Brondz, I. and Olsen, I.
Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 345(1985)119
Brooks, C.J.W., Watson, D.G. and Cole, W.J.
Analytical separation of enantiomeric polygodials by GC of pyrrole derivatives 347(1985)455
Brooks, J.B., Basta, M.T., Holler, J.S., Alley, C.C. and Kholy, A.M.
Frequency-pulsed electron-capture GLC studies of chemical changes in sera of patients with schistosomiasis 339(1985)243
Brooks, M.A., DiDonato, G. and Blumenthal, H.P.
Determination of chlorprothixene and its sulfoxide metabolite in plasma by HPLC with UV and amperometric detection 337(1985)351
Brooks, M.A. and Lee, T.L.
Determination of 4-amino-5-ethyl-3-thiophenecarboxylic acid methyl ester and its acid metabolite in plasma and urine by HPLC 345(1985)333
Brosch, D., see Kříž, J. 347(1985)297
Brosche, T., Platt, D. and Vostrowsky, O.
Methyl enol ethers as artefacts in capillary GC profiles of aldehyde dimethyl acetals 345(1985)219
Brotherton, J.E. and Widholm, J.M.
HPLC of anthranilate synthase using gel filtration and a post-column reactor 350(1985)332
Brouard, A., Barreteau, H., Merdjan, H., Paillet, M., Fredj, G. and Micoud, M.
Rapid determination of pyrazinamide in biological fluids by HPLC 345(1985)453
Brouard, A., see Paillet, M. 343(1985)455
Brown, A.S., Cho, K.Y., Cheung, H.T.A., Hemmens, V. and Vine, J.
Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary GC-MS 341(1985)139
Brown, K.F., see Tett, S.E. 344(1985)241
Brown, M.J., see Causon, R.C. 337(1985)311

Brown, N.D., Poon, B.T. and Chulay, J.D.
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 345(1985)209
Brown, P.R. and Fasching, J.L.
Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 345(1985)457
Brown, P.R., see Dwyer, M.E. 345(1985)125
Brown, P.R., see Halfpenny, A.P. 349(1985)275
Brown, P.R., see Yi, C. 339(1985)75
Brown, R.R., Storshane, R.M. and Benziger, D.P.
HPLC assay for trilostane and its major metabolite, 17-ketotrilostane, in human plasma 339(1985)440
Browne, C.A., Sim, F.R., Rae, I.D. and Keeler, R.F.
Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211
Brueggemann, E.E., see Burrows, E.P. 329(1985)285
Bruneau, P., see Meinard, C. 349(1985)105, 349(1985)109
Bryce, T.A., Burrows, J.L. and Jolley, K.W.
Determination of 1-(3'-carboxypropyl)-3,7-dimethylxanthine and 1-(4'-carboxybutyl)-3,7-dimethylxanthine, two major metabolites of oxpentifylline, in urine by HPLC 344(1985)397
Buchanan, D.N. and Thoene, J.G.
Photodiode array UV spectrophotometric profiling of carboxylic acids in physiological fluids 344(1985)23
Bucheli, F., see Dell, D. 344(1985)125
Buda, W.M., see Suprynowicz, Z. 333(1985)11
Buijten, J., see Neelakantan, L. 350(1985)307
Bullen, W.W., see Gammans, R.E. 339(1985)303, 345(1985)285
Bun, H., see Tomasini, J.-L. 343(1985)369
Bur, H., see Milon, H. 350(1985)399
Burford, R.P., Haken, J.K. and Vimalasiri, P.A.D.T.
GC analysis of isocyanate-based copolyamide resins 329(1985)132
Burford, R.P., see Haken, J.K. 349(1985)347
Burger, J.A., see Moodie, I.M. 347(1985)179
Burke, J.T. and Thénot, J.P.
Determination of antiepileptic drugs (Review) 340(1985)199
Burns, D.T., see Adair, C.G. 336(1985)429, 342(1985)447
Burrows, E.P. and Brueggemann, E.E.
Reversed-phase gradient HPLC of nitramine munitions and characterization of munitions process samples by GC-MS 329(1985)285
Burrows, J.L. and Jolley, K.W.
Determination of oxpentifylline and four metabolites in plasma and urine by automated capillary GC using nitrogen-selective detection 344(1985)187
Burrows, J.L., see Bryce, T.A. 344(1985)397
Bursey, M.M., see Parker, C.E. 347(1985)61
Burton, F.W. and Gadde, R.R.
Analysis of anthralin in dermatological products by reversed-phase HPLC 328(1985)317
Burton, J., see Soparkar, C.N.S. 337(1985)110
Bury, R.W.
LC assay of isoxicam in human plasma and urine 337(1985)156
Bury, R.W., see Kacprowicz, A.T. 337(1985)412
Büser, W. and Erbersdobler, H.F.
Determination of furosine by GLC 346(1985)363
Bush, C.A., see Dua, V.K. 328(1985)259
Busker, E., Günther, K. and Martens, J.
Application of chromatographic chiral stationary phases to pharmaceutical analysis. Enantiomeric purity of D-penicillamine 350(1985)179
Bussolo, J.M. di, see Di Bussolo, J.M. 327(1985)67
Buzello, W., see Williams, C.H. 344(1985)71

C

Cabezudo, M.D., see Reglero, G. 348(1985)327
Caccia, S., see Fong, M.H. 333(1985)191
Cachia, P.J., Van Eyk, J., McCubbin, W.D., Kay, C.M. and Hodges, R.S.
Ion-exchange HPLC purification of bovine cardiac and rabbit skeletal muscle troponin subunits 343(1985)315
Caddy, B., see Chow, W.M.L. 329(1985)213
Cafmeyer, N., see Keller, R.K. 343(1985)413
Cahill, C.M., see Harrison, P.M. 343(1985)349
Calam, D.H., Davidson, J. and Harris, R.
HPLC investigations on some enzymes of papaya latex 326(1985)103
Caldwell, R. and Flanagan, R.J.
Measurement of trazodone in biological fluids by HPLC 345(1985)187
Callewaert, K., see Daenens, P. 342(1985)79
Campos, E.M., see McConathy, W.J. 342(1985)47
Canal, M. and Flouvat, B.
Determination of betaxolol in plasma by HPLC with fluorescence detection 342(1985)212
Cannata, A., see Nota, G. 347(1985)362
Cano, J.-P., see Coassolo, P. 338(1985)347
Cano, J.-P., see Tomasini, J.-L. 343(1985)369
Cappelletti, E., see Bicchi, C. 349(1985)23
Carducci, C.N., Matejka, M., Ergüven, H., Labonia, N. and Mamianetti, A.
HPLC analysis of bile acids in hamster bile 337(1985)91
Carey, P.F., see Colthup, P.V. 345(1985)111
Carlqvist, J. and Westerlund, D.
Automated determination of amoxycillin in

biological fluids by column switching in ion-pair reversed-phase LC systems with post-column derivatization 344(1985)285
Carlsson, A. and Lundström, K.
Amperometric detector for use with small-bore LC columns 350(1985)169
Carlsson, M., see Andersson, T. 326(1985)33
Carmi, J.D., see Santa María, I. 329(1985)123
Carrel, J.E., Doom, J.P. and McCormick, J.P.
Quantitative determination of cantharidin in biological materials using capillary GC with flame ionization detection 342(1985)411
Cartoni, G., see Ronchetti, M. 348(1985)159
Cartoni, G.P., Ciardi, M., Giarrusso, A. and Rosati, F.
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315
Casanova, M.G., see Gagliardi, L. 348(1985)321
Castagnetta, L., see D'Agostino, G. 338(1985)1
Castello, G. and D'Amato, G.
Effect of the mesh-size distribution of porous polymer beads on efficiency and selectivity of stationary phases 349(1985)189
Castells, R.C.
Possibility and consequences of using different concentration scales in the study of solution thermodynamics by GLC 350(1985)339
Castiglioni, M., Giordano, R., Sappa, E. and Volpe, P.
Catalytic hydrogenation and exchange reactions on GC columns, in the presence of supported $(\eta^5\text{-}C_5H_5)NiOs_3(\mu\text{-}H)_3(CO)_9$ 349(1985)173
Castoldi, D., Monzani, V. and Tofanetti, O.
Determination of bezafibrate in human plasma and urine by HPLC 344(1985)259
Castro-Parra, M., see Von Sassen, W. 338(1985)113
Catalani, P., see Ascalone, V. 344(1985)231
Cataldi, T.R.I., see Palmisano, F. 344(1985)249
Catlow, D.A., Johnson, M., Monaghan, J.J., Porter, C. and Scrivens, J.H.
Some results from an investigation into the analytical potential of MS-MS techniques, with particular reference to industrial applications 328(1985)167
Causon, R.C., Brown, M.J. and Davies, D.S.
Reversed-phase HPLC and amperometric detection of 3-O-methyl isoprenaline sulphate: application to studies on the presystemic metabolism of *d*-isoprenaline in man 337(1985)311
Cautreels, W., see Davi, H. 342(1985)159
Cautreels, W., see Necciari, J. 341(1985)202
Cavazzutti, G., see Gagliardi, L. 348(1985)321
Cecchini, D.J., see Ehrat, M. 326(1985)311
Cee, A. and Horáková, B.
Detection of citrazinic acid by PC and TLC 331(1985)202
Ćelap, M.B., see Janjić, T.J. 331(1985)273
Čellár, P., see Matisová, E. 346(1985)177
Cerulli, N., see Laganà, A. 345(1985)251
Chabard, J.L., see Bougerolle, A.M. 345(1985)59
Chaiken, I.M., Kanmera, T., Sequeira, R.P. and Swaisgood, H.E.
HPLC and studies of neurophysin-neurohypophysial hormone pathways 336(1985)63
Chaiken, I.M., see Swaisgood, H.E. 327(1985)193
Chalmers, J.L., see Meatherall, R.C. 338(1985)141
Chamart, S., see Duez, P. 329(1985)415
Chamblee, T.S., Clark, B.C. Jr., Radford, T. and Iacobucci, G.A.
General method for the HPLC prefractionation of essential oils and flavor mixtures for GC-MS analysis. Identification of new constituents in cold pressed lime oil 330(1985)141
Chan, B.G., see Mahoney, N.E. 329(1985)91
Chan, K.
Simultaneous determination of carbamazepine and its epoxide metabolite in plasma and urine by HPLC 342(1985)341
Chan, P.H., see Chen, S.F. 344(1985)297
Chandra, D.P., see So, N. 337(1985)81
Chandrasekhara, N., see Bhat, B.G. 338(1985)259
Chang, C.A., Abdel-Aziz, H., Melchor, N., Wu, Q., Pannell, K.H. and Armstrong, D.W.
LC retention behavior of organometallic compounds and ligands with amine-, octadecylsilica- and β-cyclodextrin-bonded phase columns 347(1985)51
Chang, S.F., Welscher, T.M., Miller, A.M., McQuinn, R.L. and Fox, J.M.
HPLC method for the quantitation of a *meta*-O-dealkylated metabolite of flecainide acetate, a new antiarrhythmic 343(1985)119
Chang, T., see Hayes, A.G. 336(1985)446
Chang, T.-M., Erway, B. and Chey, W.Y.
Rapid, small-scale preparation of gastrointestinal hormones by HPLC on a C_{18} column 326(1985)121
Charlton, J.P.
Combined recycling affinity column-filtration technique 346(1985)247
Chasseaud, L.F., see Taylor, I.W. 343(1985)427
Chasseaud, L.F., see Walmsley, L.M. 338(1985)433
Chastain, J.L., Bowers-Komro, D.M. and McCormick, D.B.
HPLC of biotin and analogues 330(1985)153
Chavko, M., see Kehr, J. 345(1985)267
Cheetham, N.W.H. and Teng, G.
Some applications of reversed-phase HPLC to oligosaccharide separations 336(1985)161
Chen, S.F. and Chan, P.H.
One-step separation of free fatty acids and phospholipids in brain tissue extracts by HPLC 344(1985)297
Cheng, L.K., see Osman, M.A. 336(1985)329, 345(1985)430
Chenieux, J.C., see Montagu, M. 331(1985)437
Cheung, H.T.A., see Brown, A.S. 341(1985)139
Chey, W.Y., see Chang, T.-M. 326(1985)121

Chiavari, G. and Bergamini, C.
Determination of phenylurea herbicides by HPLC with electrochemical detection 346(1985)369
Chiavari, G., Facchini, M.C. and Fuzzi, S.
Determination of formaldehyde as its lutidine derivative in the atmospheric liquid phase by HPLC 333(1985)262
Chicarelli, M.I., see Gill, J.P. 350(1985)37
Chikai, T., see Goto, J. 348(1985)151
Chikuma, T., Ishii, Y. and Kato, T.
Highly sensitive assay for PZ-peptidase activity by HPLC 348(1985)205
Chimenti, F., see Gagliardi, L. 348(1985)321
Chino, N., see Kumagaye, K.Y. 327(1985)327
Chiong, M.A., see Bedford, G.K. 337(1985)203
Chiu, G.P., see Cwik, M.J. 338(1985)123
Chmela, Z., see Stránský, Z. 342(1985)167
Chmiel, J., see Matysik, J. 333(1985)198
Cho, K.Y., see Brown, A.S. 341(1985)139
Cho, S.H., see Smith, L.A. 346(1985)291
Choi, K.E., Sinkule, J.A., Crom, W.R., Thompson, E.I. and Evans, W.E.
HPLC assay of mitomycin in biological fluids 345(1985)197
Cholerton, T.J., Hunt, J.H. and Martin-Smith, M.
Convenient GC method for the determination of the proportions of the two racemic modifications present in labetalol hydrochloride 333(1985)178
Choporis, N., see Kotake, A.N. 337(1985)194
Chou, P.P. and Jaynes, P.K.
Determination of urinary 5-hydroxyindole-3-acetic acid using solid-phase extraction and reversed-phase HPLC with electrochemical detection 341(1985)167
Chovan, J.P., see Klett, R.P. 337(1985)182
Chow, W.M.L. and Caddy, B.
Improved method for the preparation of immobilised pyrimidonesilicones as polar stationary phases used for the capillary column GC separation of underivatised barbiturates 329(1985)213
Chow-Tung, E., see Cwik, M.J. 338(1985)123
Chowdhury, B.K., Mustapha, A. and Bhattacharyya, P.
Separation of carbazole alkaloids by GLC 329(1985)178
Chrétien, M., see Lazure, C. 326(1985)339
Christ, D.D., see Walle, T. 341(1985)213
Christie, A.E.G., see Wells, D.E. 328(1985)372
Chulay, J.D., see Brown, N.D. 345(1985)209
Chung, A., see Lam, L.K.T. 328(1985)422
Chung, T.G., Saito, A., Ogawa, H., Moriyama, A., Ohkubo, I., Sasaki, M. and Takahashi, T.
Efficient purification of human plasma β_2-microglobulin from the haemodialysate of a patient with chronic renal failure by use of HPLC 339(1985)67
Churáček, J., see Horna, A. 330(1985)439, 348(1985)141
Churchill, F.C., see Dickinson, C.M. 345(1985)91
Ciaranello, R.D., see Speedie, M.K. 327(1985)351
Ciardi, M., see Cartoni, G.P. 330(1985)315
Cięszczyk-Chmiel, A., see Matysik, J. 333(1985)198
Cifuentes, M., see Santa María, I. 329(1985)123
Cillard, J., Gobaille, J. and Cillard, P.
Separation of three dimers of α-tocopherol by HPLC 347(1985)434
Cillard, J., see Torel, J. 330(1985)425
Cillard, P., see Cillard, J. 347(1985)434
Cillard, P., see Torel, J. 330(1985)425
Cingolani, L. and Croci, D.
Simple and rapid method for the quantitation of plasma phytanic acid by GC 339(1985)375
Cirma, G., see Musumarra, G. 350(1985)151
Ciszkowski, K., see Oszczapowicz, J. 330(1985)79
Clark, B.C. Jr., see Chamblee, T.S. 330(1985)141
Clark, T. and Deas, A.H.B.
Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181
Clarke, A.E., see Harris, P.J. 350(1985)304
Claudy, P., Letoffé, J.M., Gaget, C., Morel, D. and Serpinet, J.
Long-chain alkyl grafts and mixed alkyl-alkane layers at the surface of macroporous silicas. Their GC properties below and above the phase transition 329(1985)331
Claudy, P., see Gonnet, C. 330(1985)227
Clementi, S., see Musumarra, G. 350(1985)151
Clezardin, P., see McGregor, J.L. 326(1985)179
Clezy, P.S., see Duncan, M.W. 336(1985)199
Climax, J., see Dadgar, D. 337(1985)47, 337(1985)136, 342(1985)315
Clinch, R.M., see Colenutt, B.A. 346(1985)25
Cline Love, L.J., see Arunyanart, M. 342(1985)293
Cloix, J.F., see Wainer, I.W. 338(1985)417
Clouse, S.D., Martensen, A.N. and Gilchrist, D.G.
Rapid purification of host-specific pathotoxins from *Alternaria alternata* f. sp. *lycopersici* by solid-phase adsorption on octadecylsilane 350(1985)255
Coassolo, P., Aubert, C. and Cano, J.-P.
Plasma determination of 3-methylclonazepam by capillary GC 338(1985)347
Coassolo, P., see Tomasini, J.-L. 343(1985)369
Cobelli, L., see Poy, F. 349(1985)17
Cochet, S., see Bertrand, O. 346(1985)111
Coddens, M.E., see Dhanesar, S.C. 349(1985)249
Coggins, J.R., see Mousdale, D.M. 329(1985)268
Cohen, A.I., see Jemal, M. 342(1985)186, 345(1985)299
Cohen, B.E., see Egorin, M.J. 343(1985)196
Cohen, S.A, see Bidlingmeyer, B.A. 336(1985)93
Coldwell, R.D., Trafford, D.J.H., Makin, H.L.J., Varley, M.J. and Kirk, D.N.
Specific mass fragmentographic assay for 25,26-dihydroxyvitamin D in human plasma

using a deuterated internal standard 338(1985)289
Cole, D.E.C. and Landry, D.A.
Determination of inorganic sulfate in human saliva and sweat by controlled-flow anion chromatography. Normal values in adult humans 337(1985)267
Cole, W.J., see Brooks, C.J.W. 347(1985)455
Colenutt, B.A. and Clinch, R.M.
Chromatographic properties of oxidised OV-17 stationary phase 346(1985)25
Colgan, S.T., Krull, I.S., Neue, U., Newhart, A., Dorschel, C., Stacey, C. and Bidlingmeyer, B.
Derivatization of alkyl halides and epoxides with picric acid salts for improved HPLC detection 333(1985)349
Colistra, C., see Del Principe, D. 342(1985)285
Colombo, L., see Marcucci, F. 336(1985)356
Colombo, R., see Zecca, L. 339(1985)203
Colpan, M., see Hecker, R. 326(1985)251
Colthup, P.V., Dallas, F.A.A., Saynor, D.A., Carey, P.F., Skidmore, L.F., Martin, L.E. and Wilson, K.
Determination of salbutamol in human plasma and urine by HPTLC 345(1985)111
Colussi, D., Marfil, F. and Sioufi, A.
Simultaneous determination of pirprofen and its metabolite, the pyrrole derivative, in plasma by HPLC 339(1985)157
Condra, M., see Owen, J.A. 342(1985)333
Cone, E.J., Gorodetzky, C.W., Yousefnejad, D. and Darwin, W.D.
^{63}Ni electron-capture GC assay for buprenorphine and metabolites in human urine and feces 337(1985)291
Constanzer, M.L., Vincek, W.C. and Bayne, W.F.
Determination of cyclobenzaprine in plasma and urine using capillary GC with nitrogen-selective detection 339(1985)414
Constanzer, M.L., see Vincek, W.C. 338(1985)438
Conway, P.L., see Adams, R.F. 336(1985)125
Conzentino, P., see Strojny, N. 342(1985)145
Cook, A.M., see Grossenbacher, H. 331(1985)161
Cook, L., Schafer-Mitchell, M., Angle, C. and Stohs, S.
Assay of human erythrocyte pyrimidine and deoxypyrimidine 5'-nucleotidase by isocratic reversed-phase HPLC 339(1985)293
Cookman, J.R., see Hartley, R. 342(1985)105
Coombe, R.G., Vine, J.H. and Yip, H.
Unusual fatty acids from amniotic fluid phospholipids 341(1985)146
Cooper, E.H., Johns, E.A., Itoh, Y. and Webb, J.R.
Chromatographic study of the interrelationships of immunoglobulin A and α_1-microglobulin in myelomatosis 327(1985)179
Cooper, E.H., Turner, R., Webb, J.R., Lindblom, H. and Fagerstam, L.
Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269
Cooper, H.A. and Hurtubise, R.J.
Solubility parameter and window diagram optimization methods for the reversed-phase chromatographic optimization of complex hydroxyl aromatic mixtures 328(1985)81
Cooper, S.D., see Pellizzari, E.D. 334(1985)277
Cooper, T.B., see Suckow, R.F. 338(1985)225
Coppi, S. and Betti, A.
Correlation between chemical and physical characteristics and adsorption properties of some styrene copolymers 330(1985)55
Coppi, S., see Betti, A. 349(1985)181
Corcia, A. di, see Andreolini, F. 337(1985)379
Cordonnier, R.E., see Gunata, Y.Z. 331(1985)83
Corona, G.L., see Schinelli, S. 338(1985)396
Correia, L., see Cunico, R.L. 336(1985)105
Cortes, H.J., Richter, B.E., Pfeiffer, C.D. and Jensen, D.E.
Determination of trace chlorinated benzenes in fuel oil by on-line multidimensional chromatography using packed-capillary LC and capillary GC 349(1985)55
Costa, M., see Murphy, R. 336(1985)41
Coune, A., see Brassinne, C. 339(1985)445
Coutts, R.T., see Torok-Both, G.A. 344(1985)372
Covington, R.R., see Gammans, R.E. 339(1985)303
Cowan, A.A., see Wells, D.E. 328(1985)372
Cowlishaw, M.G. and Sharman, J.R.
Doxazosin determination by HPLC using fluorescence detection 344(1985)403
Cox, J.W., see Pullen, R.H. 343(1985)271
Crabbe, M.J.C., see Lloyd, P. 343(1985)402
Crabos, M., see Wainer, I.W. 338(1985)417
Craescu, C.T., see Mahieu, J.-P. 327(1985)313
Crapper McLachlan, D.R., see Kruck, T.P.A. 341(1985)123
Crawhall, J.C., see Kremer, R. 344(1985)313
Creer, M.H. and Gross, R.W.
Reversed-phase HPLC separation of molecular species of alkyl ether, vinyl ether, and monoacyl lyosphospholipids 338(1985)61
Crevat, A., see Gallice, P. 331(1985)445
Croci, D., see Cingolani, L. 339(1985)375
Crockard, A.D., see Adair, C.G. 336(1985)429, 342(1985)447
Crom, W.R., see Choi, K.E. 345(1985)197
Cronholm, T. and Norsten, C.
GC-MS of carboxylic acids in tissues as their *tert.* -butyldimethylsilyl derivatives 344(1985)1
Croo, F. de, see De Croo, F. 329(1985)422, 349(1985)301
Crooks, P.A., see Mousa, S. 347(1985)405
Csató, E., Keresztes, P., Szabó, L.D. and Knausz, D.
Preparation of trimethylsilyl derivatives of ribonucleosides for GC 348(1985)280
Cserháti, T., Szögyi, M. and Györfi, L.
Interaction of some non-ionic tensides with

dioleoyl phosphatidyl choline, studied by charge-transfer chromatography 349(1985)295
Cubellis, M.V., Marino, G., Mayol, L., Piccialli, G. and Sannia, G.
Use of fast protein LC for the purification of synthetic oligonucleotides 329(1985)406
Cucchi, M.L., see Schinelli, S. 338(1985)396
Cucchiarini, L., see Stocchi, V. 349(1985)77
Cuddy, K.K., see Skrabalak, D.S. 341(1985)261
Cuisinaud, G., Terrier, M., Ferry, N., Proust, S. and Sassard, J.
HPLC determination of cicletanide, a new diuretic, in plasma, red blood cells, urine and saliva 341(1985)97
Cuisinaud, G., see Julien, C. 344(1985)51
Culbertson, C.R., see Rodriguez, P.A. 330(1985)365
Cullen, M.P. and Baker, R.W.R.
Confidence limits for a gel-permeation system 341(1985)420
Cullen, M.P., Turner, C. and Haycock, G.B.
GPC with interference refractometry for the rapid assay of polydisperse dextrans in biological fluids 337(1985)29
Culling-Berglund, A., see Davis, T.P. 327(1985)279
Cummings, J.
Method for the determination of 4'-deoxydoxorubicin, 4'-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 341(1985)401
Cummings, J. and Willmott, N.
Adriamycin-loaded albumin microspheres: qualitative assessment of drug incorporation and in vitro release by HPLC and high-speed multi-diode array spectrophotometric detection 343(1985)208
Cunico, R.L., Simpson, R., Correia, L. and Wehr, C.T.
High-sensitivity phenylthiohydantoin amino acid analysis using conventional and microbore chromatography 336(1985)105
Currie, B.L., see Cwik, M.J. 338(1985)123
Curry, S.H., see Karnes, H.T. 338(1985)377
Cutler, D.J., see Tett, S.E. 344(1985)241
Cutlip, R.C., see Schmerr, M.J.F. 326(1985)225
Cwik, M.J., Chiu, G.P., Fischer, J.H., Chow-Tung, E. and Currie, B.L.
Quantitative determination of tolazoline in serum and urine 338(1985)123
Czerwiec, W., see Kuśmierz, J. 331(1985)219
Czira, G., see Szederkényi, F. 333(1985)404

D

Dadgar, D., Climax, J., Lambe, R. and Darragh, A.
GC determination of mefloquine in human and dog plasma using electron-capture detection 337(1985)47
Dadgar, D., Climax, J., Lambe, R. and Darragh, A.
GLC determination of phentermine in human plasma following oral administration to healthy subjects 337(1985)136
Dadgar, D., Climax, J., Lambe, R. and Darragh, A.
HPLC determination of certain salicylates and their major metabolites in plasma following topical administration of a liniment to healthy subjects 342(1985)315
Dadone, A., see Mazzucotelli, A. 349(1985)137
Daenens, P., Laruelle, L., Callewaert, K., De Schepper, P., Galeazzi, R. and Van Rossum, J.
Determination of cotinine in biological fluids by capillary GC-MS selected-ion monitoring 342(1985)79
D'Agostino, G., Castagnetta, L., Mitchell, F. and O'Hare, M.J.
Computer-aided mobile phase optimization and chromatogram simulation in HPLC (Review) 338(1985)1
D'Agostino, P.A., Hansen, A.S., Lockwood, P.A. and Provost, L.R.
Capillary column GC-MS of tabun 347(1985)257
D'Agostino, P.A. and Provost, L.R.
GC retention indices of chemical warfare agents and simulants 331(1985)47
Dal Bo', L., see Ascalone, V. 344(1985)231
Dalas, E., Karaiskakis, G., Katsanos, N.A. and Gounaris, A.
Reversed-flow GC technique applied to measurement of mass transfer coefficients of *n*-hydrocarbons on Porapak P 348(1985)339
Dalene, M. and Skarping, G.
Trace analysis of amines and isocyanates using glass capillary GC and selective detection. IV. Determination of free aromatic amines using nitrogen-selective detection 331(1985)321
Dalene, M., see Skarping, G. 331(1985)331, 346(1985)191
Dallas, F.A.A., see Colthup, P.V. 345(1985)111
D'Amato, A., see Bicchi, C. 349(1985)23
D'Amato, G., see Castello, G. 349(1985)189
Damm, P., see Lehr, K.-H. 339(1985)451
Danesh, B., see Tebbett, I.R. 329(1985)196
Darakis, G., see Saindrenan, P. 347(1985)267
Darbord, J.C., see Pons, J.-L. 337(1985)213
D'Arcangelo, C., see Del Principe, D. 342(1985)285
Darragh, A., see Dadgar, D. 337(1985)47, 337(1985)136, 342(1985)315
Darwin, W.D., see Cone, E.J. 337(1985)291
Davi, H., Guyonnet, J., Necciari, J. and Cautreels, W.
Determination of circulating ethyl loflazepate metabolites in the baboon by

radio-HPLC with injection of crude plasma samples: comparison with solvent extraction and TLC 342(1985)159
Davídek, J., see Hajšlová, J. 348(1985)437
Davidson, J., see Calam, D.H. 326(1985)103
Davies, C.L. and Molyneux, S.G.
Determination of diprobutine in human plasma using GLC with nitrogen-selective detection 339(1985)186
Davies, D.S., see Causon, R.C. 337(1985)311
Davies, D.S., see Murray, S. 337(1985)441
Davies, R.J., see Devalia, J.L. 343(1985)407
Davis, R.J.H., see Abdullah, M. 347(1985)129
Davis, T.P. and Culling-Berglund, A.
HPLC analysis of *in vitro* central neuropeptide processing 327(1985)279
Day, R.O., see Ray, J.E. 336(1985)234
De Abreu, R.A., see Van Baal, J.M. 336(1985)422
De Croo, F., Van den Bossche, W. and De Moerloose, P.
Simultaneous determination of althiazide and spironolactone in tablets by HPLC 329(1985)422
De Croo, F., Van den Bossche, W. and De Moerloose, P.
Influence of the chain length of chemically bonded phases on the behaviour of several thiazide, potassium-sparing and loop diuretics in HPLC 349(1985)301
De Deyn, P., see Marescau, B. 345(1985)215
De Guevara, O.L., Quintero, R. and Padilla, P.
Monitoring of the production of D-phenylglycine from DL-phenylhydantoin by HPLC 329(1985)428
De Jong, A.P.J.M., see Kok, R.M. 343(1985)59
De Jong, J.W., see Harmsen, E. 337(1985)201
De la Rosa, J., see Hirschberger, L.L. 343(1985)303
De Leenheer, A.P., see Nelis, H.J.C.F. 333(1985)381
De Ligny, C.L., see Brandts, P.M. 333(1985)41
De Moerloose, P., see De Croo, F. 329(1985)422, 349(1985)301
De Moerloose, P., see De Schutter, J.A. 350(1985)135
De Reu, L., see De Schryver, E.P. 338(1985)389
De Schepper, P., see Daenens, P. 342(1985)79
De Schryver, E.P., De Reu, L. and Willems, J.L.
Determination of methyl paraoxon in dog plasma by reversed-phase HPLC 338(1985)389
De Schutter, J.A., Van der Weken, G., Van den Bossche, W. and De Moerloose, P.
Reversed-phase HPLC of mebeverine hydrochloride and related compounds 350(1985)135
De Silva, J.A.F.
Analytical strategies for therapeutic monitoring of drugs in biological fluids (Review) 340(1985)3
De Silva, J.A.F., see Puglisi, C.V. 344(1985)199
De Silva, J.A.F., see Strojny, N. 336(1985)301, 341(1985)313, 342(1985)145
De Silva, R.L.
Elution and washing in fixed bed affinity chromatography 330(1985)356
De Smet, M., Van Belle, S.J.P., Storme, G.A. and Massart, D.L.
HPLC determination of vinca-alkaloids in plasma and urine 345(1985)309
De Smet, M., see Musch, G. 348(1985)97
De Tombe, P.P., see Harmsen, E. 337(1985)201
De Villiers, L.S., see Ubbink, J.B. 342(1985)277
De Vries, J.X., Simon, M., Zimmerman, R. and Harenberg, J.
Identification of phenprocoumon metabolites in human urine by HPLC and GC-MS 338(1985)325
De Zeeuw, R.A., see Bogusz, M. 342(1985)241
De Zeeuw, R.A., see Feitsma, K.G. 333(1985)59
DeAngelis, J.J., see Sievers, R.E. 349(1985)395
Deas, A.H.B., see Clark, T. 329(1985)181
Dębowski, J., Grassini-Strazza, G. and Sybilska, D.
α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131
Debowski, J., Jurczak, J., Sybilska, D. and Żukowski, J.
Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206
Debs, J., see Jehl, F. 339(1985)347
Dechaud, H., see Sabot, J.F. 339(1985)233
Dechavanne, M., see McGregor, J.L. 326(1985)179
Dedek, W., Grahl, R. and Mothes, B.
A simple TLC method for studies of radiolabelled pesticides on soil surfaces using soil-silica layers 331(1985)200
Del Principe, D., Colistra, C., Menichelli, A., Biancini, G., D'Arcangelo, C., Multari, G., Werner, B., Lemmo, M. and Digilio, G.
Isotachophoresis as a useful tool for monitoring neurological complications of acute leukaemia in children 342(1985)285
Dell, D., Wendt, G., Bucheli, F. and Trautmann, K.-H.
Determination of pyrrolidinone in plasma by HPLC 344(1985)125
Dellacherie, E., see Mathis, R. 347(1985)291
Delpech, M., see Riffe, A. 344(1985)332
DeLucas, L.J., see Muccio, D.D. 326(1985)243
Demidov, V.V., see Potaman, V.N. 329(1985)33
Demotes-Mainard, F.M., Vinçon, G.A., Jarry, C.H. and Albin, H.C.
Micromethod for the determination of cefotaxime and desacetylcefotaxime in plasma and urine by HPLC 336(1985)438
Demotes-Mainard, F.M., Vinçon, G.A., Jarry, C.H., Bourgeois, G.L. and Albin, H.C.
HPLC determination of apalcillin in plasma and urine 342(1985)234
Deng, P. and Andrawes, F.
Integration of the deformed peak from the helium ionization detector 349(1985)415
Deng, P., see Andrawes, F. 349(1985)405

DePace, A., see Verebey, K. 343(1985)339
Derendorf, H., see Yost, R.L. 341(1985)131
Dernick, R., see Heukeshoven, J. 326(1985)91
Deruaz, D., see Sabot, J.F. 339(1985)233
Désage, M., Soubeyrand, J., Soun, A., Brazier, J.L. and Georges, Y.
Automated theophylline assay using GC and a mass-selective detector 336(1985)285
Desai, S.V., see Sane, R.T. 331(1985)432
Desai, Z.R., see Adair, C.G. 336(1985)429
Deshmukh, B.K., see Wang, J. 344(1985)412
Desideri, P.G., see Lepri, L. 347(1985)303
DeStefano, J.J., see Stout, R.W. 326(1985)63
Deutscher, J., see Alpert, C.A. 326(1985)363
Devalia, J.L., Sheinman, B.D. and Davies, R.J.
Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407
Dew, K.L., see Kilts, C.D. 342(1985)452
Dewaele, C., see Verzele, M. 329(1985)351
Deyl, Z., see Schwarz, V. 340(1985)401
Deyn, P. de, see Marescau, B. 345(1985)215
Dhanesar, S.C., Coddens, M.E. and Poole, C.F.
Evaluation of tetraalkylammonium tetrafluoroborate salts as high-temperature stationary phases for packed and open-tubular column GC 349(1985)249
D'Hoore, W., see Nelis, H.J.C.F. 333(1985)381
Di Bussolo, J.M. and Gant, J.R.
Vizualization of protein retention and migration in reversed-phase LC 327(1985)67
Di Corcia, A., see Andreolini, F. 337(1985)379
DiBella, E.E., Weissman, J.B., Joseph, M.J., Schultz, J.R. and Wenzel, T.J.
Lanthanide ions as luminescent chromophores for LC detection 328(1985)101
Dick, R.M., Freeman, J.J. and Kosh, J.W.
Use of radio GC for monitoring the *in vivo* labelling of postmorem [^{3}H]choline production 347(1985)387
Dickinson, C.M., Navin, T.R. and Churchill, F.C.
HPLC method for quantitation of pentamidine in blood serum 345(1985)91
DiDonato, G., see Brooks, M.A. 337(1985)351
Diehl, J.W., see Olson, E.S. 349(1985)337
Digilio, G., see Del Principe, D. 342(1985)285
Dijkgraaf, P.J.M., Verhaar, L.A.T., Groenland, W.P.T. and Van der Wiele, K.
High-speed LC of sugar acids on anion-exchange resins 329(1985)371
Dikkeschei, L.D., Van Veelen, H., Nagel, G.T., Willemse, P.H.B. and Wolthers, B.G.
Specific and sensitive determination of medroxyprogesterone acetate in human serum by GC-MS 345(1985)1
Dimov, N.
Unified retention index of hydrocarbons separated on squalane 347(1985)366
D'Incalci, M., see Farina, P. 345(1985)323
Diquet, B., Doare, L. and Simon, P.
Determination of clenbuterol in the high nanogram range in plasma of mice by HPLC with amperometric detection 336(1985)415
Dirks, R.J.M., see Wuis, E.W. 337(1985)341
Ditschuneit, H., see Swobodnik, W. 339(1985)263
Dixon, R., see Lucek, R. 341(1985)239
Dizdaroglu, M.
Weak anion-exchange HPLC of peptides (Review) 334(1985)49
Doare, L., see Diquet, B. 336(1985)415
Dobashi, A. and Hara, S.
Chiral recognition mechanisms in the enantioselectivity of chiral hydrogen bonding additives in LSC 349(1985)143
Dodo, G.H., Hawkes, S.J. and Thomas, L.C.
Optimized flow programming for temperature-programmed GC 328(1985)49
Dodo, G.H., Ramus, T.L. and Thomas, L.C.
Constant mass flow system for GC 330(1985)172
Doehl, J. and Greibrokk, T.
HPLC separation and determination of prostaglandins, oxidized by pyridinium dichromate. Optimization and applications 349(1985)431
Doizaki, W.M. and Levitt, M.D.
Measurement of mass and radioactivity of ^{14}CO using a reduction detector 329(1985)137
Dolanský, J., see Weidenhoffer, Z. 350(1985)324
Dolník, V., see Boček, P. 334(1985)157
Dolphin, R.J., see Mulholland, M. 350(1985)285
Dong, M.W. and Gant, J.R.
High-speed LC analysis of amino acids by post-column sodium hypochlorite-*o*-phthalaldehyde reaction 327(1985)17
Dooley, T. and Maddocks, J.L.
HPLC assay of 6-methylthioguanine, a major metabolite of 6-thioguanine, in urine 337(1985)321
Doom, J.P., see Carrel, J.E. 342(1985)411
Dordain, G., see Bougerolle, A.M. 345(1985)59
Döring, C.E., Estel, D., Pehle, W., Gaikowski, M. and Seiffarth, K.
Analysis of *tert.*-butylated cresol mixtures by capillary GC and capillary GC-MS 348(1985)430
Doris, P.A.
Separation of biologically important angiotensin peptides by HPLC on a weak cationic exchange bonded phase 336(1985)392
Dorizzi, R., see Tagliaro, F. 330(1985)323
Dorrbecker, B.R., Mercik, S.H. and Kramer, P.A.
Improved micro-method for the HPLC determination of caffeine and paraxanthine in biological fluids 336(1985)293
Dorschel, C., see Colgan, S.T. 333(1985)349
Dörschug, M., see Alpert, C.A. 326(1985)363
Dose, K., see Pabst, R. 350(1985)427
Dossena, A., see Palla, G. 349(1985)9
Double, J.A., see McDermott, B.J. 338(1985)335
Doucet, D., see Paillet, M. 343(1985)455

Douse, J.M.F.
Silica capillary GC of prostaglandins with electron-capture detection and its application to the forensic investigation of sexual offences 348(1985)111
Dow, J., Lemar, M., Frydman, A. and Gaillot, J.
Automated HPLC determination of spiramycin by direct injection of plasma, using column-switching for sample clean-up 344(1985)275
Dozier, S.E., see Williams, C.H. 344(1985)71
Dratz, E.A., see Van Kuijk, F.J.G.M. 348(1985)241
Drenth, B.F.H., see Feitsma, K.G. 333(1985)59
Dressler, D.E., see Leelavathi, D.E. 339(1985)105
Drevin, I., see Johansson, B.-L. 346(1985)255
Drewes, L.R., see Singh, A.K. 339(1985)170
Driskell, J.A., see Morrison, L.A. 337(1985)249
Drozdowicz, B., see Rotocki, P. 349(1985)155
Dryzga, M.D., see Freshour, N.L. 346(1985)376
Du Preez, J.L., Botha, S.A. and Lötter, A.P.
HPLC determination of phenazopyridine hydrochloride, tetracycline hydrochloride and sulphamethizole in combination 333(1985)249
Dua, V.K., Goso, K., Dube, V.E. and Bush, C.A.
Characterization of lacto-N-hexaose and two fucosylated derivatives from human milk by HPLC and proton NMR spectroscopy 328(1985)259
Dube, V.E., see Dua, V.K. 328(1985)259
Duddeck, H., see Kříž, J. 347(1985)297
Duez, P., Chamart, S., Hanocq, M., Molle, L., Vanhaelen, M. and Vanhaelen-Fastré, R.
Comparison between TLC-densitometry and HPLC for the determination of hyoscyamine and hyoscyne in leaves, fruits and seeds of *Datura* (*Datura* spp.) 329(1985)415
Duffield, A.M., see Low, G.K.C. 336(1985)15
Duffield, A.M., see Netting, A.G. 336(1985)115
Dufka, O., see Horna, A. 348(1985)141
Duke, R., see Zipser, R.D. 339(1985)1
Dumas, C.
The re-injection method for recovery of components adsorbed during syringe gas sampling: a statistical evaluation 329(1985)119
Dunaway, H.E., Hutchens, T.W. and Besch, P.K.
Isolation of unliganded steroid receptor proteins by high-performance size-exclusion chromatography. An investigation of steroid-dependent structural alterations 327(1985)221
Dunaway, H.E., see Hutchens, T.W. 327(1985)247
Duncan, M.W., Smythe, G.A., Nicholson, M.V. and Clezy, P.S.
Comparison of HPLC with electrochemical detection and GC-mass fragmentography for the assay of salsolinol, dopamine and dopamine metabolites in food and beverage samples 336(1985)199
Duquet, D., see Verzele, M. 329(1985)351
Durand, A., see Pok, P.R. 338(1985)171
Duval, M., see Godbillon, J. 345(1985)365
Dwyer, M.E. and Brown, P.R.
Evaluation of combined electrochemical and UV detection in the HPLC analysis of plasma for low-molecular-weight constituents 345(1985)125
Dye, D., see East, T. 338(1985)99

E

East, T. and Dye, D.
Determination of dextromethorphan and metabolites in human plasma and urine by HPLC with fluorescence detection 338(1985)99
Ebermann, R. and Gehringer, W.
Electrophoretic detection of peroxidase isoenzymes in polyacrylamide gels using hydrogen peroxide generated by rose bengal-glutathione in the presence of light 348(1985)313
Ebermann, R. and Lickl, E.
Improved staining procedure for nucleic acids in polyacrylamide gels after complexing with nitroso compounds 347(1985)384
Ebihara, M., see Goto, J. 345(1985)241
Ebinger, G., see Herregodts, P. 345(1985)33
Echols, R.E., Miller, R.H. and Thompson, L.
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89
Eddy, C.L., see Rodriguez, P.A. 330(1985)365
Edelson, J., see McCoy, L. 344(1985)211
Edols, R.W.
Simple method for the determination of diaminopimelic acid in rumen liquor hydrolysates 329(1985)199
Edwards, D.J., Mallinger, A.G., Knopf, S. and Himmelhoch, J.M.
Determination of tranylcypromine in plasma using GC-chemical-ionization MS 344(1985)356
Edwards, E.T.E., see Livesey, G. 337(1985)98
Edwards, G., Rodick, C.L., Ward, S.A., Awadzi, K., L'E Orme, M. and Breckenridge, A.M.
Determination of suramin in plasma by HPLC 343(1985)224
Edwards, G., see Mihaly, G.W. 337(1985)166
Edwards, I.R., see Schaaf, L.J. 342(1985)303
Efcavitch, J.W., see Becker, C.R. 326(1985)293
Efthymiopoulos, C., see Staveris, S. 339(1985)97

Egashira, S.
The Abbé refractometer as a LC detector using an image sensor for boundary location 332(1985)195
Eggert, F.M. and Jones, M.
Measurement of neutral sugars in glycoproteins as dansyl derivatives by automated HPLC 333(1985)123
Eglinton, G., see Gill, J.P. 350(1985)37
Egorin, M.J., Cohen, B.E., Kohlhepp, E.A. and Gutierrez, P.L.
GLC analysis of N,N',N"-triethylene thiophosphoramide and and N,N',N"-triethylene phosphoramide in biological samples 343(1985)196
Ehninger, G., Proksch, B. and Schiller, E.
Detection and separation of mitoxantrone and its metabolites in plasma and urine by HPLC 342(1985)119
Ehrat, M., Cecchini, D.J. and Giese, R.W.
Neonucleoproteins. Preparation and HPLC characterization of succinyl-lysozyme-diaminooctyl-polycytidylic acid and related polycytidylic acid conjugates 326(1985)311
Ehrsson, H., see Eksborg, S. 340(1985)31
Eichelbaum, M., see Von Sassen, W. 338(1985)113
Eichholtz, P.C.N., see Thieme, R.E. 337(1985)443
Einarsson, S.
Selective determination of secondary amino acids using precolumn derivatization with 9-fluorenylmethylchloroformate and reversed-phase HPLC 348(1985)213
Eisen, A.Z., see Seltzer, J.L. 326(1985)147
Ekberg, B., Sellergren, B. and Albertsson, P.-Å.
Direct chiral resolution in an aqueous two-phase system using the counter-current distribution principle 333(1985)211
Ekberg, B., see Sellergren, B. 347(1985)1
Eksborg, S. and Ehrsson, H.
Drug level monitoring: cytostatics (Review) 340(1985)31
El Rassi, Z. and Horváth, C.
HPLC of tRNAs on novel stationary phases 326(1985)79
Elbert, W., Breitenbach, S., Neftel, A. and Hahn, J.
4-methyl-7-methoxycoumarin as a fluorescent label for HPLC analysis of dicarboxylic acids 328(1985)111
Elgass, H., see Goetz, H. 349(1985)357
Elias, R., see Balansard, G. 348(1985)299
Elias, R., see Vidal, E. 348(1985)304
Eliason, M.S., see Karnatz, N.N. 330(1985)420
Ellington, K.R., see Patrick, K.S. 343(1985)329
Elliott, W.H., see Patrick, P.H. 347(1985)155
Ellis, B. and Jolly, J.
Purification of a uridine-specific acid nuclease from chicken liver by fast protein LC 326(1985)157
Ellis, P.P., see Rendi, M.A. 336(1985)337
Ellström, C., see Johansson, B.-L. 330(1985)360
Elworthy, P.M., Tsementzis, S.A., Westhead, D. and Hitchcock, E.R.
Determination of plasma tranexamic acid using cation-exchange HPLC with fluorescence detection 343(1985)109
Ely, T.D., see Kilts, C.D. 342(1985)452
Emran, A.M., see Boothe, T. 333(1985)269
Endoh, M., see Oyama, H. 338(1985)71
Engelen, D.L. van, see Van Engelen, D.L. 331(1985)77
Enns, R.K., see Herring, S.W. 326(1985)217
Ensminger, W.D., see Shukla, U.A. 342(1985)406
Ensminger, W.D., see Stetson, P.L. 341(1985)217, 344(1985)385
Eppert, G., Liebscher, G., Ihrke, M. and Arendt, G.
HPLC determination of contaminants in salicylic acids after computer-assisted optimization of the separation parameters (in German) 350(1985)471
Epps, D.E. and Kaluzny, M.A.
Cholesteryl ester quantitation: non-equivalent rates of transesterification of molecular species in the presence of silica gel 343(1985)143
Era, S., see Kuwata, K. 332(1985)29
Era, S., see Sogami, M. 332(1985)19
Erb, F., see Lhermitte, M. 342(1985)416
Erbersdobler, H.F., see Büser, W. 346(1985)363
Erdelmeier, C.A.J., Meier, B. and Sticher, O.
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456
Ergüven, H., see Carducci, C.N. 337(1985)91
Erickson, J.W., see Homandberg, G.A. 327(1985)343
Ericsson, Ö., see Svensson, J.-O. 341(1985)193
Eriksson, M., see Hermansson, J. 336(1985)321
Ervik, M., see Ahnoff, M. 340(1985)73
Erway, B., see Chang, T.-M. 326(1985)121
Eschbach, M.L., see Seltzer, J.L. 326(1985)147
Esclade, L., Guillochon, D. and Thomas, D.
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373
Estel, D., see Döring, C.E. 348(1985)430
Esterbauer, H., see Gleispach, H. 344(1985)11
Esterbauer, H., see Hayn, M. 329(1985)379
Estrada, G., see Ladrón de Guevara, O. 349(1985)91
Evans, D.B., see Homandberg, G.A. 327(1985)343
Evans, M.A., see Selan, F. 338(1985)213
Evans, W.E., see Choi, K.E. 345(1985)197
Everaerts, F.M., see Reijenga, J.C. 328(1985)353
Everhart, L.A., see Sisco, W.R. 348(1985)253
Evershed, R.P., see Gill, J.P. 350(1985)37
Eyk, J. van, see Cachia, P.J. 343(1985)315

F

Facchini, M.C., see Chiavari, G. 333(1985)262
Fagerstam, L., see Cooper, E.H. 327(1985)269
Faigle, J.W., see Francotte, E. 346(1985)321
Fallon, K.D., see Shao, McN.J. 345(1985)184
Fanali, S., Foret, F. and Boček, P.
Determination of sodium, ephedrine and procaine in pharmaceuticals by capillary isotachophoresis 330(1985)436
Farid, K.Z., Fasola, A.F. and Nash, J.F.
LC determination of indecainide, a new antiarrhythmic drug, and its major metabolite, desisopropyl indecainide, in biological samples 337(1985)329
Farina, P., Benfenati, E., Lassiani, L., Nisi, C. and D'Incalci, M.
HPLC assay for the determination of *p*-(3,3-dimethyl-1-triazeno)benzoic acid in mouse plasma 345(1985)323
Fasching, J.L., see Brown, P.R. 345(1985)457
Fasching, J.L., see Yi, C. 339(1985)75
Fasola, A.F., see Farid, K.Z. 337(1985)329
Faucon, G., see Al Makdessi, S. 339(1985)25
Federici, E., see Gagliardi, L. 348(1985)321
Fehrenbach, F.J., see Jürgens, D. 348(1985)363
Feitsma, K.G., Bosman, J., Drenth, B.F.H. and De Zeeuw, R.A.
Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59
Felix, A.M., Heimer, E.P., Lambros, T.J., Swistok, J., Tarnowski, S.J. and Wang, C.-T.
Analysis of different forms of recombinant human leukocyte interferons and synthetic fragments by HPLC 327(1985)359
Fell, D., Benjamin, L.E. and Steele, R.D.
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 345(1985)150
Fenemore, J., see Owen, J.A. 342(1985)333
Fernández-Sánchez, E., García-Domínguez, J.A., Menéndez, V. and Pertierra-Rimada, E.
Indirect determination of specific retention volumes in GC. Use of the retention indices of McReynolds' samples 333(1985)1
Fernández-Sánchez, E., see Reglero, G. 348(1985)327
Ferrara, F.J., see Puglisi, C.V. 344(1985)199
Ferrario, P., see Zecca, L. 337(1985)391
Ferre, J. and Jacobson, K.B.
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389
Ferreres, F., see Barberan, F.A.T. 347(1985)443
Ferry, D.G., see Schaaf, L.J. 342(1985)303
Ferry, N., see Cuisinaud, G. 341(1985)97
Fetzer, J.C. and Biggs, W.R.
LC retention behavior of large, fused polycyclic aromatics. Normal bonded phases 346(1985)81
Field, R.E., see Pirkle, W.H. 348(1985)89
Fields, M.J., see Roberts, R.F. 342(1985)25
Fields, S.M. and Lee, M.L.
Effects of density and temperature on efficiency in capillary supercritical fluid chromatography 349(1985)305
Finden, D.A.S., Fysh, R.R. and White, P.C.
Quantitative method for the detection of glucose in body fluids by HPLC with fluorescence detection 342(1985)179
Fini, C., see Palmerini, C.A. 339(1985)285
Finn, R.D., see Boothe, T. 333(1985)269
Fisch, H.-U., see Baktir, G. 339(1985)192
Fischer, J.H., see Cwik, M.J. 338(1985)123
Flanagan, R.J., see Caldwell, R. 345(1985)187
Floridi, A., see Palmerini, C.A. 339(1985)285
Flouvat, B., see Canal, M. 342(1985)212
Foglietti, M.-J., see Succari, M. 341(1985)457
Fong, M.H., Aarons, L., Urso, R. and Caccia, S.
Ionization constants and lipophilicity of 1-arylpiperazines investigated by reversed-phase HPLC 333(1985)191
Ford, C.W.
Identification of inositols and their mono-O-methyl ethers by GLC 333(1985)167
Ford, G.L.
Semi-preparative isolation of crepenynic acid, a potential inhibitor of essential fatty acid metabolism 346(1985)431
Foreman, W.T. and Bidleman, T.F.
Vapor pressure estimates of individual polychlorinated biphenyls and commercial fluids using GC retention data 330(1985)203
Foret, F., see Boček, P. 334(1985)157
Foret, F., see Fanali, S. 330(1985)436
Forgue, S.T., Van Harken, D.R. and Smyth, R.D.
Determination of tiodazosin concentrations in human plasma with a fluorescence HPLC method 342(1985)221
Först, C., see Liebich, H.M. 338(1985)33, 338(1985)187
Fořtová, H., see Přistoupil, T.I. 346(1985)401
Forward, R. and Almog, R.
Separation of two forms of glutathione peroxidase from human erythrocytes by hydrophobic chromatography 330(1985)383
Fouillit, M., see Leclercq, M. 337(1985)423
Fox, A., see Whiton, R.S. 347(1985)109
Fox, J.M., see Chang, S.F. 343(1985)119
Frache, R., see Mazzucotelli, A. 349(1985)137
Francis, F.E. and Kinsella, R.A. Jr.
Reversed-phase HPLC of C_{21} metabolites of progesterone 336(1985)361
Francotte, E., Stierlin, H. and Faigle, J.W.
Isolation and stereospecific determination of the enantiomers of oxindazac by direct LC resolution on triacetylcellulose 346(1985)321
Francotte, E., Wolf, R.M., Lohmann, D. and Mueller, R.

Chromatographic resolution of racemates on chiral stationary phases. I. Influence of the supramolecular structure of cellulose triacetate 347(1985)25
Frank, N.E., see Hollis, B.W. 343(1985)43
Frank, R., see Alpert, C.A. 326(1985)363
Franke, J.P., see Bogusz, M. 342(1985)241
Franklin, M., see Molyneux, S.G. 341(1985)160
Franklin, M.R., see Liu, Z. 342(1985)135
Franklin, R.B., see Bernstein, J.R. 342(1985)228
Frantz, S.W., see Freshour, N.L. 346(1985)376
Franz, H.E., see Kohse, K.P. 344(1985)319
Franzoi, L., see Rusconi, L. 349(1985)117
Frattini, P., see Schinelli, S. 338(1985)396
Fredj, G., see Brouard, A. 345(1985)453
Fredj, G., see Paillet, M. 343(1985)455
Freeman, J.J., see Dick, R.M. 347(1985)387
Frei, R.W., see Nielen, M.W.F. 330(1985)113
Frenz, J., Van der Schrieck, P. and Horváth, C.
Investigation of operating parameters in high-performance displacement chromatography 330(1985)1
Freshour, N.L., Langvardt, P.W., Frantz, S.W. and Dryzga, M.D.
Direct determination of acrylamide in tissue culture solution by LC using column switching 346(1985)376
Freundlich, L., see Karmen, A. 349(1985)267
Fričová, V., see Přistoupil, T.I. 346(1985)401
Frigerio, A., see Tagliaro, F. 330(1985)323
Frith, R.G. and Phillipou, G.
15-Hydroxycyproterone acetate and cyproterone acetate levels in plasma and urine 338(1985)179
Fritig, B., see Hermann, C. 348(1985)167
Fritz, J.S., see Ogawa, I. 329(1985)81
Fritz, J.S., see Sevenich, G.J. 347(1985)147
Fromark, I., see Johansson, B. 341(1985)462
Frommer-Filep, M., see Gál, S. 333(1985)239
Frydman, A., see Dow, J. 344(1985)275
Fugier, J.-C., see Al Makdessi, S. 339(1985)25
Fujimoto, C., Hirata, Y. and Jinno, K.
Supercritical fluid chromatography-IR spectroscopy of oligomers: use of buffer-memory technique 332(1985)47
Fujimura, K., Kitagawa, M., Takayanagi, H. and Ando, T.
HPLC separation of aromatic carboxylic acids with β-cyclodextrin-bonded stationary phases 350(1985)371
Fujimura, K., see Takayanagi, H. 350(1985)63, 350(1985)75
Fujioka, M., see Baba, S. 337(1985)205
Fujiwara, S., see Suzuki, H. 341(1985)341
Fujiwara, T., see Tokuma, Y. 345(1985)51
Fukita, T., see Tsuchiya, H. 339(1985)59
Fukuda, H., see Yamatodani, A. 344(1985)115
Fukuda, R., see Kurosawa, N. 343(1985)379
Fukui, K., see Okamoto, T. 342(1985)35
Fukui, T., see Ozaki, H. 332(1985)243
Fullinfaw, R.O., see Kacprowicz, A.T. 337(1985)412
Funazo, K., Tanaka, M., Morita, K., Kamino, M., Shono, T. and Wu, H.-L.
Pentafluorobenzyl *p*-toluenesulphonate as a new derivatizing reagent for GC determination of anions 346(1985)215
Fung, H.-L., see King, S.-Y.P. 343(1985)129
Furness, J.B., see Murphy, R. 336(1985)41
Fürst, P., see Kohse, K.P. 344(1985)319
Fürst, P., see Stehle, P. 346(1985)271
Furton, K.G. and Poole, C.F.
Variation in the GC stationary phase properties of tetra-*n*-butylammonium salts as a function of the anion type 349(1985)235
Furukawa, H., see Niwa, T. 337(1985)1
Furusawa, M., see Kiba, N. 328(1985)309, 329(1985)147
Furuta, H., Yamane, T. and Sugiyama, K.
Separation of cadaverine from putrescine, histamine and polyamines in rat kidney by phosphocellulose chromatography 337(1985)103
Furuta, T., see Baba, S. 337(1985)205
Fuwa, K., see Yasuhara, A. 328(1985)35
Fuzzi, S., see Chiavari, G. 333(1985)262
Fysh, R.R., see Finden, D.A.S. 342(1985)179

G

Gadde, R.R., see Burton, F.W. 328(1985)317
Gaertner, H. and Puigserver, A.
Separation of some peptides and related isopeptides by HPLC: structure-retention time relationships 350(1985)279
Gaffney, J.S., Spandau, D.J., Kelly, T.J. and Tanner, R.L.
GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121
Gaget, C., see Claudy, P. 329(1985)331
Gagliardi, L., Amato, A., Basili, A., Cavazzutti, G., Federici, E., Chimenti, F., Casanova, M.G., Gattavecchia, E. and Tonelli, D.
Determination of preservatives in cosmetic products by ion-pair reversed-phase HPLC. III 348(1985)321
Gagner, J.-P. and Gauthier, S.
Estimation of adrenal catecholamines by elevated-temperature LC with amperometric detection 342(1985)363
Gaikowski, M., see Döring, C.E. 348(1985)430
Gaillot, J., see Dow, J. 344(1985)275
Gal, J.
Determination of the enantiomeric composition of chiral epoxides using chiral derivatization and LC 331(1985)349
Gal, J., see Rendi, M.A. 336(1985)337
Gál, S., Frommer-Filep, M., Toth-Martinez, B.L., Hernádi, F.J. and Kiss, L.

Use of chromatofocusing for separation of β-lactamases. V. Inducible chromosomally mediated β-lactamase of the *Enterobacter cloacae* 53 strain 333(1985)239
Gál, S., see Kiss, L. 333(1985)244
Galeazzi, R., see Daenens, P. 342(1985)79
Gallice, P., Monti, J.-P. and Crevat, A.
Removal of counter-ion reagent after semi-preparative reversed-phase ion-pair HPLC 331(1985)445
Gallion, C., see Jehl, F. 339(1985)347
Gallo, P., Olsson, O. and Sidén, Å.
Methods for chromatofocusing of cerebrospinal fluid and serum immunoglobulin G 327(1985)293
Galushko, S.V. and Shishkina, I.P.
Reversed-phase ion-paired HPLC determination of 6-azacytidine in blood 345(1985)157
Gambardella, P., Punziano, R., Gionti, M., Guadalupi, C., Mancini, G. and Mangia, A.
Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229
Gamerith, G., Zuder, G.F.X., Giuliani, A. and Brantner, H.
GLC determination of free amino acids in fermentation broth during growth of *Clostridium oncolyticum* M 55 328(1985)241
Gammans, R.E., Kerns, E.H. and Bullen, W.W.
Capillary GC-MS determination of buspirone in plasma 345(1985)285
Gammans, R.E., Kerns, E.H., Bullen, W.W., Covington, R.R. and Russell, J.W.
GC-MS method for trazodone and a deuterated analogue in plasma 339(1985)303
Ganno, S., Hamano, Y., Kobayashi, J. and Masaki, T.
Single-column separation of aminoethylcysteine and other amino acids 332(1985)275
Gant, J.R., see Di Bussolo, J.M. 327(1985)67
Gant, J.R., see Dong, M.W. 327(1985)17
Garceau, Y., see Molles, R.J. Jr. 347(1985)414
Garcia, C.P.
Rubeanic acid: a general spray reagent for the detection of steroids on thin-layer plates 350(1985)468
García-Domínguez, J.A., see Fernández-Sánchez, E. 333(1985)1
García-Domínguez, J.A., see Reglero, G. 348(1985)327
García-Raso, A., Vázquez, M.A. and Ballester, P.
GC of bicyclo[*n*.4.0]alken-2-ones 331(1985)406
Gardner, G.J., see Siu, K.W.M. 330(1985)87
Garland, W.A., see Min, B.H. 336(1985)403
Gariand, W.A., see Rubio, F. 339(1985)313
Garraffo, R. and Lapalus, P.
Simplified LC analysis for cyclosporin A in blood and plasma with use of rapid extraction 337(1985)416
Garrettson, L.K., see Bhargava, V.O. 343(1985)219
Gaskell, S.J., see Parker, C.E. 344(1985)378, 347(1985)61
Gaspar, P. and Maghuin-Rogister, G.
Rapid extraction and purification of diethylstilboestrol in bovine urine hydrolysates using reversed-phase C_{18} columns before determination by radioimmunoassay 328(1985)413
Gattavecchia, E., see Gagliardi, L. 348(1985)321
Gaudry, D., Wantiez, D., Richard, J. and Metayer, J.P.
Simultaneous determination of metoprolol and deuterium-labelled metoprolol in human plasma by GC-negative-ion MS 339(1985)404
Gauron, S., see Godbillon, J. 338(1985)151
Gauthier, S., see Gagner, J.-P. 342(1985)363
Gaydou, E.M., see Bianchini, J.-P. 329(1985)231
Gazdag, M., Szepesi, G., Varsányi-Riedl, K., Végh, Z. and Pap-Sziklay, Z.
Chromatography of bis-quaternary amino steroids. I. Separation on silica by TLC and HPLC 328(1985)279
Gazdag, M., Varsányi-Riedl, K. and Szepesi, G.
Chromatography of bis-quaternary amino steroids. II. Separation on chemically bonded silica phases 347(1985)284
Gazzotti, G., Sonnino, S. and Ghidoni, R.
Normal-phase HPLC separation of non-derivatized ganglioside mixtures 348(1985)371
Gebauer, P., see Boček, P. 334(1985)157
Geerdink, R.B., see Maris, F.A. 328(1985)93, 347(1985)75
Gehas, J., see Metz, P.A. 330(1985)307
Gehringer, W., see Ebermann, R. 348(1985)313
Gehrke, C.W., see Williams, C.H. 344(1985)71
Geissler, W., see Wahl, R. 329(1985)153
Gelder, W.M.J. van, see Van Gelder, W.M.J. 331(1985)285
Gelsema, W.J., see Brandts, P.M. 333(1985)41
Geng, X. and Regnier, F.E.
Stoichiometric displacement of solvent by non-polar solutes in reversed-phase LC 332(1985)147
Genin-Ramakers, J., see Kumps, A. 342(1985)469
Geoffroy, P., see Hermann, C. 348(1985)167
Georges, Y., see Désage, M. 336(1985)285
Gerhardt, K.O., see Williams, C.H. 344(1985)71
Gerlo, E. and Malfait, R.
HPLC assay of free norepinephrine, epinephrine, dopamine, vanillylmandelic acid and homovanillic acid 343(1985)9
Gerner, T.H., see Boeren, E.G. 349(1985)377
Gerritse, R.G. and Adeney, J.A.
Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419
Gerry, N., see Robinson, Y. 338(1985)219
Gharfeh, S.G., see Pearson, C.D. 329(1985)142
Ghatge, B.B., see Bendre, S.B. 346(1985)359

Ghïdoni, R., see Gazzotti, G. 348(1985)371
Gianazza, E., see Bosisio, A.B. 330(1985)299
Gianazza, E., see Righetti, P.G. 334(1985)71
Giannola, L.I., see Bousquet, E. 344(1985)325
Giarrusso, A., see Cartoni, G.P. 330(1985)315
Gierz, J., see Bogusz, M. 342(1985)241
Giese, R.W., see Adams, J. 347(1985)99
Giese, R.W., see Ehrat, M. 326(1985)311
Giese, R.W., see Sentissi, A. 331(1985)396
Gifford, R., see Swagzdis, J.E. 345(1985)203
Gilbart, J., see Whiton, R.S. 347(1985)109
Gilbert, M., see Grech-Bélanger, O. 337(1985)172
Gilchrist, D.G., see Clouse, S.D. 350(1985)255
Giliáš, J. and Novák, J.
Determination of ethylene by a GC enrichment technique with a flame ionization detector (in German) 346(1985)43
Gill, J.P., Evershed, R.P., Chicarelli, M.I., Wolff, G.A., Maxwell, J.R. and Eglinton, G.
Computerised capillary GC-MS studies of the petroporphyrins of the Gilsonite bitumen (Eocene, U.S.A.) 350(1985)37
Gill, R. and Key, J.A.
HPLC system for the separation of ergot alkaloids with applicability to the analysis of illicit lysergide (LSD) 346(1985)423
Gionti, M., see Gambardella, P. 348(1985)229
Giordano, R., see Castiglioni, M. 349(1985)173
Girard, F., see Godin, J. 343(1985)424
Giuliani, A., see Gamerith, G. 328(1985)241
Giulietti, G., see Musumarra, G. 350(1985)151
Glad, M., Norrlöw, O., Sellergren, B., Siegbahn, N. and Mosbach, K.
Use of silane monomers for molecular imprinting and enzyme entrapment in polysiloxane-coated porous silica 347(1985)11
Glebko, L.I., see Pokushalova, T.V. 329(1985)189
Gleispach, H., Moser, R. and Leis, H.J.
Contribution to the measurement of prostaglandins by GC-MS 342(1985)245
Gleispach, H., Moser, R., Mayer, B., Esterbauer, H., Skriletz, U., Zierman, L. and Leis, H.J.
Qualitative and quantitative measurement of hydroxy fatty acids, thromboxanes and prostaglandins using stable isotope dilutions and detection by GC-MS 344(1985)11
Glennon, J.D., see Hojabri, H. 342(1985)97
Gluckman, J.C. and Novotny, M.
Nitrogen-sensitive thermionic detection in microcolumn LC 333(1985)291
Gmur, D.J., Yee, G.C. and Kennedy, M.S.
Modified column-switching HPLC method for the measurement of cyclosporine in serum 344(1985)422
Go, J.G., see Koopman, B.J. 337(1985)259
Gobaille, J., see Cillard, J. 347(1985)434
Godbillon, J., Duval, M. and Gosset, G.
Determination of oxprenolol in human plasma by HPLC, in comparison with GC and GC-MS 345(1985)365
Godbillon, J., Gauron, S. and Metayer, J.P.
HPLC determination of diclofenac and its monohydroxylated metabolites in biological fluids 338(1985)151
Goddard, C., see Slack, J.A. 337(1985)178
Godel, H.G., see Kohse, K.P. 344(1985)319
Godin, J., Girard, F. and Hellier, G.
Rapid GC method for the determination of nicotine and cotinine in urine (in French) 343(1985)424
Goebel, K.-J. and Kölle, E.U.
HPLC determination of diltiazem and four of its metabolites in plasma. Application to pharmacokinetics 345(1985)355
Goetz, H., Elgass, H. and Huber, L.
Determination of molecular weight data in size-exclusion chromatography by use of braod-molecular-weight-distributed standards 349(1985)357
Goewie, C.E. and Hogendoorn, E.A.
LC determination of bromide in human milk and plasma 344(1985)157
Goffic, F. le, see Léonil, J. 347(1985)316
Goheen, S.C. and Matson, R.S.
Purification of human serum gamma globulins by hydrophobic interaction HPLC 326(1985)235
Golay, M.J.E.
A problem of optimization in capillary GC 348(1985)416
Gold, H.S., see Stadalius, M.A. 327(1985)27
Goldstein, S., Stivaletta, L.A. and Phillips, L.S.
Separation of somatomedins and somatomedin inhibitors by size exclusion HPLC 339(1985)388
Gombert, J., see Picard, M. 341(1985)445
Gonnet, C., Morel, D., Ramamonjinirina, E., Serpinet, J., Claudy, P. and Letoffé, J.M.
Insertion of various long alkyl chain molecules in brush-type grafted monolayers. Chromatographic study of the resulting materials 330(1985)227
Gonnet, C., see Boulieu, R. 339(1985)380
Gooding, K.M. and Schmuck, M.N.
Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139
Goodwin, K.R., see Schmerr, M.J.F. 326(1985)225
Gopal, N.G.S. and Sharma, G.
Shelf-life of unused HPLC columns 346(1985)461
Gorham, J. and McDonnell, E.
HPLC method for the separation and estimation of choline, glycinebetaine aldehyde and related compounds 350(1985)245
Gorodetzky, C.W., see Cone, E.J. 337(1985)291
Gorp, L. van, see Marescau, B. 345(1985)215
Goshorn, C., see Park, Y.H. 343(1985)359
Goso, K., see Dua, V.K. 328(1985)259
Gosset, G., see Godbillon, J. 345(1985)365
Goto, J., Suzaki, K., Chikai, T., Nagase, K. and Nambara, T.
Studies on steroids. CCXVI. Separation of

bile acid 3-glucuronides by HPLC 348(1985)151
Goto, J., Suzaki, K., Ebihara, M., Nambara, T. and Masu, A.
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 345(1985)241
Goto, M., Irino, K. and Ishii, D.
Multichannel spectrophotometric detector for fused-silica capillary tube isotachophoresis 346(1985)167
Goto, M., see Ito, Y. 346(1985)161
Goulle, J.P., see Lacroix, C. 345(1985)436
Gounaris, A., see Dalas, E. 348(1985)339
Grahl, R., see Dedek, W. 331(1985)200
Grant, A. and Bhattacharyya, P.K.
Application of derivative spectroscopy to the determination of chromatographic peak purity 347(1985)219
Grant, H.G., see Jefford, C.W. 347(1985)183
Graser, T.A., see Kohse, K.P. 344(1985)319
Grassini-Strazza, G., see Dębowski, J. 349(1985)131
Gray, J.E., see Schallinger, L.E. 342(1985)67
Grech-Bélanger, O., Turgeon, J. and Gilbert, M.
HPLC assay for mexiletine enantiomers in human plasma 337(1985)172
Greef, J. van der, see Van der Greef, J. 343(1985)397
Green, J.P., see Khandelwal, J.K. 343(1985)249
Greenblatt, D.J., see Locniskar, A. 337(1985)131
Greene, S.V. and Smith, J.M.
High-performance liquid chromatographic detection of XMP as a basis for an improved IMP dehydrogenase assay 343(1985)160
Grego, B. and Hearn, M.T.W.
HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25
Gregory, V.M., Larsen, B. and Benson, B.
Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 345(1985)140
Greibrokk, T., see Blilie, A.L. 349(1985)317
Greibrokk, T., see Doehl, J. 349(1985)431
Greibrokk, T., see Langseth-Manrique, K. 349(1985)421
Greibrokk, T., see Lundanes, E. 349(1985)439
Greig, N.H., see Sweeney, D.J. 339(1985)434
Griffiths, R., see Allen, I.C. 336(1985)385
Grimvall, A., see Borén, H. 348(1985)67
Grimvall, A., see Sävenhed, R. 328(1985)219
Grinberg, N.
TLC separations using a temperature gradient 333(1985)69
Gringore, A., see Lacroix, C. 345(1985)436
Grisham, C.M., Marquard, F. and Jørgensen, P.L.
Versatile HPLC computer system for solvent delivery, gradient control and data acquisition using the Commodore 64 personal computer 333(1985)301
Grob, G., see Grob, K. 347(1985)351
Grob, K. and Grob, G.
New approach to capillary columns for GC? Condensation of hydroxyl-terminated stationary phases 347(1985)351
Grob, K., see Blum, W. 346(1985)341
Grob, K. Jr.
Connecting capillaries by fusion 330(1985)217
Grob, K. Jr., Karrer, G. and Riekkola, M.-L.
On-column injection of large sample volumes using the retention gap technique in capillary GC (Review) 334(1985)129
Grob, K. Jr. and Läubli, T.
Rejection of the sample plug after on-column injection into capillaries at high temperatures 333(1985)156
Grob, R.L., see Bigley, F.P. 350(1985)407
Groen, G., see Welling, G.W. 326(1985)173
Groeningen, C.J. van, see Kok, R.M. 343(1985)59
Groenland, W.P.T., see Dijkgraaf, P.J.M. 329(1985)371
Gronlund, S., see McGregor, J.L. 326(1985)179
Groppi, A. and Papa, P.
One-step extraction procedure for GC determination of viloxazine as its acetyl derivative in human plasma 337(1985)142
Gross, R.W., see Creer, M.H. 338(1985)61
Grossenbacher, H., Cook, A.M. and Hütter, R.
Determination of biuret and urea by HPLC 331(1985)161
Guadalupi, C., see Gambardella, P. 348(1985)229
Guay, D.R.P., see Meatherall, R.C. 338(1985)141
Guereca, L., see Ladrón de Guevara, O. 349(1985)91
Guerra, M.C., see Barbaro, A.M. 347(1985)209
Guevara, O. Ladrón de, see Ladrón de Guevara, O. 349(1985)91
Guevara, O.L. de, see De Guevara, O.L. 329(1985)428
Guigues, M., see Vidal, E. 348(1985)304
Guillén, M.D., see Bermejo, J. 331(1985)237
Guillochon, D., see Esclade, L. 341(1985)373
Guinet, R., see Leclercq, M. 337(1985)423
Guinn, G. and Hendrix, D.L.
Bracketing, a simple loading technique that increases sample recovery, improves resolution, and increases sensitivity in HPLC 348(1985)123
Guiochon, G. and Martin, M.
Theoretical investigation of the optimum particle size for the resolution of proteins by size-exclusion chromatography 326(1985)3
Guirakhoo, F., see Winkler, G. 326(1985)113
Gunasingham, H., Tay, B.T. and Ang, K.P.
Determination of oestriol in pregnancy urine by normal-phase HPLC with electrochemical detection 341(1985)271

Gunata, Y.Z., Bayonove, C.L., Baumes, R.L. and Cordonnier, R.E.
The aroma of grapes. I. Extraction and determination of free and glycosidically bound fractions of some grape aroma components 331(1985)83
Günther, K., see Busker, E. 350(1985)179
Gupta, R.N.
Drug level monitoring: Sedative hypnotics (Review) 340(1985)139
Gupta, R.N. and Lew, M.
Determination of trazodone in human plasma by LC with fluorescence detection 342(1985)442
Gupta, R.N. and Lew, M.
LC determination of mexiletine and tocainide in human plasma with fluorescence detection after reaction with a modified *o*-phthalaldehyde reagent 344(1985)221
Gurley, L.R., see Jackson, P.S. 326(1985)199
Gutierrez, P.L., see Egorin, M.J. 343(1985)196
Guyonnet, J., see Davi, H. 342(1985)159
Gyllenhaal, O.
Direct derivatization of alprenolol and its 4-hydroxy metabolite in urine with phosgene and methanol prior to analysis by capillary column GC 349(1985)447
Gyllenhaal, O., König, W.A. and Vessman, J.
Enantiomer separation of metoprolol and its analogues and metabolites by capillary column GC after derivatization with phosgene 350(1985)328
Györfi, L., see Cserháti, T. 349(1985)295

H

Ha, Y.C. and Barter, P.J.
Rapid separation of plasma lipoproteins by GPC on agarose gel Superose 6B 341(1985)154
Haagsma, N. and Van de Water, C.
Rapid determination of five sulphonamides in swine tissue by HPLC 333(1985)256
Haan, H.B.P., see Lingeman, H. 336(1985)241
Haddad, G.R., see Santa María, I. 329(1985)123
Haddad, P.R., Jackson, P.E. and Heckenberg, A.L.
Performance characteristics of some commercially available low-capacity anion-exchange columns suitable for non-suppressed ion chromatography 346(1985)139
Haddad, P.R., see Heckenberg, A.L. 330(1985)95
Haddad, P.R., see Jackson, P.E. 346(1985)125
Haddad, P.R., see Low, G.K.C. 336(1985)15
Hagardorn, A.N., see Robert, T.A. 344(1985)177
Hagel, L., see Andersson, T. 326(1985)33
Hagemeier, E., see Kemper, K. 336(1985)374
Hagen, B., Walseth, F., Walstad, R.A. and Iversen, T.
GC assay of triethylenethiophosphoramide in serum and urine 345(1985)173
Haginaka, J., Wakai, J., Yasuda, H., Uno, T. and Nakagawa, T.
HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 341(1985)115
Hahn, J., see Elbert, W. 328(1985)111
Haikala, V.
New sensitive method to determine noscapine in serum by reversed-phase LC 337(1985)429
Haines-Nutt, R.F., see Adams, P.S. 329(1985)438
Hajšlová, J., Rathouská, Z. and Davídek, J.
GLC method for the determination of dodine in fruit 348(1985)437
Hakam, A. and Kun, E.
HPLC of *in vitro* synthesized poly(ADP-ribose) on ion-exchange columns, separation of oligomers of varying chain length and estimation of apparent branching 330(1985)287
Haken, J.K., Burford, R.P. and Vimalasiri, P.A.D.T.
Chromatographic analysis of thermoplastic, medical-grade, polyether-based polyurethane elastomers 349(1985)347
Haken, J.K., see Burford, R.P. 329(1985)132
Haken, J.K., see Itsikson, L.B. 334(1985)1
Haken, J.K., see Smith, R.J. 328(1985)11, 331(1985)389, 334(1985)95
Halász, A., see Szakács-Dobozi, M. 347(1985)310
Halbert, M.K. and Baldwin, R.P.
Determination of cysteine and glutathione in plasma and blood by LC with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 345(1985)43
Haldna, Ü., Palvadre, R., Pentshuk, J. and Kleemeier, T.
Preparation of low-capacity anion-exchange resins for ion chromatography on a methacrylic copolymer matrix 350(1985)296
Hale, P.W. Jr. and Poklis, A.
Determination of diastereoisomeric pairs of thioridazine 5-sulfoxide by HPLC 336(1985)452
Halfpenny, A.P. and Brown, P.R.
Simultaneous HPLC assay of the activities of erythrocytic hypoxanthine-guanine phosphoribosyl transferase and purine nucleoside phosphorylase 349(1985)275
Halkiewicz, J., see Ośmiałowski, K. 346(1985)53
Hallowell, S.F., see Bossle, P.C. 330(1985)388
Hamada, C., Iwasaki, M., Kuroda, N. and Ohkura, Y.
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for

the assay of 17-oxosteroids in urine 341(1985)426
Hamano, Y., see Ganno, S. 332(1985)275
Hamilton, G., Roth, E., Wallisch, E. and Tichy, F.
Semi-automated HPLC determination of cyclosporine A in whole blood using one-step sample purification and column-switching 341(1985)411
Hanai, T.
Energy effects in the retention of aromatic acids in LC 332(1985)189
Hanai, T., Arai, Y., Kirukawa, M., Noguchi, K. and Yanagihara, Y.
Reversed-phase LC on rigid vinyl alcohol copolymer gels 349(1985)323
Hancock, W.S., see Knighton, D.R. 347(1985)237
Hancock, W.S., see Poll, D. 343(1985)149
Handelman, G.J., see Van Kuijk, F.J.G.M. 348(1985)241
Hanke, F.J. and Kubo, I.
Increasing the speed of droplet counter-current chromatography separations 329(1985)395
Hanna, M.H., see Tameesh, A.H.H. 328(1985)207
Hannan, S.F., see Poll, D. 343(1985)149
Hanocq, M., see Duez, P. 329(1985)415
Hansen, A.S., see D'Agostino, P.A. 347(1985)257
Hansen, B., see Welinder, B.S. 348(1985)347
Hansen, E.B. Jr., see Rushing, L.G. 337(1985)37
Hansen, L.B., see Larsen, N.-E. 341(1985)244
Hara, I., see Okazaki, M. 336(1985)151
Hara, S., Takemori, Y., Yamaguchi, M., Nakamura, M. and Ohkura, Y.
Determination of α-keto acids in serum and urine by HPLC with fluorescence detection 344(1985)33
Hara, S., see Dobashi, A. 349(1985)143
Hara, S., see Iwata, T. 344(1985)351
Hara, S., see Oka, K. 339(1985)253
Hara, S., see Yamaguchi, M. 346(1985)227
Harada, T., Iwamori, M., Nagai, Y. and Nomura, Y.
Analysis of aminoglycoside antibiotics as benzoyl derivatives by HPLC and its application to the quantitation of neomycin in the perilymph 337(1985)187
Haralambidis, J., see Scanlon, D. 336(1985)189
Harding, D.R.K., see Knighton, D.R. 347(1985)237
Harding, D.R.K., see Poll, D. 343(1985)149
Harenberg, J., see De Vries, J.X. 338(1985)325
Harken, D.R. van, see Forgue, S.T. 342(1985)221
Härkönen, H., see Pfäffli, P. 337(1985)146
Harmsen, E., De Tombe, P.P. and De Jong, J.W.
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides 337(1985)201
Harriott, M., see Adair, C.G. 336(1985)429, 342(1985)447
Harris, P.J., Bacic, A. and Clarke, A.E.
Capillary GC of partially methylated alditol acetates on a SP-2100 wall-coated open-tubular column 350(1985)304
Harris, R., see Calam, D.H. 326(1985)103
Harrison, P.M., Tonkin, A.M., Cahill, C.M. and McLean, A.J.
Rapid and simultaneous extraction of propranolol, its neutral and basic metabolites from plasma and assay by HPLC 343(1985)349
Harrison, P.M., Tonkin, A.M. and McLean, A.J.
Determination of 4-amino-3-(*p*-chlorophenyl)butyric acid (baclofen) in plasma by HPLC 339(1985)424
Harrison, P.M., Tonkin, A.M. and McLean, A.J.
Simple and rapid analysis of atenolol and metoprolol in plasma using solid-phase extraction and HPLC 339(1985)429
Hartley, R., Smith, I.J. and Cookman, J.R.
Improved HPLC method for the simultaneous determination of caffeine and its N-demethylated metabolites in plasma using solid-phase extraction 342(1985)105
Harvey, V.J., Joel, S.P., Johnston, A. and Slevin, M.L.
HPLC of etoposide in plasma and urine 339(1985)419
Hasegawa, A., see Tanaka, T. 328(1985)271
Hashimoto, T., see Kato, Y. 332(1985)39, 333(1985)93, 333(1985)202
Hashimoto, Y., see Ishii, D. 332(1985)3
Hashimoto, Y., see Takeuchi, T. 331(1985)99
Hashizume, M., see Takayanagi, H. 350(1985)63
Hatada, K., see Okamoto, Y. 350(1985)127
Hatake, K., Motoyoshi, K., Ishizaka, Y., Sato, M., Takaku, F. and Miura, Y.
Purification of human urinary colony-stimulating factor by HPLC 344(1985)339
Hatano, H.
Recent progress in gel packing materials and detectors for modern LC in Japan 332(1985)227
Hatano, H., see Makino, K. 332(1985)71
Hatano, H., see Naikwadi, K.P. 331(1985)69
Hatano, H., see Rokushika, S. 332(1985)15
Hatem, S.F., see Munavalli, S. 346(1985)463
Hauser, A.R., see Llyod, C.W. 339(1985)139
Hawkes, S.J., see Dodo, G.H. 328(1985)49
Hayano, S., see Sogami, M. 332(1985)19
Hayashi, K., Miyamoto, M. and Sekine, Y.
Determination of captopril and its mixed disulphides in plasma and urine by HPLC 338(1985)161
Hayashi, M., Imai, Y., Minami, Y., Kawata, S., Matsuzawa, Y. and Tarui, S.
Determination of bile acids in rat bile by HPLC 338(1985)195
Hayashi, R., see Nagashima, H. 345(1985)373
Hayashi, T., see Tsuchiya, H. 339(1985)59
Haycock, G.B., see Cullen, M.P. 337(1985)29
Hayes, A.G., Mehta, S. and Chang, T.
Determination of 4,5-dihydro-6[4-(1H-

imidazol-1-yl)phenyl]-3(2H)-pyridazinone hydrochloride, a new cardiotonic, in plasma and urine by reversed-phase HPLC 336(1985)446
Hayn, M. and Esterbauer, H.
Separation and partial characterization of *Trichoderma reesei* cellulase by fast chromatofocusing 329(1985)379
Hearn, M.T.W., Hodder, A.N. and Aguilar, M.-I.
HPLC of amino acids, peptides and proteins. LXVI Investigations on the effects of chromatographic dwell in the reversed-phase HPLC separation of proteins 327(1985)47
Hearn, M.T.W., see Aguilar, M.-I. 327(1985)115
Hearn, M.T.W., see Grego, B. 336(1985)25
Heazlewood, R.L. and Lemass, R.W.J.
Simple HPLC assay for the routine monitoring of clonazepam in plasma 336(1985)229
Heckenberg, A.L. and Haddad, P.R.
Studies on sample preconcentration in ion chromatography. II. An automated, single pump preconcentration system for non-suppressed ion chromatography with conductivity detection 330(1985)95
Heckenberg, A.L., see Haddad, P.R. 346(1985)139
Hecker, R., Colpan, M. and Riesner, D.
HPLC of DNA restriction fragments 326(1985)251
Hegemann, J.H. and Linscheid, M.
Reversed-phase LC of protected oligonucleotide diesters 348(1985)286
Heimer, E.P., see Felix, A.M. 327(1985)359
Heimler, D., see Lepri, L. 347(1985)303
Heinemann, M.A., Whitaker, J.N. and Seyer, J.M.
Separation and isolation of human myelin basic protein peptides by reversed-phase HPLC in volatile buffers 329(1985)295
Heinemann, V., see Schwandt, P. 345(1985)145
Heiner, C.R., see Becker, C.R. 326(1985)293
Heinola, K., see Humppi, T. 331(1985)410
Heiny, M.E., see Takahashi, N. 326(1985)373
Heinz, F.X., see Winkler, G. 326(1985)113, 347(1985)83
Hellier, G., see Godin, J. 343(1985)424
Hem, G.K. van der, see Koopman, B.J. 337(1985)259
Hemmens, V., see Brown, A.S. 341(1985)139
Hendrix, D.L., see Guinn, G. 348(1985)123
Hendrix, P.Y., see Ansari, G.A.S. 346(1985)435
Henegouwen, R. Beijersbergen van, see Boeren, E.G. 349(1985)377
Hengstenberg, W., see Alpert, C.A. 326(1985)363
Hengy, H. and Kölle, E.-U.
Determination of levobunolol and dihydrolevobunolol in blood and urine by HPLC using fluorescence detection 338(1985)444
Hengy, H. and Kölle, E.-U.
Determination of gabapentin in plasma and urine by HPLC and pre-column labelling for UV detection 341(1985)473
Henion, J.D., see Skrabalak, D.S. 341(1985)261
Henn, R., see Battista, H.J. 345(1985)77
Henning, H.V., see Schneider, K. 345(1985)19
Henriks-Eckerman, M.-L. and Laijoki, T.
Ion-pair HPLC determination of dansylated aliphatic polyamines with fluorescence and UV detection 333(1985)220
Henry, J.A., see Storey, G.C.A. 341(1985)223
Herbranson, D.E., see Karnatz, N.N. 330(1985)420
Herilier, H., see Al Makdessi, S. 339(1985)25
Herington, A.C., see Kuffer, A.D. 336(1985)87
Hermann, C., Legrand, M., Geoffroy, P. and Fritig, B.
HPLC of proteins. Rapid chromatofocusing of plant enzymes 348(1985)167
Hermansson, J., Eriksson, M. and Nyquist, O.
Determination of (*R*)- and (*S*)-disopyramide in human plasma using a chiral α_1-acid glycoprotein column 336(1985)321
Hermens, W.A.J.J., see Overdiek, J.W.P.M. 341(1985)279
Hernádi, F.J., see Gál, S. 333(1985)239
Hernádi, F.J., see Kiss, L. 333(1985)244
Hernandez, L., see Barberan, F.A.T. 347(1985)443
Herraiz, M., see Reglero, G. 348(1985)327
Herranen, J., see Penttilä, I. 338(1985)265
Herregodts, P., Michotte, Y. and Ebinger, G.
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33
Herring, S.W., Shitanishi, K.T., Moody, K.E. and Enns, R.K.
Isolation of human factor VIII:C by preparative high-performance size-exclusion chromatography 326(1985)217
Herslöf, B., see Sotirhos, N. 331(1985)313
Hessey, G.A. II, see Vincek, W.C. 338(1985)438
Hester, R.D., see Vaidya, R.A. 333(1985)152
Heukeshoven, J. and Dernick, R.
Characterization of a solvent system for separation of water-insoluble poliovirus proteins by reversed-phase HPLC 326(1985)91
Heusler, H.
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273
Higuchi, S., Urano, C. and Kawamura, S.
Determination of plasma protein binding of propafenone in rats, dogs and humans by highly sensitive GC-MS 341(1985)305
Hilgers, H., see Schmitz, F.P. 346(1985)69
Hill, J.A. and Kitto, G.B.
HPLC assays of phenylpyruvate and phenylpyruvate oxidase 337(1985)397
Hill, J.A., see Abdullah, M. 347(1985)129
Himmel, M.E., see Baker, J.O. 346(1985)93
Himmelhoch, J.M., see Edwards, D.J. 344(1985)356

Hines, J.W., see Voyksner, R.D. 347(1985)137
Hinsvark, O., see Park, Y.H. 343(1985)359
Hiraga, Y. and Kinoshita, T.
HPLC analysis of guanidino compounds using ninhydrin reagent. II. Guanidino compounds in blood of patients on haemodialysis therapy 342(1985)269
Hiraoka, E., see Okamoto, T. 342(1985)35
Hirata, Y., see Fujimoto, C. 332(1985)47
Hirayama, C. and Ihara, H.
Novel GPC packing composed of cross-linked and porous spherical particles of poly-γ-methyl-L-glutamate 347(1985)357
Hirose, S., see Ishimitsu, T. 337(1985)239
Hirschberger, L.L., De la Rosa, J. and Stipanuk, M.H.
Determination of cysteinesulfinate, hypotaurine and taurine in physiological samples by reversed-phase HPLC 343(1985)303
Hitchcock, E.R., see Elworthy, P.M. 343(1985)109
Hjertén, S.
High-performance electrophoresis. Elimination of electroendosmosis and solute adsorption 347(1985)191
Hjertén, S. and Wu, B.-L.
Studies of fish zona pellucida by high-performance ion-exchange chromatography on agarose columns and free zone electrophoresis 341(1985)295
Hjertén, S. and Zhu, M.-D.
Micropreparative version of high-performance electrophoresis. The electrophoretic counterpart of narrow-bore HPLC 327(1985)157
Hjertén, S. and Zhu, M.-D.
Adaptation of the equipment for high-performance electrophoresis to isoelectric focusing 346(1985)265
Hoang, M.D. le, see Le Hoang, M.D. 346(1985)382
Hobo, T., Ishikawa, K. and Suzuki, S.
?In situ *generation of alkyl sulphides and alkyl disulphides* for GC identification 330(1985)131
Hoch, B., see Bialer, M. 337(1985)408
Hodder, A.N., see Aguilar, M.-I. 327(1985)115
Hodder, A.N., see Hearn, M.T.W. 327(1985)47
Hodges, R.S., see Cachia, P.J. 343(1985)315
Hodges, R.S., see Ingraham, R.H. 327(1985)77
Hodges, R.S., see Mant, C.T. 326(1985)349, 327(1985)147
Hogendoorn, E.A., see Goewie, C.E. 344(1985)157
Höhn, M., see Van der Greef, J. 343(1985)397
Hojabri, H. and Glennon, J.D.
Determination of cianopramine in human plasma by HPLC and GLC with UV, fluorescence and electron-capture detection 342(1985)97
Holdiness, M.R.
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321
Holland, M.L. and Ng, K.T.
Automated capillary GC assay using nitrogen-phosphorus detection for the determination of cetiedil in plasma 345(1985)178
Holler, J.S., see Brooks, J.B. 339(1985)243
Hollis, B.W. and Frank, N.E.
Solid phase extraction system for vitamin D and its major metabolites in human plasma 343(1985)43
Holmes, R.P., see Kou, I-L. 330(1985)339
Holt, G., see Rogers, M.E. 326(1985)163
Holt, N.J., see McCracken, M.S. 348(1985)221
Homandberg, G.A., Evans, D.B., Kramer, J. and Erickson, J.W.
Interaction between fluorescence-labeled fibronectin fragments studied by gel HPLC 327(1985)343
Honda, S., see Okamoto, Y. 350(1985)127
Hondo, T., see Hoshino, T. 332(1985)139
Hondo, T., see Sugiyama, K. 332(1985)107
Honeyman, T.W., see Ryan, P.J. 331(1985)177
Hoogmartens, J., see Kibwage, I.O. 330(1985)275, 346(1985)309
Hook, W.A. van, see Van Hook, W.A. 328(1985)333
Hoppel, C.L., see Minkler, P.E. 336(1985)271
Horai, Y. and Ishizaki, T.
Rapid and sensitive LC method for the determination of dapsone and monoacetyldapsone in plasma and urine 345(1985)447
Horáková, B., see Cee, A. 331(1985)202
Horie, M., see Baba, S. 341(1985)251
Horino, M., see Oyama, H. 338(1985)71
Horna, A., Pechová, H. and Churáček, J.
Chromatography of monomers. III. TLC separation of C_6-C_{18} straight-chain acrylates from corresponding methacrylates and alcohols 330(1985)439
Horna, A., Táborský, J., Churáček, J. and Dufka, O.
Chromatography of monomers. IV. GLC studies of C_1-C_6 *n*-alkyl and C_3-C_6 isoalkyl acrylates and their hydrogen halide and halogen addition derivatives 348(1985)141
Horstmann, M.
Simple HPLC method for rapid determination of nicotine and cotinine in urine 344(1985)391
Horváth, C., see El Rassi, Z. 326(1985)79
Horváth, C., see Frenz, J. 330(1985)1
Horváth, C., see Huang, J.-X. 330(1985)181
Horváth, G., see Szederkényi, F. 333(1985)404
Hoshino, T., Hondo, T., Senda, M., Saito, M. and Tohei, S.
Quantitative deconvolution of heavily fused chromatographic peaks of biological components using a multi-wavelength UV detector 332(1985)139
Hosoi, M., see Maeda, K. 345(1985)11
Hosoi, M., see Shiraishi, S. 338(1985)51
Hostettmann, K., see Schaufelberger, D. 346(1985)396
Houdi, A.A., see Mousa, S. 347(1985)405

Houdret, N., Lhermitte, M., Lalau, G., Izydorczak, J. and Roussel, P. Determination of thiopental and pentobarbital in plasma using HPLC 343(1985)437
Hough, B.J., see Moodie, I.M. 347(1985)179
Houston, J.B., see Adedoyin, A. 345(1985)192
Howes, L.G., Miller, S. and Reid, J.L. Simultaneous assay of 3,4-dihydroxyphenylethylene glycol and norepinephrine in human plasma by HPLC with electrochemical detection 338(1985)401
Hryhorczuk, L.M., see Novak, E.A. 339(1985)457
Hsia, C.J.C., see Wong, L.T. 338(1985)410, 341(1985)452
Hsieh, J.Y.-K., see Slobodzian, D.K. 338(1985)253, 345(1985)345
Hsu, F.-L., see Munavalli, S. 346(1985)463
Huang, D.Y., see Rokushika, S. 332(1985)15
Huang, J.-X., Bouvier, E.S.P., Stuart, J.D., Melander, W.R. and Horváth, C. HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. II. Retention behavior, quantitative structure-retention relationships and octanol-water partition coefficients 330(1985)181
Hubbard, J.W., see Lim, H.K. 328(1985)378
Huber, L. and Obbens, H. On-line analysis of sulphur compounds in natural gas with flame photometric detection 349(1985)465
Huber, L., see Goetz, H. 349(1985)357
Hubert, P., see Mathis, R. 347(1985)291
Huebner, F.R. and Bietz, J.A. Detection of quality differences among wheats by HPLC 327(1985)333
Hughes, D.E. Determination of α-2-deoxy-D-glucose in topical formulations by HPLC with UV detection 331(1985)183
Huhtikangas, A., see Penttilä, I. 338(1985)265
Hulshoff, A., see Lingeman, H. 336(1985)241
Hulst, A.G., see Wils, E.R.J. 330(1985)379
Hultin, T.A., Mehta, R.G. and Moon, R.C. Simple HPLC method for the separation of retinoids including N-(4-hydroxyphenyl)-all-*trans*-retinamide 341(1985)187
Humphreys, E., see Blatchford, C. 329(1985)281
Humppi, T. and Heinola, K. Synthesis and GC-MS determination of polychlorinated dibenzo-*p*-dioxins and related compounds in the technical chlorophenol formulation Ky-5 331(1985)410
Hung, C.T., see Schaaf, L.J. 342(1985)303
Hunt, J.H., see Cholerton, T.J. 333(1985)178
Hurtig, H., see Beers, M.F. 336(1985)380
Hurtubise, R.J., see Cooper, H.A. 328(1985)81
Hurtubise, R.J., see Ruckmick, S.C. 331(1985)55
Husain, S., Akella, K.S.R. and Narsimha, R. Analysis of some chlorinated pyridine isomers in mixtures by reversed-phase HPLC 348(1985)309
Hušek, P., see Mareš, P. 350(1985)87
Husseini, H.S. and Balzer, H.O. 1-DOPA decarboxylase and dopamine-β-hydroxylase isolated from sympathetic splenic nerves estimated by sodium dodecyl sulphate polyacrylamide gel electrophoresis with and without mercaptoethanol 329(1985)300
Hutchens, T.W., Dunaway, H.E. and Besch, P.K. High-performance chromatofocusing of steroid receptor proteins in the presence and absence of steroid. Investigation of steroid-dependent alterations in surface charge heterogeneity 327(1985)247
Hutchens, T.W., see Dunaway, H.E. 327(1985)221
Hutchings, A.L., see Littlewood, T.J. 336(1985)434
Hutte, R.S., see Sievers, R.E. 349(1985)395
Hütter, R., see Grossenbacher, H. 331(1985)161
Huysmans, M.B. and Spicer, W.J. Assessment of head-space GLC for the rapid detection of growth in blood cultures 337(1985)223
Hwang, D.-G., see Yinon, J. 339(1985)127
Hyder, S.M., Wiehle, R.D., Brandt, D.W. and Wittliff, J.L. High-performance hydrophobic-interaction chromatography of steroid hormone receptors 327(1985)237
Hyland, K. Estimation of tetrahydro, dihydro and fully oxidised pterins by HPLC using sequential electrochemical and fluorometric detection 343(1985)35
Hyun, M.H., see Pirkle, W.H. 328(1985)1
Hyver, K.J., Parcher, J.F., Johnson, D.M. and Lin, P.J. Adsorption of mixtures of acetone and benzene vapors on graphitized carbon black 328(1985)63

I

Iacobucci, G.A., see Chamblee, T.S. 330(1985)141
Ichihashi, Y., see Yoshida, A. 343(1985)155
Igarashi, T., see Inouye, K. 327(1985)301
Ihara, H., see Hirayama, C. 347(1985)357
Ihrke, M., see Eppert, G. 350(1985)471
Ijitsu, T., see Oka, K. 339(1985)253
Ikenaka, T., see Omichi, K. 336(1985)368
Ikuno, Y., Maoka, T., Shimizu, M., Komori, T. and Matsuno, T. Direct diastereomeric resolution of carotenoids. II. All ten stereoisomers of tunaxanthin (ε,ε-carotene-3,3'-diol) 328(1985)387

Imai, Y., see Hayashi, M. 338(1985)195
Improta, C., see Nota, G. 347(1985)362
Imre, L., see Juvancz, Z. 337(1985)121
Ingalls, S.T., see Minkler, P.E. 336(1985)271
Ingraham, R.H., Lau, S.Y.M., Taneja, A.K. and Hodges, R.S.
Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77
Inoue, T., Ishiwata, H., Yoshihira, K. and Tanimura, A.
HPLC determination of melamine extracted from cups made of malemine resin 346(1985)450
Inoue, T. and Niwaguchi, T.
Determination of nitrazepam and its main metabolites in urine by TLC and direct densitometry 339(1985)163
Inouye, H., see Kuwata, K. 332(1985)29
Inouye, K., Nakamura, K., Mitoma, Y., Matsumoto, M. and Igarashi, T.
Application of a new ion exchanger TSK-GEL DEAE-5PW, to the purification of Cu,Zn-superoxide dismutase of bovine erythrocytes 327(1985)301
Ioffe, B.V. and Zenkevich, I.G.
New methods of interpretation of chromatographic-MS data 349(1985)385
Irino, K., see Goto, M. 346(1985)167
Ishida, J., Kai, M. and Ohkura, Y.
Determination of *p*-hydroxybestatin in human serum by HPLC using fluorescence detection 344(1985)267
Ishida, J., see Kai, M. 345(1985)259
Ishii, D., Watanabe, K., Asai, H., Hashimoto, Y. and Takeuchi, T.
Advanced microcolumn techniques in LC. Use of glass-lined stainless-steel tubing columns 332(1985)3
Ishii, D., see Goto, M. 346(1985)167
Ishii, D., see Ito, Y. 346(1985)161
Ishii, D., see Takeuchi, T. 331(1985)99
Ishii, I., Takahashi, N., Kato, S., Akamatsu, N. and Kawazoe, Y.
HPLC analysis of changes of asparagine-linked oligosaccharides in regenerating rat liver 345(1985)134
Ishii, S.-I., see Shimura, K. 350(1985)265
Ishii, Y., see Chikuma, T. 348(1985)205
Ishikawa, K., see Hobo, T. 330(1985)131
Ishimitsu, T. and Hirose, S.
Simultaneous assay of 3,4-dihydroxyphenylalanine, catecholamines and O-methylated metabolites in human plasma using HPLC 337(1985)239
Ishioka, N., see Takahashi, N. 326(1985)407
Ishiwata, H., see Inoue, T. 346(1985)450
Ishiwatari, R., see Takada, H. 346(1985)281
Ishizaka, Y., see Hatake, K. 344(1985)339
Ishizaki, T., see Horai, Y. 345(1985)447
Ishizaki, T., see Kushida, K. 338(1985)131
Ishizu, H., see Ageta, T. 343(1985)186
Ito, K., see Kurosawa, N. 343(1985)379
Ito, M., see Takata, Y. 332(1985)117
Ito, M., see Tsukida, K. 331(1985)265
Ito, N., Noguchi, K., Kazama, M., Shimura, K. and Kasai, K.-I.
Separation of human Glu-plasminogen, Lys-plasminogen and plasmin by high-performance affinity chromatography on Asahipak GS gel coupled with *p*-aminobenzamidine 348(1985)199
Ito, N., Noguchi, K., Shimura, K. and Kasai, K.-I.
High-performance affinity chromatography of trypsins on Asahipak GS-gel coupled with *p*-aminobenzamidine 333(1985)107
Ito, Y.
Interaction of synthetic glycoproteins with immobilized lectins 330(1985)392
Ito, Y., Takeuchi, T., Ishii, D. and Goto, M.
Direct coupling of micro HPLC with fast atom bombardment MS 346(1985)161
Ito, Y., see Sagara, K. 328(1985)289
Ito, Y., see Sandlin, J.L. 348(1985)131
Itoh, Y., see Cooper, E.H. 327(1985)179
Itokawa, H., see Sagara, K. 328(1985)289
Itokawa, Y., see Kimura, M. 332(1985)181
Itsikson, L.B., Berezkin, V.G. and Haken, J.K.
Soviet polysiloxane stationary phases for GC (Review) 334(1985)1
Ivashkiv, E., see Jemal, M. 345(1985)299
Iversen, T., see Hagen, B. 345(1985)173
Iwaeda, T., see Yamatodani, A. 344(1985)115
Iwai, K., see Kawada, T. 329(1985)99
Iwamori, M., see Harada, T. 337(1985)187
Iwasaki, M., see Hamada, C. 341(1985)426
Iwasaki, S., Tanaka, H., Nakazawa, K. and Arima, M.
Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 341(1985)182
Iwashita, H., see Antoku, Y. 342(1985)359
Iwata, T., Yamaguchi, M., Hara, S., Nakamura, M. and Ohkura, Y.
Determination of total ascorbic acid in human serum by HPLC with fluorescence detection 344(1985)351
Iyengar, J.R., see Lawrence, J.F. 350(1985)237
Izydorczak, J., see Houdret, N. 343(1985)437

J

Jackson, G.E.D., see Young, N.M. 336(1985)397
Jackson, P.E. and Haddad, P.R.
Occurrence and origin of system peaks in non-suppressed ion chromatography of inorganic anions with indirect UV absorption detection 346(1985)125
Jackson, P.E., see Haddad, P.R. 346(1985)139
Jackson, P.S. and Gurley, L.R.
Analysis of nucleoproteins by direct

injection of dissolved nuclei or chromosomes into a HPLC system 326(1985)199
Jacob, K., Sommer, W., Meyer, H.D. and Vogt, W.
Ion-pair HPLC separation of porphyrin isomers 349(1985)283
Jacobson, K.B., see Ferre, J. 350(1985)389
Jadczak, J., see Walendziak, L. 331(1985)193
Jamali, F., see Torok-Both, G.A. 344(1985)372
James, S. and Bennett, H.P.J.
Use of reversed-phase and ion-exchange batch extraction in the purification of bovine pituitary peptides 326(1985)329
Jamet, G., see Staveris, S. 338(1985)79, 339(1985)97
Janda, V. and Marha, K.
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186
Janjić, T.J., Tešić, Ž.L., Malinar, M.J., Radivojša, P.N. and Ćelap, M.B.
Phenol effect in PC of metal complexes 331(1985)273
Jansen, E.H.J.M., Van Blitterswijk, H., Zoontjes, P.W., Both-Miedema, R. and Stephany, R.W.
Selectivity of a diol phase HPLC system in trace analysis of anabolic compounds 347(1985)375
Jansen, E.H.J.M., Zoontjes, P.W., Both-Miedema, R., Van Blitterswijk, H. and Stephany, R.W.
Screening of parts-per-billion levels of dietyhlstilbestrol in bovine urine by HPLC with UV detection 347(1985)379
Jansen, G., see Koopman, B.J. 337(1985)259
Janson, J.-C., see Andersson, T. 326(1985)33
Janssen, G., see Kibwage, I.O. 346(1985)309
Janus, E.D., see Poll, D. 343(1985)149
Jardine, L., see Labadarios, D. 339(1985)366
Jarry, C.H., see Demotes-Mainard, F.M. 336(1985)438, 342(1985)234
Jarvis, C.A., see Bosin, T.R. 341(1985)287
Jastorff, B., see Braumann, T. 329(1985)321, 350(1985)105
Javors, M.A., Bowden, C.L. and Maas, J.W.
3-Methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, and homovanillic acid in human cerebrospinal fluid. Storage and measurement by reversed-phase HPLC and coulometric detection using 3-methoxy-4-hydroxyphenyllactic acid as an internal standard 336(1985)259
Jaynes, P.K., see Chou, P.P. 341(1985)167
Jee, M.H., see Ritchie, A.S. 329(1985)273
Jefford, C.W., Grant, H.G., Kohmoto, S. and Boukouvalas, J.
Determination of the thermal stability of some 1,2-dioxetanes by HPLC 347(1985)183
Jehl, F., Gallion, C., Debs, J., Brogard, J.M., Monteil, H. and Minck, R.
HPLC method for determination of ciprofloxacin in biological fluids 339(1985)347
Jelínková, M., see Kuchař, M. 347(1985)335
Jemal, M. and Cohen, A.I.
Determination of S-benzoyl captopril in human urine by capillary GC with electron-capture detection 342(1985)186
Jemal, M., Ivashkiv, E., Ribick, M. and Cohen, A.I.
Determination of SQ 27,519, the active phosphinic acid-carboxylic acid of the prodrug SQ 28,555, in human serum by capillary GC with nitrogen-phosphorus detection after a two-step derivatization 345(1985)299
Jenkins, K.M., see Tsuji, K. 333(1985)365
Jensen, D.E., see Cortes, H.J. 349(1985)55
Jeppsson, J.-O., Lilja, H. and Johansson, M.
Isolation and characterization of two minor fractions of α_1-antitrypsin by HPLC chromatofocusing 327(1985)173
Jikumaru, N., see Ohta, K. 350(1985)292
Jindal, S.P., see Bagchi, S.P. 344(1985)362
Jinno, K., see Fujimoto, C. 332(1985)47
Jirsa, M., see Kotal, P. 333(1985)141
Jizba, J.V., Přikrylová, V. and Lipavská, H.
Preparative chromatography of epimers and anomers of daunomycin derivatives on Sephadex LH-20 329(1985)193
Joel, S.P., see Harvey, V.J. 339(1985)419
Joelsson, M., see Johansson, G. 331(1985)11
Johansson, B. and Fromark, I.
Determination of meprobamate as an *n*-butylboronate ester derivative in serum by GLC 341(1985)462
Johansson, B.-L. and Drevin, I.
Three independent methods for quantitative determination of octyl covalently coupled to Sepharose CL-4B 346(1985)255
Johansson, B.-L. and Ellström, C.
Column lifetime of a new agarose medium for high-performance gel filtration chromatography at basic pH 330(1985)360
Johansson, G., Joelsson, M., Olde, B. and Shanbhag, V.P.
Affinity partitioning of biopolymers and membranes in Ficoll-dextran aqueous two-phase systems 331(1985)11
Johansson, M., see Jeppsson, J.-O. 327(1985)173
Johns, E.A., see Cooper, E.H. 327(1985)179
Johns, R.B., see Strachan, M.G. 329(1985)65
Johnson, D.M., see Hyver, K.J. 328(1985)63
Johnson, M., see Catlow, D.A. 328(1985)167
Johnsson, H., see Österdahl, B.-G. 337(1985)151
Johnston, A., see Harvey, V.J. 339(1985)419
Jolley, K.W., see Bryce, T.A. 344(1985)397
Jolley, K.W., see Burrows, J.L. 344(1985)187
Jolly, J., see Ellis, B. 326(1985)157
Jones, A.J., see O'Keefe, J.H. 336(1985)73
Jones, K. and Malcolme-Lawes, D.J.
High-performance UV absorption detector for LC. I. Preliminary experiments 329(1985)25
Jones, M., see Eggert, F.M. 333(1985)123
Jones, R.L., see Adams, R.F. 336(1985)125
Jong, A.P.J.M. de, see Kok, R.M. 343(1985)59

Jong, J.W. de, see Harmsen, E. 337(1985)201
Jørgensen, P.L., see Grisham, C.M. 333(1985)301
Joseph, M.H.
Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370
Joseph, M.J., see DiBella, E.E. 328(1985)101
Josephson, M.W., see Tsai, M.Y. 341(1985)1
Julien, C., Rodriguez, C., Cuisinaud, G., Bernard, N. and Sassard, J.
Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51
Jung, D. and Zafar, N.U.
Micro HPLC assay of acetaminophen and its major metabolites in plasma and urine 339(1985)198
Jung, D., see Shah, A. 344(1985)408
Jung, L., see Staveris, S. 338(1985)79, 339(1985)97
Jurczak, J., see Debowski, J. 329(1985)206
Jürgens, D., Shalaby, F.Y.Y.I. and Fehrenbach, F.J.
Purification and characterization of CAMP-factor from *Streptococcus agalactiae* by hydrophobic interaction chromatography and chromatofocusing 348(1985)363
Jurlina, J.L. and Paxton, J.W.
Determination of N-5-dimethyl-9-[(2-methoxy-4-methylsulfonylamino)phenylamino]-4-acridinecarboxamide in plasma by HPLC 342(1985)431
Justice, J.B. Jr., see Alexander, L.R. 342(1985)1
Juvancz, Z., Ledniczky, M., Imre, L., Bihari, M. and Vajda, M.
Identification of 1,3-diphenyl-2-aminopropane metabolites by GC, GC with Fourier transform IR spectroscopy and GC-MS 337(1985)121

K

Kabaya, Y., see Mizutani, H. 331(1985)371
Kabra, P.M. and Nzekwe, E.U.
LC analysis of clonazepam in human serum with solid-phase (Bond-Elut) extraction 341(1985)383
Kacprowicz, A.T., Fullinfaw, R.O. and Bury, R.W.
HPLC measurement of verapamil in plasma using a diol column 337(1985)412
Kadam, A.N., see Bendre, S.B. 346(1985)359
Kadofuku, T. and Sato, T.
Detection of the changes in cellular proteins in regenerating rat liver by high-resolution two-dimensional electrophoresis 343(1985)51
Kafil, J.B. and Last, T.A.
LC with voltammetric detection for quantitation of phenolic acids 348(1985)397
Kageyu, A., see Tanaka, Y. 347(1985)275
Kai, M., Miura, T., Ishida, J. and Ohkura, Y.
HPLC method for monitoring leupeptin in mouse serum and muscle by pre-column fluorescence derivatization with benzoin 345(1985)259
Kai, M., see Ishida, J. 344(1985)267
Kaiser, N.F., see Becker, C.R. 326(1985)293
Kaliszan, R., see Knox, J.H. 349(1985)211
Kaliszan, R., see Ośmiałowski, K. 346(1985)53
Kalow, W., see Kruck, T.P.A. 341(1985)123
Kalpaxis, D.L.
Comparative study of affinity chromatography of components of the hyaluronate-proteoglycan complex to immobilized hyaluronate 350(1985)227
Kaluzny, M.A., see Epps, D.E. 343(1985)143
Kameyama, M., see Yoshizawa, I. 338(1985)404
Kamga, R., see Bossard, A. 330(1985)400
Kamino, M., see Funazo, K. 346(1985)215
Kammer, J., see Schweer, H. 338(1985)273
Kamminga, D., see Brinkman, U.A.T. 330(1985)375
Kamperman, G. and Kraak, J.C.
Simple and fast analysis of adrenaline and noradrenaline in plasma on microbore HPLC columns using fluorimetric detection 337(1985)384
Kanamori, N., see Okamoto, T. 342(1985)35
Kaneda, N. and Nagatsu, T.
Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 341(1985)23
Kaneda, N., Noro, Y. and Nagatsu, T.
Highly sensitive assay for acetylcholinesterase activity by HPLC with electrochemical detection 344(1985)93
Kangas, J.
GC and MS identification of sulphur gases in kraft digestion plants 346(1985)405
Kanmera, T., see Chaiken, I.M. 336(1985)63
Kaplan, E. and Ansari, K.
HPLC as a tool for identification of linolenic acid hydroperoxide prepared with soybean lipoxygenase 350(1985)435
Kapur, P.A., Law, T. and Watson, E.
Simultaneous quantitation of verapamil, norverapamil, and N-dealkylated metabolites in human plasma following oral administration 337(1985)160
Karaiskakis, G., see Dalas, E. 348(1985)339
Karaiskakis, G., see Katsanos, N.A. 349(1985)369
Karasek, F.W. and Tong, H.Y.
Semi-preparative HPLC analysis of complex organic mixtures 332(1985)169
Karasek, F.W., see Tong, H.Y. 350(1985)27

Karger, B.L., see Miller, N.T. 326(1985)45
Kärkkäinen, S.
HPLC determination of sotalol in biological fluids 336(1985)313
Karlaganis, G., see Baktir, G. 339(1985)192
Karlaganis, G., see Wahlländer, A. 338(1985)369
Karmen, A., Malikin, G., Freundlich, L. and Lam, S.
High-sensitivity radioassay of chromatographic effluents. Automatic fraction collector/concentrator for quantitative autoradiography 349(1985)267
Karmen, A., see Smith, S.L. 336(1985)351
Karnatz, N.N., Baaske, D.M., Herbranson, D.E. and Eliason, M.S.
HPLC method for the determination of esmolol hydrochloride in solutions and parenteral formulations 330(1985)420
Karnes, H.T., Riley, C.M., Curry, S.H. and Schulman, S.G.
Analysis of N-benzoyl-L-tyrosyl-*p*-aminobenzoic acid (bentiromide) metabolites in urine by ion-pair HPLC 338(1985)377
Karrer, G., see Grob, K. Jr. 334(1985)129
Kasai, K.-I., see Ito, N. 333(1985)107, 348(1985)199
Kasai, K.-I., see Shimura, K. 350(1985)265
Kasai, M., see Yasukawa, K. 332(1985)287
Kastner, F., see Rimböck, K.-H. 329(1985)307
Kasuya, Y., see Minagawa, K. 343(1985)231
Kasuya, Y., see Shinohara, Y. 338(1985)281
Kataoka, H., Ohnishi, N. and Makita, M.
Electron-capture GC of taurine as its N-pentafluorobenzoyl di-*n*-butylamide derivative 339(1985)370
Kato, S. and Yamamoto, K.
GC separation of monomethyl ethers of dihydroxycoumarin isomers 333(1985)175
Kato, S., see Ishii, I. 345(1985)134
Kato, T., see Chikuma, T. 348(1985)205
Kato, Y., Kitamura, T. and Hashimoto, T.
Resin-based support for reversed-phase chromatography of proteins 333(1985)93
Kato, Y., Kitamura, T. and Hashimoto, T.
Preparative high-performance hydrophobic interaction chromatography of proteins on TSKgel Phenyl-5PW 333(1985)202
Kato, Y., Matsuda, T. and Hashimoto, T.
New gel permeation column for the separation of water-soluble polymers 332(1985)39
Kato, Y., see Nakamura, K. 333(1985)29
Kato, Y., see Ozaki, H. 332(1985)243
Katsanos, N.A., Karaiskakis, G. and Agathonos, P.
Measurement of activity coefficients by reversed-flow GC 349(1985)369
Katsanos, N.A., see Dalas, E. 348(1985)339
Katsura, K., see Kawada, T. 329(1985)99
Kauffman, R., see Stout, M. 342(1985)424
Kauffman, R.E., see Aravind, M.K. 344(1985)428
Kawada, T., Watanabe, T., Katsura, K., Takami, H. and Iwai, K.
Formation and metabolism of pungent principle of *Capsicum* fruits. XV. Microdetermination of capsaicin by HPLC with electrochemical detection 329(1985)99
Kawahara, K., see Ohta, K. 350(1985)292
Kawai, S., see Nishioka, R. 342(1985)89
Kawamura, S., see Higuchi, S. 341(1985)305
Kawanabe, K.
Application of thin-layer stick chromatographic identification test methods to drugs contained in preparations in the Pharmacopoeia of Japan 333(1985)115
Kawata, S., see Hayashi, M. 338(1985)195
Kawazoe, Y., see Ishii, I. 345(1985)134
Kay, C.M., see Cachia, P.J. 343(1985)315
Kazama, M., see Ito, N. 348(1985)199
Keeler, R.F., see Browne, C.A. 336(1985)211
Kehr, J. and Chavko, M.
Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column LC under isocratic conditions 345(1985)267
Keller, P.L., see Link, G.W. Jr. 331(1985)253
Keller, R.K., Brennan, S., Cafmeyer, N. and Adair, W.L.
Quantitation of tissue levels of dolichyl pyrophosphate by HPLC on silica 343(1985)413
Kelly, T.J., see Gaffney, J.S. 347(1985)121
Kempen, G.M.J. van, see Pennings, E.J.M. 341(1985)172
Kemper, K., Hagemeier, E., Boos, K.-S. and Schlimme, E.
Direct clean-up and analysis of urinary catecholamines 336(1985)374
Kennedy, M.S., see Gmur, D.J. 344(1985)422
Keresztes, P., see Csató, E. 348(1985)280
Kerns, E.H., see Gammans, R.E. 339(1985)303, 345(1985)285
Key, J.A., see Gill, R. 346(1985)423
Khachik, F. and Beecher, G.R.
Decapreno-β-carotene as an internal standard for the quantification of the hydrocarbon carotenoids by HPLC 346(1985)237
Khalil, S.N., see Shao, McN.J. 345(1985)184
Khandelwal, J.K., Kline, T. and Green, J.P.
Measurement of imidazoleacetic acid in urine by GC-MS 343(1985)249
Kharbade, B.V. and Agrawal, O.P.
Identification of natural red dyes in old Indian textiles. Evaluation of TLC systems 347(1985)447
Kholy, A.M., see Brooks, J.B. 339(1985)243
Kiba, N., Mizuno, T. and Furusawa, M.
Determination of catecholamines by HPLC with catalytic photometric detection 329(1985)147
Kiba, N., Takamatsu, M. and Furusawa, M.
Determination of anthraquinones in pulping materials by HPLC using online post-column derivatization 328(1985)309
Kibwage, I.O., Janssen, G., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.
Isolation of erythromycins and related

substances from fermentation residues of *Streptomyces erythreus* by HPLC on silica gel 346(1985)309
Kibwage, I.O., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.
Separation of erythromycin and related substances by HPLC on poly(styrene-divinylbenzene) packing materials 330(1985)275
Kida, K., see Sogami, M. 332(1985)19
Kilts, C.D., Dew, K.L., Ely, T.D. and Mailman, R.B.
Quantification of *R*-(+)-7-chloro-8-hydroxy-1-phenyl-2,3,4,5-tetrahydro-1H-3-methyl-3-benzazepine in brain and blood by use of reversed-phase HPLC with electrochemical detection 342(1985)452
Kilts, C.D., see Patrick, K.S. 343(1985)329
Kimura, M. and Itokawa, Y.
Determination of thiamine and its phosphate esters in human and rat blood by HPLC with post-column derivatization 332(1985)181
Kimura, T., see Kumagaye, K.Y. 327(1985)327
King, S.-Y.P. and Fung, H.-L.
Examination of the in vitro degradation of [^{14}C]pentaerythritol tetranitrate in rat and human blood with an improved thin-layer radiochromatographic procedure 343(1985)129
Kinoshita, H., see Tanaka, N. 332(1985)57
Kinoshita, K., see Taniguchi, N. 339(1985)394
Kinoshita, M., see Okazaki, M. 336(1985)151
Kinoshita, T., see Hiraga, Y. 342(1985)269
Kinoshita, T., see Yoshida, T. 348(1985)425
Kinsella, R.A. Jr., see Francis, F.E. 336(1985)361
Kirk, D.N., see Coldwell, R.D. 338(1985)289
Kirkland, J.J., see Schallinger, L.E. 342(1985)67
Kirret, O., see Murel, A. 347(1985)325
Kirukawa, M., see Hanai, T. 349(1985)323
Kishi, H., see Okamoto, T. 342(1985)35
Kishi, T., see Okamoto, T. 342(1985)35
Kiss, L., Toth-Martinez, B.L., Gál, S. and Hernádi, F.J.
Use of chromatofocusing for separation of β-lactamases. VI. Comparative studies on two chromatofocusing polybuffer exchanger matrices with the R46 plasmid coded β-lactamase carried by the *Escherichia coli* K12 J5-3 strain 333(1985)244
Kiss, L., see Gál, S. 333(1985)239
Kitada, Y., Mizobuchi, M., Ueda, Y. and Nakazawa, H.
Analysis of isoflavones in *Puerariae radix* by HPLC with amperometric detection 347(1985)438
Kitagawa, M., see Fujimura, K. 350(1985)371
Kitamura, T., see Kato, Y. 333(1985)93, 333(1985)202
Kitto, G.B., see Hill, J.A. 337(1985)397
Klaus, R.
Bile acids as an example to demonstrate the methodological aspects of the quantitative HPTLC analysis of fluorescent and non-fluorescent compounds (in German) 333(1985)276
Klee, C.B., see Manalan, A.S. 326(1985)387
Kleemeier, T., see Haldna, Ü. 350(1985)296
Klein, C.M. and Lau-Cam, C.A.
Simple TLC method for the investigation of six related compounds in flurazepam hydrochloride and its capsules 350(1985)273
Klesper, E., see Schmitz, F.P. 346(1985)69
Klett, R.P. and Chovan, J.P.
Modification of a new HPLC method for bleomycin to separate epi-, iso-, desamido-, and unmodified analogues 337(1985)182
Kleyn, E. van der, see Wuis, E.W. 337(1985)341
Kline, T., see Khandelwal, J.K. 343(1985)249
Kloer, H.U., see McConathy, W.J. 342(1985)47
Klüppelberg, U., see Swobodnik, W. 339(1985)263
Knapp, G., see Minagawa, K. 343(1985)231
Knausz, D., see Csató, E. 348(1985)280
Knighton, D.R., Harding, D.R.K., Napier, J.R. and Hancock, W.S.
Purification of synthetic lipid associating peptides and the monitoring of the deformylation of N^{in}-formyltryptophan by reversed-phase HPLC 347(1985)237
Knödgen, B., see Seiler, N. 339(1985)45
Knodgen, B., see Seiler, N. 341(1985)11
Knoll, R. and Brandl, M.
Rapid and simple method for the routine determination of dobutamine in human plasma by HPLC 345(1985)425
Knopf, S., see Edwards, D.J. 344(1985)356
Knowles, J.A., see Leelavathi, D.E. 339(1985)105
Knowlton, S., see Schallinger, L.E. 342(1985)67
Knox, J.H. and Kaliszan, R.
Theory of solvent disturbance peaks and experimental determination of thermodynamic dead-volume in column LC 349(1985)211
Knudsen, P., see Larsen, N.-E. 341(1985)244
Kobayashi, A., Sugita, S. and Nakazawa, K.
HPLC determination of imipramine and desipramine in human serum 336(1985)410
Kobayashi, A., see Yonezawa, K. 339(1985)219
Kobayashi, J., see Ganno, S. 332(1985)275
Kočan, A., see Matisová, E. 346(1985)177
Köck, A. and Luger, T.A.
HPLC separation of distinct epidermal cell-derived cytokines 326(1985)129
Kocourek, J., see Malý, P. 347(1985)343
Kodama, H., see Ageta, T. 343(1985)186
Kodama, H., see Mikasa, H. 343(1985)179
Koffel, J.C., see Staveris, S. 338(1985)79, 339(1985)97
Kohlhepp, E.A., see Egorin, M.J. 343(1985)196
Kohmoto, S., see Jefford, C.W. 347(1985)183
Kohse, K.P., Graser, T.A., Godel, H.G., Rössle, C., Franz, H.E. and Fürst, P.
HPLC determination of plasma free trimethyllysine in humans 344(1985)319
Koizumi, K., Kubota, Y., Okada, Y. and Utamura, T.

Microanalyses of β-cyclodextrin in plasma by HPLC 341(1985)31
Koizumi, Y., see Nakamura, Y. 333(1985)83
Kojima, K., see Suzuki, H. 341(1985)176
Kok, R.M., De Jong, A.P.J.M., Van Groeningen, C.J., Peters, G.J. and Lankelma, J.
Highly sensitive determination of 5-fluorouracil in human plasma by capillary GC and negative ion chemical ionization MS 343(1985)59
Kolanitch, R., see Kremer, R. 344(1985)313
Kolbe, H.V.J., Lu, R.C. and Wohlrab, H.
Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1
Kölle, E.U., see Goebel, K.-J. 345(1985)355
Kölle, E.-U., see Hengy, H. 338(1985)444, 341(1985)473
Komers, R., see Tameesh, A.H.H. 328(1985)207
Komori, T., see Ikuno, Y. 328(1985)387
Komuro, C., Ono, K., Shibamoto, Y., Nishidai, T., Takahashi, M. and Abe, M.
Rapid and simple method for quantitative determination of non-protein sulphydryls in mouse liver by reversed-phase HPLC 338(1985)209
Kondo, K., see Nishioka, R. 342(1985)89
Kondo, S., see Suzuki, H. 341(1985)341
König, W.A. and Sturm, U.
Determination of optical purity by enantioselective capillary GC: panthenol and related compounds 328(1985)357
König, W.A., see Gyllenhaal, O. 350(1985)328
Koopman, B.J., Jansen, G., Wolthers, B.G., Beukhof, J.R., Go, J.G. and Van der Hem, G.K.
Determination of inorganic sulphate in plasma by reversed-phase chromatography using UV detection and its application to plasma samples of patients receiving different types of haemodialysis 337(1985)259
Koordes, R.C.A., see Nielen, M.W.F. 330(1985)113
Kopanica, M., see Trojánek, A. 328(1985)127
Korach, K.S., see Parker, C.E. 344(1985)378
Kordač, V., see Kotal, P. 333(1985)141
Koren, E., see McConathy, W.J. 342(1985)47
Korhonen, I.O.O.
GLC analyses. XLIII. Retention increments for 2-chloro-, 2,2-dichloro- and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns 329(1985)43
Korhonen, I.O.O.
GLC analyses. XLIV. Benzoyl and pentafluorobenzoyl esters of lower saturated branched-chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 329(1985)359
Korhonen, I.O.O. and Lind, M.A.
GLC analyses. XLII. Retention behaviour of C_1-C_{12} *n*-alkyl esters of benzoic and pentafluorobenzoic acids on SE-30 and OV-351 capillary columns 328(1985)325
Kormos, L.S., see Minkler, P.E. 336(1985)271
Kościelski, T. and Sybilska, D.
Resolution of *cis*- and *trans*-dimethylcyclohexanes by partition GC through cyclodextrin complexes 349(1985)3
Kosh, J.W., see Dick, R.M. 347(1985)387
Kotake, A.N., Vogelzang, N.J., Larson, R.A. and Choporis, N.
New HPLC assay for plasma doxorubicin 337(1985)194
Kotal, P., Porsch, B., Jirsa, M. and Kordač, V.
Rapid isocratic HPLC of porphyrin esters on aminopropyl-bonded silica 333(1985)141
Kothari, P.J., see Boothe, T. 333(1985)269
Kou, I-L. and Holmes, R.P.
Analysis of 25-hydroxycholesterol in plasma and cholesterol-containing foods by HPLC 330(1985)339
Kovács, S., see Siklósi, A. 347(1985)393
Kovář, J.
Determination of alcoholic strength in alcoholic beverages by GLC. Precision and accuracy 333(1985)389
Kowalski, W.J.
Selectivity of complexing GC stationary phases containing lanthanide β-diketonates 349(1985)457
Kozloski, R.P.
Simple method of concentrating volatiles in water for GC analysis by vacuum distillation 346(1985)408
Kozma, M., see Rónai-Lukács, S. 338(1985)428
Kraak, J.C., see Kamperman, G. 337(1985)384
Kramer, J., see Homandberg, G.A. 327(1985)343
Kramer, P.A., see Dorrbecker, B.R. 336(1985)293
Kramlová, M., see Přistoupil, T.I. 346(1985)401
Kranz, T., see Angelo, H.R. 345(1985)413
Kraus, E., see Kuchař, M. 347(1985)335
Krause, W., see Lemm, U. 342(1985)393
Krauss, F. and Schmidt, A.
Simple separation of adenosine-5'-phosphosulphate, adenosine-5'-phosphoramidate and cyclic adenosine-5'-monophosphate nucleotides involved in adenosine-5'-phosphosulphate degradation 348(1985)296
Krawczyk, W., see Oszczapowicz, J. 330(1985)79
Kremer, R., Crawhall, J.C. and Kolanitch, R.
Rapid and reliable estimation of urinary free catecholamines in patients with pheochromocytoma: comparison with plasma catecholamines and vanillylmandelic acid excretion 344(1985)313
Krishnan, S. and Vijayalakshmi, M.A.
Purification of an acid protease and a serine carboxypeptidase from *Aspergillus niger* using metal-chelate affinity chromatography 329(1985)165
Kristian, P., see Podhradský, D. 328(1985)402

Kříž, J., Březina, M., Brosch, D., Vodička, L. and Duddeck, H.
Separation of isomeric halogenobicyclononene carbonitriles 347(1985)297
Kroschel, D.M., see Powis, G. 345(1985)402
Kroviarski, Y., see Bertrand, O. 346(1985)111
Kruck, T.P.A., Kalow, W. and McLachlan, D.R.C.
Determination of desferoxamine and a major metabolite by HPLC. Application to the treatment of aluminium-related disorders 341(1985)123
Krüger, G., see Berndt, H. 348(1985)275
Kruggel, W.G. and Lewis, R.V.
HPLC support for the baseline separation of all phenylthiohydantoin amino acids 342(1985)376
Kruh, J., see Riffe, A. 344(1985)332
Krull, I.S., see Colgan, S.T. 333(1985)349
Krupčík, J. and Bohov, P.
Use of equivalent chain lengths for the characterization of fatty acid methyl esters separated by linear temperature-programmed GC 346(1985)33
Krupčík, J., see Matisová, E. 346(1985)177
Krýsl, S. and Smolková-Keulemansová, E.
Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167
Kubo, I., see Hanke, F.J. 329(1985)395
Kubota, Y., see Koizumi, K. 341(1985)31
Kuchař, M., Kraus, E., Jelínková, M., Rejholec, V. and Miller, V.
TLC and HPLC in the evaluation of the lipophilicity of aryloxoalkanoic and arylhydroxyalkanoic acids 347(1985)335
Kuchař, M., Tomková, H., Rejholec, V. and Skalická, O.
Relationships between GLC behaviour and structure of arylaliphatic acids 333(1985)21
Kuchař, M., see Tomková, H. 329(1985)113
Kucharczyk, N. and Segelman, F.H.
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243
Kudo, S., see Akiyama, H. 338(1985)456
Kuffer, A.D. and Herington, A.C.
Partial purification of a specific inhibitor of the insulin-like growth factors by reversed-phase HPLC 336(1985)87
Kuhn, R.W., see Riemer, R.K. 338(1985)236
Kuhnz, W., Zimmermann, B. and Nau, H.
Improved separation of phospholipids by HPLC 344(1985)309
Kuijk, F.J.G.M. van, see Van Kuijk, F.J.G.M. 348(1985)241
Kumagaye, K.Y., Takai, M., Chino, N., Kimura, T. and Sakakibara, S.
Comparison of reversed-phase and cation-exchange HPLC for separating closely related peptides: separation of Asp^{76}-human parathyroid hormone (1-84) from Asn^{76}-human parathyroid hormone (1-84) 327(1985)327
Kumps, A., Genin-Ramakers, J. and Mardens, Y.
Simultaneous determination of anticonvulsant drugs and metabolites in plasma by HPLC 342(1985)469
Kun, E., see Hakam, A. 330(1985)287
Kunz, C., see Winkler, G. 326(1985)113, 347(1985)83
Kurauchi, O., see Taira, H. 339(1985)399
Kuribayashi, S., see Kurosawa, N. 343(1985)379
Kuroda, N., see Hamada, C. 341(1985)426
Kuroda, N., see Matsuoka, C. 341(1985)432
Kurosawa, N., Kuribayashi, S., Owada, E., Ito, K., Nioka, M., Arakawa, M. and Fukuda, R.
Determination of streptomycin in serum by HPLC 343(1985)379
Kurowski, M.
Simultaneous determination of ketanserin and ketanserinol in biological fluids using ion-pair LC and fluorometric detection 341(1985)208
Kushida, K. and Ishizaki, T.
Concurrent determination of valproic acid with other antiepileptic drugs by HPLC 338(1985)131
Kuśmierz, J., Maliński, E., Czerwiec, W. and Szafranek, J.
Kováts retention indices of high-molecular-weight monomethyl-, cyclopentyl-, cyclohexyl- and phenylalkanes 331(1985)219
Kuwamoto, T., see Okada, T. 350(1985)317
Kuwata, K., Era, S., Inouye, H., Sogami, M. and Sasaki, H.
Ion-exchange HPLC studies on sulphydryl-catalysed structural alterations of bovine mercaptalbumin 332(1985)29
Kuwata, K., see Sogami, M. 332(1985)19

L

Laarveld, B., see Misir, R. 331(1985)141
Labadarios, D., Shephard, G.S., Moodie, I.M., Jardine, L. and Botha, E.
Plasma amino acid analysis by GC. Removal of glucose interference 339(1985)366
Labadarios, D., see Moodie, I.M. 347(1985)179
Labadie, R.P., see Van den Berg, A.J.J. 329(1985)311
Labonia, N., see Carducci, C.N. 337(1985)91
Lacroix, C., Levert, P., Laine, G., Goulle, J.P. and Gringore, A.
Micro determination of 5-fluorocytosine by HPLC and fluorimetric detection (in French) 345(1985)436
Ladrón de Guevara, O., Estrada, G., Antonio, S., Alvarado, X., Guereca, L.,

Zamudio, F. and Bolívar, F.
Identification and isolation of human insulin A and B chains by HPLC 349(1985)91
Laduron, C., see Brassinne, C. 339(1985)445
Laffi, G., see Zipser, R.D. 339(1985)1
Laganà, A., Liberti, A., Politi, L., Scandurra, R. and Cerulli, N.
Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251
Lagerström, P.-O., see Ahnoff, M. 340(1985)73
Lahm, H.-W. and Stein, S.
Characterization of recombinant human interleukin-2 with micromethods 326(1985)357
Laijoki, T., see Henriks-Eckerman, M.-L. 333(1985)220
Laine, G., see Lacroix, C. 345(1985)436
Lalau, G., see Houdret, N. 343(1985)437
Laliberté, C., see Massé, R. 339(1985)11
Lam, L.K.T., Yee, C., Chung, A. and Wattenberg, L.W.
Use of silver nitrate impregnated silica cartridges in the separation of kahweol and cafestol esters by preparative LC 328(1985)422
Lam, S., see Karmen, A. 349(1985)267
Lambe, R., see Dadgar, D. 337(1985)47, 337(1985)136, 342(1985)315
Lambrechts, M. and Rasmussen, K.E.
Use of bonded-phase silica sorbents for rapid sampling of impurities in illicit amphetamine for HPLC analyses 331(1985)339
Lambros, T.J., see Felix, A.M. 327(1985)359
Landi, M., see Bianchi, G. 338(1985)230
Landry, D.A., see Cole, D.E.C. 337(1985)267
Lang, H.R.M., see Rizzi, A. 331(1985)33
Lang, J.F., see Smithers, J.A. 341(1985)232
Langenhove, H. van, see Van der Straeten, D. 331(1985)207
Langrené, S., see Léonil, J. 347(1985)316
Langseth-Manrique, K., Bredesen, J.E. and Greibrokk, T.
Rapid, isothermal GLC determination of nitroglycerine in plasma using an electron-capture detector. II 349(1985)421
Langvardt, P.W., see Freshour, N.L. 346(1985)376
Lankelma, J., see Kok, R.M. 343(1985)59
Lapalus, P., see Garraffo, R. 337(1985)416
Laperashvili, L.Y., see Andronikashvili, T.G. 331(1985)402
Larsen, B., see Gregory, V.M. 345(1985)140
Larsen, N.-E., Hansen, L.B. and Knudsen, P.
Quantitative determination of perphenazine and its dealkylated metabolite using HPLC 341(1985)244
Larson, R.A., see Kotake, A.N. 337(1985)194
Laruelle, L., see Daenens, P. 342(1985)79
LaRusso, N.F., see Tietz, P.S. 336(1985)249
Lassiani, L., see Farina, P. 345(1985)323
Last, T.A., see Kafil, J.B. 348(1985)397
Lau, P., see Whiton, R.S. 347(1985)109
Lau, S.Y.M., see Ingraham, R.H. 327(1985)77
Lau-Cam, C.A., see Klein, C.M. 350(1985)273
Lau-Cam, C.A., see Yuen, S. 329(1985)107
Läubli, T., see Grob, K. Jr. 333(1985)156
Laufen, H., see Räder, K. 344(1985)416
Laurance, A., see Bassene, E. 346(1985)428
Lauren, D.R. and Watkinson, J.H.
Elemental sulphur analysis using HPLC on 10-μm rigid polymer particles 348(1985)317
Law, T., see Kapur, P.A. 337(1985)160
Lawrence, J.F. and Iyengar, J.R.
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237
Lawson, G.J.
Rapid method for the measurement of methylprednisolone and its hemisuccinate in plasma and urine following "pulse therapy" by HPLC 342(1985)251
Lazure, C., Rochemont, J., Seidah, N.G. and Chrétien, M.
Novel approach to rapid and sensitive localization of protein disulfide bridges by HPLC and electrochemical detection 326(1985)339
Le Goffic, F., see Léonil, J. 347(1985)316
Le Hoang, M.D., Postaire, E., Prognon, P., Pradeau, D., Blondin, A. and Sauziere, J.
Separation and determination of the various constituents of troxerutin by HPLC. Application to the pharmaceutical control (in French) 346(1985)382
Le Moing, J.P., see Vajta, S. 343(1985)239
L'E Orme, M., see Edwards, G. 343(1985)224
Leboda, R., see Tracz, E. 346(1985)346
Leclercq, M., Fouillit, M., Panteix, G. and Guinet, R.
Reproducible measurement of amphotericin B in serum by HPLC in alkaline buffer 337(1985)423
Ledniczky, M., see Juvancz, Z. 337(1985)121
Lee, D.M., see McConathy, W.J. 342(1985)47
Lee, D.W., see Weber, A.F. 337(1985)434
Lee, M., Nohta, H., Ohkura, Y. and Yoo, B.
Determination of phenylethanolamine N-methyltransferase by HPLC with fluorescence detection 348(1985)407
Lee, M.L., see Fields, S.M. 349(1985)305
Lee, T.L., see Brooks, M.A. 345(1985)333
Lee, U.-S., see Tanaka, T. 328(1985)271
Leelavathi, D.E., Dressler, D.E., Soffer, E.F., Yachetti, S.D. and Knowles, J.A.
Determination of promethazine in human plasma by automated HPLC with electrochemical detection and by GC-MS 339(1985)105
Leenheer, A.P. de, see Nelis, H.J.C.F. 333(1985)381
Leeuwen, M.B. van, see Van Baal, J.M. 336(1985)422
Lefevere, M.F., see Nelis, H.J.C.F. 333(1985)381
Legator, M.S., see Rinkus, S.J. 342(1985)465
Legrand, M., see Hermann, C. 348(1985)167
Lehmkuhl, H.D., see Schmerr, M.J.F. 326(1985)225

Lehr, K.-H. and Damm, P.
Quantification of ciclopirox by HPLC after pre-column derivatization. An example of efficient clean-up using silica-bonded cyano phases 339(1985)451
Lehtonen, P.
Properties of six octadecylsilane stationary phases for the separation of dansylamides 330(1985)243
Leis, H.J., see Gleispach, H. 342(1985)245, 344(1985)11
Leluan, G., see Pons, J.-L. 337(1985)213
Lemar, M., see Dow, J. 344(1985)275
Lemass, R.W.J., see Heazlewood, R.L. 336(1985)229
Lemm, U., Tenczer, J., Baudisch, H. and Krause, W.
Antibody-mediated extraction of the main tetrahydrocannabinol metabolite, 11-nor-Δ^9-tetrahydrocannabinol-9-carboxylic acid, from human urine and its identification by GC-MS in the sub-nanogram range 342(1985)393
Lemmo, M., see Del Principe, D. 342(1985)285
Lempiäinen, M. and Mäkelä, A.-L.
Determination of proquazone and its *m*-hydroxy metabolite by HPLC. Clinical application: pharmacokinetics of proquazone in children with juvenile rheumatoid arthritis 341(1985)105
Lendero, L., see Obbens, H. 349(1985)425
Lennard, L.
Assay of 6-mercaptopurine in human plasma 345(1985)441
Lennard, M.S.
Quantitative analysis of metoprolol and three of its metabolites in urine and liver microsomes by HPLC 342(1985)199
Lennard, M.S. and Parkin, S.
Timolol determination in plasma and urine by HPLC with UV detection 338(1985)249
Léonil, J., Langrené, S., Sicsic, S. and Le Goffic, F.
Purification of D-amino acid oxidase apoenzyme by affinity chromatography on Cibacron Blue Sepharose 347(1985)316
Lepri, L., Desideri, P.G. and Heimler, D.
HPTLC of phenylethylamines and phenolic acids on silanized silica and on ammonium tungstophosphate 347(1985)303
Lester, D.C., see Nierenberg, D.W. 345(1985)275
Letoffé, J.M., see Claudy, P. 329(1985)331
Letoffé, J.M., see Gonnet, C. 330(1985)227
Levert, P., see Lacroix, C. 345(1985)436
Levillain, P., see Montagu, M. 331(1985)437
Levitt, M.D., see Doizaki, W.M. 329(1985)137
Levy, L.A., see Parker, C.E. 344(1985)378
Levy-Favatier, F., see Riffe, A. 344(1985)332
Lew, M., see Gupta, R.N. 342(1985)442, 344(1985)221
Lewin, M.J.M., see Robert, J.C. 338(1985)315
Lewis, N.G. and Yean, W.Q.
High-performance size-exclusion chromatography of lignosulphonates 331(1985)419
Lewis, P.A., Woodward, A.J. and Maddock, J.
Improved method for the determination of N-acetylcysteine in human plasma by HPLC 327(1985)261
Lewis, R.V., see Kruggel, W.G. 342(1985)376
Leykam, J.F., see McCourt, D.W. 327(1985)9
Lhermitte, M., Bernier, J.L., Mathieu, D., Mathieu-Nolf, M., Erb, F. and Roussel, P.
Colchicine quantitation by HPLC in human plasma and urine 342(1985)416
Lhermitte, M., see Houdret, N. 343(1985)437
Liberti, A., see Laganà, A. 345(1985)251
Lickl, E., see Ebermann, R. 347(1985)384
Liebich, H.M. and Först, C.
N-Methylation and N,N-dimethylation of amino acids. An artifact production in the analysis of organic acids using diazomethane as derivatizing agent 338(1985)33
Liebich, H.M. and Först, C.
Urinary excretion of N-acetylamino acids 338(1985)187
Liebich, H.M. and Pickert, A.
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25
Liebscher, G., see Eppert, G. 350(1985)471
Light, N., see Avery, N.C. 328(1985)347
Ligny, C.L. de, see Brandts, P.M. 333(1985)41
Lilja, H., see Jeppsson, J.-O. 327(1985)173
Lim, H.K., Sardessai, M., Hubbard, J.W. and Midha, K.K.
Enantiomeric resolution of *dl-threo*-methylphenidate, U.S.P. (Ritalin), by HPLC 328(1985)378
Lin, J.H., see Musson, D.G. 337(1985)363
Lin, K.-T., Momparler, R.L. and Rivard, G.E.
Sample preparation for the determination of 5-aza-2'-deoxycytidine in plasma by HPLC 345(1985)162
Lin, L., Saller, C.F. and Salama, A.I.
Rapid automated HPLC analysis of cyclic adenosine 3',5'-monophosphate. Synthesis in brain tissues 341(1985)43
Lin, P.J., see Hyver, K.J. 328(1985)63
Lind, M.A., see Korhonen, I.O.O. 328(1985)325
Lindberg, R.L.P.
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 341(1985)333
Lindblom, H., see Cooper, E.H. 327(1985)269
Linde, S., see Welinder, B.S. 348(1985)347
Lingeman, H., Haan, H.B.P. and Hulshoff, A.
Rapid and selective derivatization method for the nitrogen-selective detection of carboxylic acids in biological fluids prior to GC analysis 336(1985)241
Link, G.W., see Banes, A.J. 326(1985)419
Link, G.W. Jr., Keller, P.L., Stout, R.W. and Banes, A.J.
Effects of solutions used for storage of size-exclusion columns on subsequent chromatography of peptides and proteins 331(1985)253

Linscheid, M., see Hegemann, J.H. 348(1985)286
Lintz, W. and Uragg, H.
Quantitative determination of tramadol in human serum by GC-MS 341(1985)65
Lipavská, H., see Jizba, J.V. 329(1985)193
Lipinski, J., see Stan, H.-J. 349(1985)49
Littlewood, T.J., Hutchings, A.L., Bentley, D.P. and Spragg, B.P.
HPLC determination of etoposide in plasma using electrochemical detection 336(1985)434
Liu, Z. and Franklin, M.R.
HPLC separation and determination of 2-(diethylamino)ethyl diphenylpropylacetate and its metabolites from biological samples 342(1985)135
Livesey, G. and Edwards, E.T.E.
Quantification of branched-chain α-keto acids as quinoxalinols: importance of excluding oxygen during derivatization 337(1985)98
Ljungquist, C., see Sjöquist, B. 343(1985)1
Lloyd, J.B.F.
Microcolumn clean-up and recovery techniques for organic explosives compounds and for propellants traces in firearms discharge residues 330(1985)121
Lloyd, P. and Crabbe, M.J.C.
HPLC analysis of glucose, fructose, sorbitol and low levels of the aldose reductase inhibitor sorbinil in human lens and plasma 343(1985)402
Llyod, C.W., Martin, W.J., Nagle, J. and Hauser, A.R.
Determination of cimetidine and metabolites in plasma by reversed-phase HPLC radial compression technique 339(1985)139
Loadman, P.M., see McDermott, B.J. 338(1985)335
Lockwood, P.A., see D'Agostino, P.A. 347(1985)257
Locniskar, A., Greenblatt, D.J. and Ochs, H.R.
Simplified GC assay of underivatized nitrazepam in plasma 337(1985)131
Logan, R.I. and Rivett, D.E.
Investigation of the synthesis of tryptathionine using HPLC 348(1985)421
Lohmann, D., see Francotte, E. 347(1985)25
Lomić, S., see Petrović, S.M. 348(1985)49
Loon, G.R. van, see Mousa, S. 347(1985)405
Lorenschat, B., see Schmitz, F.P. 346(1985)69
Lötter, A.P., see Du Preez, J.L. 333(1985)249
Lottspeich, F.
Microscale isocratic separation of phenylthiohydantoin amino acid derivatives 326(1985)321
Love, L.J. Cline, see Arunyanart, M. 342(1985)293
Low, G.K.C., Haddad, P.R. and Duffield, A.M.
Proposed model for peak splitting in reversed-phase ion-pair HPLC with computer prediction of eluted peak profiles 336(1985)15
Lowenthal, A., see Marescau, B. 345(1985)215
Lowry, L.K., see Sollenberg, J. 343(1985)175, 343(1985)419
Lu, K., Savaraj, N. and Newman, R.A.
HPLC separation and quantitation of benzisoquinolinedione (Nafidimide) from biological fluids 345(1985)408
Lu, R.C., see Kolbe, H.V.J. 327(1985)1
Lubli, G., see Tagliaro, F. 330(1985)323
Lucek, R. and Dixon, R.
Quantitation of levorphanol in plasma using HPLC with electrochemical detection 341(1985)239
Luger, T.A., see Köck, A. 326(1985)129
Lum, J.T., Vassanji, N.A. and Wells, P.G.
Analysis of the toxicologically relevant metabolites of phenytoin in biological samples by HPLC 338(1985)242
Lundanes, E. and Greibrokk, T.
Group separation of oil residues by supercritical fluid chromatography 349(1985)439
Lundström, K., see Carlsson, A. 350(1985)169
Lutz, M.P., see Abramson, F.P. 339(1985)87
Lutz, T., see Bagchi, S.P. 344(1985)362
Lyon, K.A., see Musson, D.G. 337(1985)363

M

Maas, J.W., see Javors, M.A. 336(1985)259
Maasch, H.J., see Wahl, R. 329(1985)153
McConathy, W.J., Koren, E., Wieland, H., Campos, E.M., Lee, D.M., Kloer, H.U. and Alaupovic, P.
Evaluation of immunoaffinity chromatography for isolating human lipoproteins containing apolipoprotein B 342(1985)47
McCormick, D.B., see Chastain, J.L. 330(1985)153
McCormick, J.P., see Carrel, J.E. 342(1985)411
McCourt, D.W., Leykam, J.F. and Schwartz, B.D.
Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9
McCoy, L., Skee, D. and Edelson, J.
Determination of fezolamine and its desmethyl metabolite in human plasma and urine by HPLC. Intravenous pharmacokinetics in the beagle hound 344(1985)211
McCracken, M.S. and Holt, N.J.
HPLC determination of lipids in vesicles 348(1985)221
McCubbin, W.D., see Cachia, P.J. 343(1985)315

McDermott, B.J., Double, J.A., Bibby, M.C., Wilman, D.E.V., Loadman, P.M. and Turner, R.L.
GC analysis of triethylenethiophosphoramide and triethylenephosphoramide in biological specimens 338(1985)335
McDonald, M.J., see Turci, S.M. 343(1985)168
McDonnell, E., see Gorham, J. 350(1985)245
Macek, J. and Smolková-Keulemansová, E.
Correlation of the chromatographic retention of some phenylacetic and phenylpropionic acid derivatives with molecular structure 333(1985)309
Macek, K., see Schwarz, V. 340(1985)401
McEvoy-Bowe, E.
Cooling plate for cellulose thin-layer electrophoresis and its application to amino acid analysis 347(1985)199
Macewen, G.D., see Salzman, S.K. 343(1985)285
McGee, W.W., see Quinn, M.E. 350(1985)187
McGregor, G.A., see Tucker, F.A. 342(1985)193
McGregor, J.L., Clezardin, P., Manach, M., Gronlund, S. and Dechavanne, M.
Tandem separation of labelled human blood platelet membrane glycoproteins by anion-exchange and gel fast protein LC 326(1985)179
McGregor, W.C., see Roy, S.K. 327(1985)189
McIntyre, I.M., see Novak, E.A. 339(1985)457
McLachlan, D.R.C., see Kruck, T.P.A. 341(1985)123
McLean, A.J., see Harrison, P.M. 339(1985)424, 339(1985)429, 343(1985)349
McNally, M.E. and Rogers, L.B.
Examination of the effect of solvent composition on bonded phase LC packings by ^{13}C Fourier transform NMR spectroscopy 331(1985)23
McQuinn, R.L., see Chang, S.F. 343(1985)119
McVie, J.G., see Van Belle, S.J.P. 337(1985)73
Madden, B.G., see Smith, R.J. 328(1985)11
Madden, J., see Alexander, L.R. 342(1985)1
Maddock, J., see Lewis, P.A. 327(1985)261
Maddocks, J.L., see Dooley, T. 337(1985)321
Maeda, K., Shiraishi, S., Sakamoto, N., Ohki, T., Hosoi, M., Ohta, K. and Yamanaka, N.
Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 345(1985)11
Maeda, K., see Shiraishi, S. 338(1985)51
Mafune, E., see Takasugi, N. 342(1985)349
Magg, H. and Ballschmiter, K.
Retention indices of some ergopeptines in reversed-phase HPLC 331(1985)245
Maghuin-Rogister, G., see Gaspar, P. 328(1985)413
Maglietto, S.M., see Musson, D.G. 338(1985)357
Magnani, M., see Stocchi, V. 349(1985)77
Maguire, J.H. and Wilson, D.C.
Urinary dihydrodiol metabolites of phenytoin: HPLC assay of diastereometic composition 342(1985)323
Maguire, K.P., see Norman, T.R. 340(1985)173
Mahieu, J.-P., Sebille, B., Craescu, C.T., Rhoda, M.-D. and Beuzard, Y.
Determination of the dissociation constant of oligomeric proteins by size-exclusion HPLC: application to human haemoglobin 327(1985)313
Mahler, G.S., see Pirkle, W.H. 348(1985)89
Mahoney, N.E. and Chan, B.G.
HPLC analysis of terpene aldehydes in cotton 329(1985)91
Mailman, R.B., see Kilts, C.D. 342(1985)452
Makdessi, S. al, see Al Makdessi, S. 339(1985)25
Mäkelä, A.-L., see Lempiäinen, M. 341(1985)105
Makin, H.L.J., see Coldwell, R.D. 338(1985)289
Makin, H.L.J., see Street, J.M. 343(1985)259
Makino, K., Moriya, F. and Hatano, H.
Separation of free radicals by HPLC with electron spin resonance detection 332(1985)71
Makino, K., see Ozaki, H. 332(1985)243
Makino, K., see Sasaki, I. 332(1985)237
Makita, A., see Taniguchi, N. 339(1985)394
Makita, M., see Kataoka, H. 339(1985)370
Malcolme-Lawes, D.J., see Blatchford, C. 329(1985)281
Malcolme-Lawes, D.J., see Jones, K. 329(1985)25
Malfait, R., see Gerlo, E. 343(1985)9
Malikin, G., see Karmen, A. 349(1985)267
Malinar, M.J., see Janjić, T.J. 331(1985)273
Maliński, E., see Kuśmierz, J. 331(1985)219
Malison, R.T., see Soparkar, C.N.S. 337(1985)110
Mallinger, A.G., see Edwards, D.J. 344(1985)356
Malspeis, L., Abdel-Hay, M.H. and Wahbi, A.M.
GC determination of pyrazoloimidazole 343(1985)190
Malý, E.
PC of the essential oils occurring in the genus Stachys 333(1985)288
Malý, P., Tichá, M. and Kocourek, J.
Studies on lectins. LVIII. Sugar-binding properties, as determined by affinity electrophoresis, of α-D-galactosidases from *Vicia faba* seeds possessing erythroagglutinating activity 347(1985)343
Mamianetti, A., see Carducci, C.N. 337(1985)91
Manach, M., see McGregor, J.L. 326(1985)179
Manalan, A.S., Newton, D.L. and Klee, C.B.
Purification and peptide mapping of calmodulin and its chemically modified derivatives by reversed-phase HPLC 326(1985)387
Mancini, G., see Gambardella, P. 348(1985)229
Mangia, A., see Gambardella, P. 348(1985)229
Mangia, A., see Verzella, G. 346(1985)417, 349(1985)83
Mannschreck, A., see Rimböck, K.-H. 329(1985)307
Mant, C.T. and Hodges, R.S.
General method for the separation of cyanogen bromide digests of proteins by

HPLC. Rabbit skeletal troponin I 326(1985)349
Mant, C.T. and Hodges, R.S.
Separation of peptides by strong cation-exchange HPLC 327(1985)147
Maoka, T., see Ikuno, Y. 328(1985)387
Mar, T., see Brill, J.H. 349(1985)31, 349(1985)39
Marchelli, R., see Palla, G. 349(1985)9
Marcucci, F., Colombo, L. and Mussini, E.
GC quantitative determination of 1- and 3-methylhistidine in urine and muscles: comparison with glass capillary determination 336(1985)356
Mardarowicz, M., see Suprynowicz, Z. 333(1985)11
Mardens, Y., see Kumps, A. 342(1985)469
Mareš, P. and Hušek, P.
Quantitative capillary GLC of triglycerides on a fused-silica column with a chemically bonded stationary phase 350(1985)87
Marescau, B., De Deyn, P., Wiechert, P., Van Gorp, L. and Lowenthal, A.
Quantitative overestimation of free taurocyamine using short liquid cation-exchange chromatography columns? 345(1985)215
Marfil, F., see Colussi, D. 339(1985)157
Marha, K., see Janda, V. 329(1985)186
María, I. Santa, see Santa María, I. 329(1985)123
Marigo, M., see Tagliaro, F. 330(1985)323
Marino, G., see Cubellis, M.V. 329(1985)406
Marino, G., see Pucci, P. 331(1985)425
Maris, F.A., Geerdink, R.B. and Brinkman, U.A.T.
Selection of mobile phases for reversed-phase LC with on-line electron-capture detection 328(1985)93
Maris, F.A., Van der Vliet, A., Geerdink, R.B. and Brinkman, U.A.T.
Narrow-bore normal phase LC with on-line electron-capture detection 347(1985)75
Marko, V., Štefek, M. and Šoltés, L.
GC determination of pentacaine in rat serum 339(1985)410
Marnot, B., see Masson, P. 328(1985)135
Marquard, F., see Grisham, C.M. 333(1985)301
Martens, J., see Busker, E. 350(1985)179
Martensen, A.N., see Clouse, S.D. 350(1985)255
Martin, L.E., see Colthup, P.V. 345(1985)111
Martin, M., see Guiochon, G. 326(1985)3
Martin, W.J., see Llyod, C.W. 339(1985)139
Martin-Smith, M., see Cholerton, T.J. 333(1985)178
Marzo, A. and Treffner, E.
GLC determination of isosorbide-5-mononitrate derivatized with perfluorinated anhydrides for pharmacokinetic and bioavailability investigations 345(1985)390
Marzo, A., Treffner, E., Quadro, G. and Rapelli, S.
GLC evaluation of bencyclane in biological samples for pharmacokinetic and bioavailability investigations: comparison of two analytical methods 343(1985)77
Masaki, T., see Ganno, S. 332(1985)275
Mashiba, H., see Tatsuhara, T. 339(1985)149
Massart, D.L., see De Smet, M. 345(1985)309
Massart, D.L., see Michotte, Y. 344(1985)367
Massart, D.L., see Musch, G. 348(1985)97
Massé, R., Laliberté, C. and Tremblay, L.
GC-MS of epimeric 19-norandrostan-3-ol-17-ones as the trimethylsilyl ether, methyloxime-trimethylsilyl ether and trimethylsilyl-enol trimethylsilyl ether derivatives 339(1985)11
Masson, P. and Marnot, B.
Affinity electrophoresis in polyacrylamide gel. Influence of the gel concentration on the apparent affinity of cholinesterase for an anionic site ligand (in French) 328(1985)135
Masu, A., see Goto, J. 345(1985)241
Matejka, M., see Carducci, C.N. 337(1985)91
Mathews, S.E., see Bowers, L.D. 333(1985)231
Mathiasson, L., see Skarping, G. 346(1985)191
Mathieu, D., see Lhermitte, M. 342(1985)416
Mathieu-Nolf, M., see Lhermitte, M. 342(1985)416
Mathis, R., Hubert, P. and Dellacherie, E.
Polyethylene oxide: a ligand for mild hydrophobic interaction chromatography? 347(1985)291
Matisová, E., Krupčík, J., Čellár, P. and Kočan, A.
Quantitative analysis of hydrocarbons in gasolines by capillary GLC. II. Isothermal and temperature-programmed analyses 346(1985)177
Matson, R.S., see Goheen, S.C. 326(1985)235
Matsuda, T., see Kato, Y. 332(1985)39
Matsuki, M., see Oyama, H. 338(1985)71
Matsuki, Y., see Tanaka, T. 328(1985)271
Matsumoto, M., see Inouye, K. 327(1985)301
Matsumura, G., see Shimada, E. 328(1985)73
Matsunaga, R., see Yamaguchi, M. 346(1985)227
Matsuno, T., see Ikuno, Y. 328(1985)387
Matsuoka, C., Nohta, H., Kuroda, N. and Ohkura, Y.
Simultaneous determination of cholestanol and cholesterol in human serum by HPLC with fluorescence detection 341(1985)432
Matsushima, Y., Nagata, Y., Niyomura, M., Takakusagi, K. and Takai, N.
Analysis of antipyretics by semimicro LC 332(1985)269
Matsushima, Y., Nagata, Y., Takakusagi, K., Niyomura, M. and Takai, N.
HPLC of antipyretics on chemically modified porous glass 332(1985)265
Matsuzawa, Y., see Hayashi, M. 338(1985)195
Matthaei, D., see Schneider, K. 345(1985)19
Matysik, J., Cięszczyk-Chmiel, A. and Chmiel, J.
Application of interferometry in studies of transport processes near ionites and adsorbents 333(1985)198
Maxwell, J.R., see Gill, J.P. 350(1985)37

Maycock, R. and Utley, D.
Analysis of some trichothecene mycotoxins by LC 347(1985)429
Mayer, B., see Gleispach, H. 344(1985)11
Mayfield, H.T., see Brill, J.H. 349(1985)31, 349(1985)39
Mayol, L., see Cubellis, M.V. 329(1985)406
Maza, M.P., see Vioque, E. 331(1985)187
Mazurkiewicz, B.
Paper electrophoretic characterization of ethylenic compounds as dimethylsulphoniomethylthio derivatives 328(1985)436
Mazza, G.
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179
Mazza, G.
GC and MS studies of the constituents of the rhizome of calamus. II. The volatile constituents of alcoholic extracts 328(1985)195
Mazzucotelli, A., Dadone, A., Frache, R. and Baffi, F.
Determination of trace amounts of lanthanides in rocks and minerals by HPLC 349(1985)137
Meatherall, R.C., Guay, D.R.P. and Chalmers, J.L.
Analysis of meperidine and normeperidine in serum and urine by HPLC 338(1985)141
Mehta, R.G., see Hultin, T.A. 341(1985)187
Mehta, S., see Hayes, A.G. 336(1985)446
Meier, B., see Erdelmeier, C.A.J. 346(1985)456
Meinard, C., Bruneau, P. and Perronnet, J.
High-performance liquid chromatograph coupled with two detectors: a UV spectrometer and a polarimeter. Example in the field of pyrethroids: identification of enantiomers 349(1985)109
Meinard, C., Bruneau, P. and Roche, M.
Coupling a high-performance liquid chromatograph with a liquid scientillation detector: example in the field of pyrethroids 349(1985)105
Meineke, I., see Brendel, E. 339(1985)359
Melander, W.R., see Huang, J.-X. 330(1985)181
Melchor, N., see Chang, C.A. 347(1985)51
Melvin, G., see Beers, M.F. 336(1985)380
Men, L.-C. and Rogers, L.B.
Role of polar groups in fractionating isomeric polystyrene oligomers 347(1985)39
Menéndez, V., see Fernández-Sánchez, E. 333(1985)1
Menichelli, A., see Del Principe, D. 342(1985)285
Mercik, S.H., see Dorrbecker, B.R. 336(1985)293
Merdjan, H., see Brouard, A. 345(1985)453
Merdjan, H., see Paillet, M. 343(1985)455
Merkus, F.W.H.M., see Overdiek, J.W.P.M. 341(1985)279
Merkwitz, H., see Beger, J. 333(1985)319
Merkwitz, H., see Szymanowski, J. 330(1985)61
Mery, D., see Necciari, J. 341(1985)202
Metayer, J.P., see Gaudry, D. 339(1985)404
Metayer, J.P., see Godbillon, J. 338(1985)151
Metz, P.A., Gehas, J., Spriggle, J.E. and Veening, H.
Off-line LC-MS studies of fluorescent β-aminothiol-*o*-phthalaldehyde derivatives 330(1985)307
Meyer, H.D., see Jacob, K. 349(1985)283
Meyer, V.R.
HPLC theory for the practitioner (Review) 334(1985)197
Meyerhoff, G., see Van der Greef, J. 343(1985)397
Miceli, J.N., see Aravind, M.K. 344(1985)428
Michelsen, P., Aronsson, E., Odham, G. and Åkesson, B.
Diastereomeric separations of natural glycero derivatives as their 1-(1-naphthyl)ethyl carbamates by HPLC 350(1985)417
Michelsen, P. and Odham, G.
Highly sensitive diastereoisomeric analysis of secondary alcohols and short-chain 1,2-diacylglycerols by capillary GC-MS 331(1985)295
Michotte, Y. and Massart, D.L.
Capillary GC determination of vincamine in plasma 344(1985)367
Michotte, Y., see Herregodts, P. 345(1985)33
Mico, B.A., see Swagzdis, J.E. 345(1985)203
Micoud, M., see Brouard, A. 345(1985)453
Midha, K.K., see Lim, H.K. 328(1985)378
Mihaly, G.W., Nicholl, D.D., Edwards, G., Ward, S.A., Orme, M.L'E., Warrell, D.A. and Breckenridge, A.M.
HPLC analysis of amodiaquine in human plasma 337(1985)166
Mikalsen, A., see Stray, H. 349(1985)331
Mikasa, H., Sasaki, K., Arata, J., Yamamoto, Y., Ohno, T. and Kodama, H.
Simultaneous measurement of prolidase and prolinase activities in erythrocytes using an isotachophoretic analyser 343(1985)179
Miki, T., see Tanaka, M. 330(1985)253
Milborrow, B.V., see Vaughan, G.T. 336(1985)221
Millán, F., see Vioque, E. 331(1985)187
Miller, A.M., see Chang, S.F. 343(1985)119
Miller, K.J., see Ames, M.M. 341(1985)89
Miller, L.J., see Tietz, P.S. 336(1985)249
Miller, N.T. and Karger, B.L.
High-performance hydrophobic-interaction chromatography on ether-bonded phases. Chromatographic characteristics and gradient optimization 326(1985)45
Miller, R.H., see Echols, R.E. 347(1985)89
Miller, S., see Howes, L.G. 338(1985)401
Miller, T.G.
Characterization of neutralized β-naphthalenesulfonic acid and formaldehyde condensates 347(1985)249
Miller, V., see Kuchař, M. 347(1985)335
Milon, H. and Bur, H.
Direct liquid introduction MS of some

underivatized dipeptides and polypeptides 350(1985)399
Min, B.H. and Garland, W.A.
Quantitation of cibenzoline in human plasma by GC-negative-ion chemical-ionization MS 336(1985)403
Minagawa, K., Kasuya, Y., Baba, S., Knapp, G. and Skelly, J.P.
Determination of dexamethasone in human plasma and urine by electron-impact MS 343(1985)231
Minagawa, K., see Oka, K. 339(1985)253
Minami, Y., see Hayashi, M. 338(1985)195
Minck, R., see Jehl, F. 339(1985)347
Minkler, P.E., Ingalls, S.T., Kormos, L.S., Weir, D.E. and Hoppel, C.L.
Determination of carnitine, butyrobetaine, and betaine as 4'-bromophenacyl ester derivatives by HPLC 336(1985)271
Minty, P.S.B., see Tucker, F.A. 342(1985)193
Misaki, T., see Sagara, K. 328(1985)289
Misir, R., Laarveld, B. and Blair, R.
Evaluation of a rapid method for preparation of fatty acid methyl esters for analysis by GLC 331(1985)141
Mitchell, F., see D'Agostino, G. 338(1985)1
Mitoma, Y., see Inouye, K. 327(1985)301
Mitsui, A., Nohta, H. and Ohkura, Y.
HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61
Miura, K., see Sogami, M. 332(1985)19
Miura, T., see Kai, M. 345(1985)259
Miura, Y., see Hatake, K. 344(1985)339
Miwa, H.
Reversed-phase ion-pair chromatography of some biologically important carboxylic acids as 2-nitrophenylhydrazides 333(1985)215
Miyamoto, G., Sasabe, H., Odomi, M. and Shimizu, T.
HPLC procedure for the determination of a new positive inotropic agent, 3,4-dihydro-6-[4-(3,4-(dimethoxybenzoyl)-1-piperazinyl]-2(1*H*)-quinolinone, in human plasma and urine 338(1985)450
Miyamoto, M., see Hayashi, K. 338(1985)161
Miyata, S., see Sogami, M. 332(1985)19
Miyazaki, K., see Tadano, K. 341(1985)228
Miyazaki, T., see Yoshioka, M. 344(1985)345
Mizobuchi, M., see Kitada, Y. 347(1985)438
Mizoguchi, N., see Ageta, T. 343(1985)186
Mizuno, T., see Kiba, N. 329(1985)147
Mizutani, H., Kabaya, Y. and Wada, E.
HPLC isolation of uric acid from soil for isotopic determination 331(1985)371
Mizutani, H. and Wada, E.
HPLC determination of uric acid in soil 331(1985)359
Mizutani, S., see Taira, H. 339(1985)399
Mizutani, T., Tajima, K., Okino, N. and Yamamoto, K.
Derivatization of butylated hydroxytoluene for GC with electron-capture detection and HPLC with UV detection 333(1985)171
Moerloose, P. de, see De Croo, F. 329(1985)422, 349(1985)301
Moerloose, P. de, see De Schutter, J.A. 350(1985)135
Moertel, D.M., see Ames, M.M. 341(1985)89
Moilanen, O., see Penttilä, I. 338(1985)265
Moing, J.P. le, see Vajta, S. 343(1985)239
Moll, H., see Shukla, A.K. 337(1985)231
Molle, L., see Duez, P. 329(1985)415
Molles, R.J. Jr. and Garceau, Y.
Quantitation of hydralazine hydrochloride in pharmaceutical dosage forms by HPLC 347(1985)414
Molyneux, S.G. and Franklin, M.
Routine determination of unconjugated 3-methoxy-4-hydroxyphenylglycol in plasma using HPLC with electrochemical detection 341(1985)160
Molyneux, S.G., see Davies, C.L. 339(1985)186
Momparler, R.L., see Lin, K.-T. 345(1985)162
Monaghan, J.J., see Catlow, D.A. 328(1985)167
Moncrieff, J.
Assay of nadolol in serum by reversed-phase HPLC with fluorometric detection 342(1985)206
Montagu, M., Levillain, P., Chenieux, J.C. and Rideau, M.
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437
Montay, G. and Tassel, J.P.
Improved HPLC determination of pefloxacin and its metabolite norfloxacin in human plasma and tissue 339(1985)214
Montecucchi, P.C., see Rusconi, L. 346(1985)390, 349(1985)117
Monteil, H., see Jehl, F. 339(1985)347
Monti, J.-P., see Gallice, P. 331(1985)445
Monzani, V., see Castoldi, D. 344(1985)259
Moodie, I.M., Burger, J.A., Hough, B.J., Shephard, G.S. and Labadarios, D.
GLC of amino acids. Variable arginine response 347(1985)179
Moodie, I.M., see Labadarios, D. 339(1985)366
Moody, K.E., see Herring, S.W. 326(1985)217
Moon, R.C., see Hultin, T.A. 341(1985)187
Moore, D.J., see Powis, G. 342(1985)129, 345(1985)402
Moore, T., see Whelpton, R. 341(1985)361
Morales, A., see Owen, J.A. 342(1985)333
More, N.S., see Phillips, T.M. 327(1985)205, 327(1985)213
Morel, D., see Claudy, P. 329(1985)331
Morel, D., see Gonnet, C. 330(1985)227
Morelli, A., see Palmerini, C.A. 339(1985)285
Morgan, E.D., see Scalia, S. 346(1985)301
Morgan, S.L., see Whiton, R.S. 347(1985)109
Mori, M., see Niwa, T. 336(1985)345
Mori, Y., see Baba, S. 341(1985)251
Moriguchi, S. and Takei, S.
Contributions to hydrocarbon retention in

gas-liquid-solid chromatography with polyethylene glycol 6000 as stationary phase 350(1985)15
Moriguchi, S., see Naito, K. 330(1985)193
Morita, K., see Funazo, K. 346(1985)215
Morita, M., see Yasuhara, A. 328(1985)35
Moriya, F., see Makino, K. 332(1985)71
Moriyama, A., see Chung, T.G. 339(1985)67
Morreal, C.E., Sinha, D.K., White, C.J. and Nemoto, D.T.
Assay of prostaglandins in the epithelial cells and fibroblasts of the rat mammary gland 345(1985)380
Morris, G.S. and Simmonds, H.A.
Use of a fundamental elution protocol for the development of reversed-phase HPLC enabling rapid simultaneous determination of purines, pyrimidines and allied compounds commonly found in human biological fluids 344(1985)101
Morris, R.G.
Estimation of plasma hydroxychloroquine by HPLC with UV detection 338(1985)422
Morrison, A., see Zipser, R.D. 339(1985)1
Morrison, L.A. and Driskell, J.A.
Quantities of B_6 vitamers in human milk by HPLC. Influence of maternal vitamin B_6 status 337(1985)249
Mosbach, K., see Glad, M. 347(1985)11
Mosbach, K., see Sellergren, B. 347(1985)1
Moschidis, M.C.
Silicic acid CC of phosphinolipids. I. Phosphinate analogues of lecithin and cephalin, phosphonic acid and the phosphonate analogue of bis(diacylglycero)phosphate: their separation from other phospholipids and related phosphonolipids 329(1985)160
Moseley, M.A., see Pellizzari, E.D. 334(1985)277
Moser, R., see Gleispach, H. 342(1985)245, 344(1985)11
Mostow, N.D., O'Neill, R., Noon, D.L. and Bacon, B.R.
Determination of α-tocopherol in rat liver by HPLC utilizing UV detection 344(1985)137
Mothes, B., see Dedek, W. 331(1985)200
Motoyoshi, K., see Hatake, K. 344(1985)339
Mould, G., see Read, J. 341(1985)437
Mousa, S., Van Loon, G.R., Houdi, A.A. and Crooks, P.A.
HPLC with electrochemical detection for the determination of nicotine and N-methylnicotinium ion 347(1985)405
Mousdale, D.M. and Coggins, J.R.
HPLC of shikimate pathway intermediates 329(1985)268
Muccio, D.D. and DeLucas, L.J.
Isolation of detergent-solubilized monomers of bacteriorhodopsin by size-exclusion HPLC 326(1985)243
Muchtar, A., see Svensson, J.-O. 341(1985)193
Mueller, R., see Francotte, E. 347(1985)25
Mulé, S.J., see Verebey, K. 343(1985)339
Mulholland, M. and Dolphin, R.J.
Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285
Müller, M. and Sorrell, T.C.
Quantitation of sulfidopeptide leukotrienes by reversed-phase HPLC 343(1985)213
Multari, G., see Del Principe, D. 342(1985)285
Munavalli, S., Hsu, F.-L., Hatem, S.F. and Poziomek, E.J.
Separation of organic quaternary salts by reversed-phase TLC 346(1985)463
Murakami, K., see Nakagawa, T. 330(1985)43
Murao, T., see Takata, Y. 332(1985)117
Murata, H., see Nakagawa, T. 330(1985)43
Murayama, H., see Sagara, K. 328(1985)289
Murayama, H., see Yoshida, T. 348(1985)425
Murel, A., Vilde, S., Pank, M., Shevchuk, I. and Kirret, O.
Chromatophoresis: a new approach to the theroy and practice of chromatofocusing. I. General principles 347(1985)325
Murphy, R., Furness, J.B. and Costa, M.
Measurement and chromatographic characterization of vasoactive intestinal peptide from guinea-pig enteric nerves 336(1985)41
Murray, D.K.
Improved reproducibility and quantitative analysis of phospholipids by flame ionization detection 331(1985)303
Murray, S. and Davies, D.S.
Quantitative analysis of (3-methoxy-4-sulphooxyphenyl)ethylene glycol (MHPG sulphate) in human urine 337(1985)441
Musch, E., see Von Sassen, W. 338(1985)113
Musch, G., De Smet, M. and Massart, D.L.
Expert system for pharmaceutical analysis. I. Selection of the detection system in HPLC analysis: UV *versus* amperometric detection 348(1985)97
Musenger, C., see Allinquant, B. 326(1985)281
Muskiet, F.A.J., see Van den Berg, G.A. 339(1985)223
Mussini, E., see Marcucci, F. 336(1985)356
Musson, D.G., Lin, J.H., Lyon, K.A., Tocco, D.J. and Yeh, K.C.
Assay methodology for quantification of the ester and ether glucuronide conjugates of diflunisal in human urine 337(1985)363
Musson, D.G., Maglietto, S.M. and Bayne, W.F.
Determination of the antibiotic fludalanine in plasma and urine by HPLC using a packed-bed, post-column reactor with *o*-phthalaldehyde and 2-mercaptoethanol 338(1985)357
Mustapha, A., see Chowdhury, B.K. 329(1985)178
Musumarra, G., Scarlata, G., Cirma, G., Romano, G., Palazzo, S., Clementi, S. and Giulietti, G.
Qualitative organic analysis. I. Identification of drugs by principal components analysis of standardized TLC data in four eluent systems 350(1985)151

Muto, N. and Tan, L.
Purification of oestrogen synthetase by HPLC. Two membrane-bound enzymes from the human plancenta 326(1985)137
Muto, Y., see Sogami, M. 332(1985)19
Mutschler, E., see Schladitz-Keil, G. 345(1985)99
Mutschler, E., see Spahn, H. 342(1985)458

N

Na, H., see Shah, P. 329(1985)5
Nadiradze, N.A., see Andronikashvili, T.G. 331(1985)402
Nagai, Y., see Harada, T. 337(1985)187
Nagaoka, S., see Sogami, M. 332(1985)19
Nagase, K., see Goto, J. 348(1985)151
Nagashima, H., Tanaka, Y. and Hayashi, R.
Column LC for the simultaneous determination of the enantiomers of loxoprofen sodium and its metabolites in human urine 345(1985)373
Nagata, Y., see Matsushima, Y. 332(1985)265, 332(1985)269
Nagatsu, T., see Kaneda, N. 341(1985)23, 344(1985)93
Nagatsu, T., see Suzuki, H. 341(1985)176
Nagel, G.T., see Dikkeschei, L.D. 345(1985)1
Nagel, G.T., see Van den Berg, G.A. 339(1985)223
Nagle, J., see Llyod, C.W. 339(1985)139
Naidu, R., see Salzman, S.K. 343(1985)285
Naikwadi, K.P., Rokushika, S., Hatano, H. and Ohshima, M.
New liquid crystalline stationary phases for GC of positional and geometrical isomers having similar volatilities 331(1985)69
Naito, C., see Okazaki, M. 336(1985)151
Naito, K., Ohwada, N., Moriguchi, S. and Takei, S.
Adsorption effects in GLC. Solute retention in the hydrocarbon solute-polar stationary phase (polypropylene glycol 2000) system 330(1985)193
Nakagawa, T., Murata, H., Shibukawa, A., Murakami, K. and Tanaka, H.
LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43
Nakagawa, T., see Haginaka, J. 341(1985)115
Nakamoto, M., see Okamoto, T. 342(1985)35
Nakamura, K. and Kato, Y.
Preparative high-performance ion-exchange chromatography 333(1985)29
Nakamura, K., see Inouye, K. 327(1985)301
Nakamura, M., see Hara, S. 344(1985)33
Nakamura, M., see Iwata, T. 344(1985)351
Nakamura, M., see Nishioka, R. 342(1985)89
Nakamura, M., see Yamaguchi, M. 346(1985)227
Nakamura, Y. and Koizumi, Y.
Radioactivity detection system with a CaF_2 (Eu) scintillator for HPLC 333(1985)83
Nakano, K., Shindo, K., Yasaka, T. and Yamamoto, H.
Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127
Nakano, K., Shindo, K., Yasaka, T. and Yamamoto, H.
Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21
Nakatsu, S.L., see Owen, J.A. 342(1985)333
Nakazawa, H., see Kitada, Y. 347(1985)438
Nakazawa, K., see Iwasaki, S. 341(1985)182
Nakazawa, K., see Kobayashi, A. 336(1985)410
Nambara, T., see Goto, J. 345(1985)241, 348(1985)151
Napier, J.R., see Knighton, D.R. 347(1985)237
Narita, O., see Taira, H. 339(1985)399
Narsimha, R., see Husain, S. 348(1985)309
Nash, J.F., see Farid, K.Z. 337(1985)329
Nau, H., see Kuhnz, W. 344(1985)309
Navin, T.R., see Dickinson, C.M. 345(1985)91
Nawrocki, J. and Szczepaniak, W.
Coordination polymers as stationary phases and adsorbents for GC. VIII. An attempt to bind poly[chromium(III) diphenylphosphinate] to silica 330(1985)370
Nayak, V.G., see Sane, R.T. 331(1985)432
Necciari, J., Mery, D., Sales, Y., Berthet, D. and Cautreels, W.
Determination of propisomide, a new antiarrhythmic agent, in biological samples by GC with a thermionic detector 341(1985)202
Necciari, J., see Davi, H. 342(1985)159
Neelakantan, L., Buijten, J. and Ahlborg, U.G.
Analysis of 2-acetylaminofluorene and its metabolites by GC with electron-capture detection 350(1985)307
Neftel, A., see Elbert, W. 328(1985)111
Nelis, H.J.C.F., Lefevere, M.F., Baert, E., D'Hoore, W. and De Leenheer, A.P.
Chromatographic determination of N-acetyl-DL-tryptophan and octanoic acid in human albumin solutions 333(1985)381
Nemoto, D.T., see Morreal, C.E. 345(1985)380
Netting, A.G. and Duffield, A.M.
HPLC separation of fatty acids as pentafluorobenzyl esters 336(1985)115
Neue, U., see Colgan, S.T. 333(1985)349
Neupert, M., see Schneider, K. 345(1985)19
Nevejans, F. and Verzele, M.
Swelling propensity (SP factor) of semi-rigid chromatographic packing materials 350(1985)145
Newhart, A., see Colgan, S.T. 333(1985)349

Newman, R.A., see Lu, K. 345(1985)408
Newton, D.L., see Manalan, A.S. 326(1985)387
Ng, K.T., see Holland, M.L. 345(1985)178
Ng, S., see Ng, T.-L. 329(1985)13
Ng, T.-L. and Ng, S.
Computer simulation of abnormal high-performance liquid chromatograms caused by solvents 329(1985)13
Nicholl, D.D., see Mihaly, G.W. 337(1985)166
Nicholson, M.V., see Duncan, M.W. 336(1985)199
Nieder, M., see Albert, K. 346(1985)17
Nielen, M.W.F., Koordes, R.C.A., Frei, R.W. and Brinkman, U.A.T.
Automated pre-column system for trace enrichment and clean-up of plasma samples in narrow-bore LC 330(1985)113
Nienstedt, W.
Characterization of C_{19} steroids by two-dimensional TLC 329(1985)171
Nierenberg, D.W.
Serum and plasma β-carotene levels measured with an improved method of HPLC 339(1985)273
Nierenberg, D.W. and Lester, D.C.
Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 345(1985)275
Nimura, N., see Yoshida, T. 348(1985)425
Nioka, M., see Kurosawa, N. 343(1985)379
Nishida, S., see Oyama, H. 338(1985)71
Nishidai, T., see Komuro, C. 338(1985)209
Nishimura, M., see Yamamoto, T. 342(1985)261
Nishioka, R., Takeuchi, M., Kawai, S., Nakamura, M. and Kondo, K.
Improved direct injection method and extractive methylation method for determination of valproic acid in serum by GC 342(1985)89
Nisi, C., see Farina, P. 345(1985)323
Nissinen, E.
Assay of aromatic L-amino acid decarboxylase by HPLC with radiochemical detection 342(1985)175
Niwa, T., Yamada, K., Ohki, T. and Furukawa, H.
3-Hydroxyhexanoic acid: an abnormal metabolite in urine and serum of diabetic ketoacidotic patients 337(1985)1
Niwa, T., Yamada, K., Ohki, T., Saito, A. and Mori, M.
Identification of 6-deoxyallitol and 6-deoxygulitol in human urine. Electron-impact mass spectra of eight isomers of 6-deoxyhexitol 336(1985)345
Niwaguchi, T., see Inoue, T. 339(1985)163
Niyomura, M., see Matsushima, Y. 332(1985)265, 332(1985)269
Noguchi, H., see Tokuma, Y. 345(1985)51
Noguchi, K., see Hanai, T. 349(1985)323
Noguchi, K., see Ito, N. 333(1985)107, 348(1985)199
Noguchi, K., see Seki, T. 332(1985)9, 332(1985)283
Noguchi, K., see Sogami, M. 332(1985)19
Noguchi, K., see Yasukawa, K. 332(1985)287
Noguchi, M., see Oka, K. 339(1985)253, 345(1985)419
Nohta, H., see Lee, M. 348(1985)407
Nohta, H., see Matsuoka, C. 341(1985)432
Nohta, H., see Mitsui, A. 344(1985)61
Nomura, Y., see Harada, T. 337(1985)187
Noon, D.L., see Mostow, N.D. 344(1985)137
Nordlander, I., see Österdahl, B.-G. 337(1985)151
Norman, E.J., Wee, E.L., Berry, H.K. and Zimmerman, E.F.
Rapid GC-MS quantitation of γ-aminobutyric acid in biological specimens 337(1985)21
Norman, H.A., see Smith, L.A. 346(1985)291
Norman, T.R. and Maguire, K.P.
Analysis of tricyclic antidepressant drugs in plasma and serum by chromatographic techniques (Review) 340(1985)173
Normandin, G., see Swobodnik, W. 339(1985)263
Noro, Y., see Kaneda, N. 344(1985)93
Norris, K.J., see Thompson, J.A. 341(1985)349
Norrlöw, O., see Glad, M. 347(1985)11
Norsten, C., see Cronholm, T. 344(1985)1
Nota, G., Improta, C. and Cannata, A.
Liquefied gases as eluents for TLC 347(1985)362
Novak, E.A., Stanley. M., McIntyre, I.M. and Hryhorczuk, L.M.
HPLC method for quantification of cyproheptadine in serum or plasma 339(1985)457
Novák, J., see Giliáš, J. 346(1985)43
Novotny, M., see Gluckman, J.C. 333(1985)291
Novotný, M., see Smith, S.L. 336(1985)351
Nyarady, S.A., see Sievers, R.E. 349(1985)395
Nye, E.R., see Poll, D. 343(1985)149
Nyquist, O., see Hermansson, J. 336(1985)321
Nzekwe, E.U., see Kabra, P.M. 341(1985)383

O

Oates, J.A., see Roberts, L.J. II 338(1985)41
Obbens, H. and Lendero, L.
Novel approaches to the determination of polychlorinated biphenyls 349(1985)425
Obbens, H., see Huber, L. 349(1985)465
Ochs, H.R., see Locniskar, A. 337(1985)131
O'Connell, K., see Sentissi, A. 331(1985)396
Odham, G., see Michelsen, P. 331(1985)295, 350(1985)417
Odomi, M., see Akiyama, H. 338(1985)456
Odomi, M., see Miyamoto, G. 338(1985)450
Oehme, M., see Stray, H. 349(1985)331
Ogawa, H., see Chung, T.G. 339(1985)67

Ogawa, I. and Fritz, J.S.
Determination of low concentrations of low-molecularweight aldehydes and ketones in aqueous samples 329(1985)81
O'Gorman Hughes, D.W., see So, N. 337(1985)81
O'Hare, M.J., see D'Agostino, G. 338(1985)1
O'Hare, M.J., see Smith, J.A. 345(1985)168
Ohashi, S., see Baba, Y. 348(1985)27, 350(1985)119, 350(1985)461
Ohki, T., see Maeda, K. 345(1985)11
Ohki, T., see Niwa, T. 336(1985)345, 337(1985)1
Ohki, T., see Shiraishi, S. 338(1985)51
Ohkubo, I., see Chung, T.G. 339(1985)67
Ohkura, Y., see Hamada, C. 341(1985)426
Ohkura, Y., see Hara, S. 344(1985)33
Ohkura, Y., see Ishida, J. 344(1985)267
Ohkura, Y., see Iwata, T. 344(1985)351
Ohkura, Y., see Kai, M. 345(1985)259
Ohkura, Y., see Lee, M. 348(1985)407
Ohkura, Y., see Matsuoka, C. 341(1985)432
Ohkura, Y., see Mitsui, A. 344(1985)61
Ohkura, Y., see Yamaguchi, M. 346(1985)227
Ohnishi, N., see Kataoka, H. 339(1985)370
Ohno, T., see Mikasa, H. 343(1985)179
Ohshima, M., see Naikwadi, K.P. 331(1985)69
Ohta, K., Yamaguchi, K., Jikumaru, N. and Kawahara, K.
Heterogeneities of blue dextran and differences among samples 350(1985)292
Ohta, K., see Maeda, K. 345(1985)11
Ohta, K., see Shiraishi, S. 338(1985)51
Ohwada, N., see Naito, K. 330(1985)193
Oikawa, K., see Saitoh, H. 329(1985)247
Oka, K., Aoshima, S. and Noguchi, M.
Highly sensitive determination of ketoprofen in human serum and urine and its application to pharmacokinetic study 345(1985)419
Oka, K., Ijitsu, T., Minagawa, K., Hara, S. and Noguchi, M.
Efficient chromatographic fractionation of steroids in human serum through regulation of liquid-liquid distribution ratios 339(1985)253
Okada, T. and Kuwamoto, T.
Trace analysis of anions by use of a back-flush method and large injection volumes in ion chromatography 350(1985)317
Okada, Y., see Koizumi, K. 341(1985)31
Okamoto, T., Fukui, K., Nakamoto, M., Kishi, T., Okishio, T., Yamagami, T., Kanamori, N., Kishi, H. and Hiraoka, E.
HPLC of coenzyme Q-related compounds and its application to biological materials 342(1985)35
Okamoto, Y., Honda, S., Hatada, K. and Yuki, H.
Chromatographic resolution. IX. HPLC resolution of enantiomers on optically active poly(triphenylmethyl methacrylate) 350(1985)127
Okamura, Y., see Tatsuhara, T. 339(1985)149
Okazaki, M., Kinoshita, M., Naito, C. and Hara, I.
HPLC of apolipoproteins in serum high-density lipoproteins 336(1985)151
O'Keefe, J.H., Sharry, L.F. and Jones, A.J.
Macroscale HPLC separation and instrumental identification of components of diethylaminoethyl murine epidermal growth factor 336(1985)73
Okerholm, R.A., see Smithers, J.A. 341(1985)232
Okino, N., see Mizutani, T. 333(1985)171
Okishio, T., see Okamoto, T. 342(1985)35
Olde, B., see Johansson, G. 331(1985)11
Olichon, D., see Picard, M. 341(1985)445
Oliphant, C., see Tsai, M.Y. 341(1985)1
Oliw, E.H.
Analysis of epoxyeicosatrienoic acids by GC-MS using chlorohydrin adducts 339(1985)175
Olschwang, D., see Bassene, E. 346(1985)428
Olsen, I., see Brondz, I. 342(1985)13, 345(1985)119
Olson, E.S. and Diehl, J.W.
Size-exclusion chromatography-low-angle laser light-scattering photometry of lignite macromolecules 349(1985)337
Olson, K.L. and Swarin, S.J.
Determination of aldehydes and ketones by derivatization and LC-MS 333(1985)337
Olsson, O., see Gallo, P. 327(1985)293
Omichi, K. and Ikenaka, T.
Ion-exchange chromatography of fluorogenic derivatives of maltooligosaccharides for preparation of α-amylase substrates 336(1985)368
Omile, C.I., see Tebbett, I.R. 329(1985)196
O'Neill, R., see Mostow, N.D. 344(1985)137
Ono, K., see Komuro, C. 338(1985)209
Ono, M., see Taniguchi, N. 339(1985)394
Opheim, K., see Weber, A.F. 337(1985)434
Ordway, F.S., see Schilsky, R.L. 337(1985)63
Orichowskyj, S.T., see Roy, S.K. 327(1985)189
Orme, M. L'E., see Edwards, G. 343(1985)224
Orme, M.L'E., see Mihaly, G.W. 337(1985)166
Orsini, F. and Verotta, L.
Separation of natural polar substances by reversed-phase HPLC, centrifugal TLC and droplet counter-current chromatography 349(1985)69
Osek, J., see Oszczapowicz, J. 330(1985)79
Oshima, T., see Sagara, K. 328(1985)289
Osman, M.A., Pinchbeck, F.M., Cheng, L.K. and Wright, G.J.
Simultaneous determination of fluzinamide and three of its active metabolites in plasma by HPLC 336(1985)329
Osman, M.A., Pinchbeck, F.M., Cheng, L.K. and Wright, G.J.
Determination of lenperone in dog plasma by HPLC 345(1985)430
Ośmiałowski, K., Halkiewicz, J., Radecki, A. and Kaliszan, R.
Quantum chemical parameters in correlation analysis of GLC retention indices of amines 346(1985)53

Osswald, W.F.
HPLC of isoflavonoid phytoalexins from soybean (*Glycine max*, L. Harosoy 63) 333(1985)225
Österdahl, B.-G., Johnsson, H. and Nordlander, I.
Rapid Extrelut column method for determination of levamisole in milk using HPLC 337(1985)151
Ostrowski, M., see Oszczapowicz, J. 330(1985)79
Oszczapowicz, J., Osek, J., Ciszkowski, K., Krawczyk, W. and Ostrowski, M.
Retention indices of dimethylbenzamidines and benzylideneamines on a non-polar column 330(1985)79
Otsuka, K., Terabe, S. and Ando, T.
Electrokinetic chromatography with micellar solutions. Separation of phenylthiohydantoin-amino acids 332(1985)219
Otsuka, K., Terabe, S. and Ando, T.
Electrokinetic chromatography with micellar solutions. Retention behaviour and separation of chlorinated phenols 348(1985)39
Otsuka, K., see Terabe, S. 332(1985)211
Overdiek, J.W.P.M., Hermens, W.A.J.J. and Merkus, F.W.H.M.
Determination of the serum concentration of spironolactone and its metabolites by HPLC 341(1985)279
Owada, E., see Kurosawa, N. 343(1985)379
Owen, A.J., see Andrews, A.T. 348(1985)177
Owen, J.A., Nakatsu, S.L., Condra, M., Surridge, D.H., Fenemore, J. and Morales, A.
Sub-nanogram analysis of yohimbine and related compounds by HPLC 342(1985)333
Owens, J.A. and Robinson, J.S.
Isolation and quantitation of carbohydrates in sheep plasma by HPLC 338(1985)303
Oyama, H., Endoh, M., Yoneda, M., Matsuki, M., Satoh, A., Nishida, S. and Horino, M.
New application of HPLC for analysis of urinary C-peptide 338(1985)71
Ozaki, H., Wada, H., Takeuchi, T., Makino, K., Fukui, T. and Kato, Y.
Behaviour of single-stranded oligodeoxyribonucleotides on a DEAE-5PW anion-exchange column 332(1985)243
Ozaki, H., see Terabe, S. 332(1985)211

P

Paalman, A.C.A., see Van Belle, S.J.P. 337(1985)73
Paar, W.D., see Von Unruh, G.E. 345(1985)67
Pabst, R. and Dose, K.
Rapid delipidation of the membrane protein bacteriorhodopsin by HPLC 350(1985)427
Pacáková, V., see Tomková, H. 329(1985)113
Padilla, P., see De Guevara, O.L. 329(1985)428
Paillet, M., Merdjan, H., Brouard, A., Doucet, D., Barreteau, H. and Fredj, G.
Rapid determination of alminoprofen in plasma by HPLC 343(1985)455
Paillet, M., see Brouard, A. 345(1985)453
Palazzo, S., see Musumarra, G. 350(1985)151
Palla, G., Dossena, A. and Marchelli, R.
Oligo-oxaalkanoyl tetraamides derived from L-phenylalanine as stationary phases in capillary GC resolution of D,L-amino acids 349(1985)9
Palmborg, J., see Borén, H. 348(1985)67
Palmerini, C.A., Fini, C., Floridi, A., Morelli, A. and Vedovelli, A.
HPLC analysis of free hydroxyproline and proline in blood plasma and of free and peptide-bound hydroxyproline in urine 339(1985)285
Palmisano, F., Cataldi, T.R.I. and Zambonin, P.G.
Determination of the antineoplastic agent methotrexate in body fluids by HPLC with electrochemical detection 344(1985)249
Palvadre, R., see Haldna, Ü. 350(1985)296
Pank, M., see Murel, A. 347(1985)325
Pannell, K.H., see Chang, C.A. 347(1985)51
Panteix, G., see Leclercq, M. 337(1985)423
Pao, J., see Puglisi, C.V. 344(1985)199
Pap-Sziklay, Z., see Gazdag, M. 328(1985)279
Papa, P., see Groppi, A. 337(1985)142
Parcher, J.F., see Hyver, K.J. 328(1985)63
Parekh, N.J. and Rogers, L.B.
Effects of various strengths of bases on the electrophoretic behavior of weak acids in a non-aqueous solvent 330(1985)19
Park, C.M., Reid, P.E. and Applegarth, D.A.
Evidence for the degradation of sugars during the resin hydrolysis of glycoproteins 339(1985)182
Park, Y.H., Goshorn, C. and Hinsvark, O.
Two HPLC methods for quantitation of theophylline in plasma 343(1985)359
Parker, C.E., Bursey, M.M., Smith, R.W. and Gaskell, S.J.
Gaseous acidity and basicity scales as guides to chemical ionization in reversed-phase LC-MS with direct liquid introduction 347(1985)61
Parker, C.E., Levy, L.A., Smith, R.W., Yamaguchi, K., Gaskell, S.J. and Korach, K.S.
Separation and detection of enantiomers of stilbestrol analogues by combined HPLC-thermospray MS 344(1985)378
Parkin, S., see Lennard, M.S. 338(1985)249
Parkinson, D.
Sensitive analysis of butyrophenone neuroleptics by HPLC with UV detection at 254 nm 341(1985)465

Pasutto, F.M., see Torok-Both, G.A. 344(1985)372
Paterson, M.D., see Wheals, B.B. 350(1985)205
Patrick, K.S., Ellington, K.R., Breese, G.R. and Kilts, C.D.
GC-MS analysis of methylphenidate and *p*-hydroxymethylphenidate using deuterated internal standards 343(1985)329
Patrick, P.H. and Elliott, W.H.
Bile acids. LXXVI. Analyses of bile acids and sterols by HPLC with a microbore column 347(1985)155
Patrykiejew, A., see Suprynowicz, Z. 333(1985)11
Patsalides, E. and Robards, K.
Rapid chromatographic determination of benzotiazoles in automotive cooling waters and cooling water formulations 331(1985)149
Patsalides, E. and Robards, K.
Applications of fused-silica wall-coated open-tubular columns. I. Schiff base chelates 350(1985)353
Paulsen, B.S., see Barsett, H. 329(1985)315
Paxton, J.W., see Jurlina, J.L. 342(1985)431
Pearson, C.D. and Gharfeh, S.G.
Determination of trimer in *Z*-11-hexadecenal pheromone by HPLC with a rotating disc flame ionization detector 329(1985)142
Peč, P., see Stránský, Z. 342(1985)167
Pechová, H., see Horna, A. 330(1985)439
Peel, H.W., see Perrigo, B.J. 341(1985)81
Pehle, W., see Döring, C.E. 348(1985)430
Pellinen, J. and Salkinoja-Salonen, M.
High-performance size-exclusion chromatography of lignin and its derivatives 328(1985)299
Pellizzari, E.D., Moseley, M.A. and Cooper, S.D.
Recent advances in the analysis of polychlorinated biphenyls in environmental and biological media (Review) 334(1985)277
Pennings, E.J.M., Verhagen, J.C.M. and Van Kempen, G.M.J.
Assay of urinary phenylacetic acid by HPLC 341(1985)172
Pentshuk, J., see Haldna, Ü. 350(1985)296
Penttilä, I., Huhtikangas, A., Herranen, J. and Moilanen, O.
GC determination of serum branched-chain α-keto acids derivatized by extractive alkylation 338(1985)265
Percheron, F., see Succari, M. 341(1985)457
Pernemalm, P.-Å., see Andersson, T. 326(1985)33
Perrier, D.G., see Schaaf, L.J. 342(1985)303
Perrigo, B.J., Peel, H.W. and Ballantyne, D.J.
Use of dual-column fused-silica capillary GC in combination with detector response factors for analytical toxicology 341(1985)81
Perronnet, J., see Meinard, C. 349(1985)109
Perseo, G., see Rusconi, L. 349(1985)117
Persson, B.-A., see Ahnoff, M. 340(1985)73
Pertierra-Rimada, E., see Fernández-Sánchez, E. 333(1985)1
Peters, G.J., see Kok, R.M. 343(1985)59
Petersen, D.R., see Thompson, J.A. 341(1985)349
Petit, J., see Bougerolle, A.M. 345(1985)59
Petrović, S.M., Lomić, S. and Šefer, I.
Utilization of the functional group contribution concept in LC on chemically bonded reversed phases 348(1985)49
Pfäffli, P., Härkönen, H. and Savolainen, H.
Urinary dimethylbenzoic acid excretion as an indicator of occupational exposure to white spirit 337(1985)146
Pfäffli, P. and Svartling, N.
Assay of methyl methacrylate in blood samples by headspace capillary GC 345(1985)386
Pfeiffer, C.D., see Cortes, H.J. 349(1985)55
Phillipou, G., see Frith, R.G. 338(1985)179
Phillips, L.S., see Goldstein, S. 339(1985)388
Phillips, T.M., More, N.S., Queen, W.D. and Thompson, A.M.
Isolation and quantitation of serum IgE levels by high-performance immunoaffinity chromatography 327(1985)205
Phillips, T.M., Queen, W.D., More, N.S. and Thompson, A.M.
Protein A-coated glass beads. Universal support medium for high-performance immunoaffinity chromatography 327(1985)213
Phipps, F.C., see Sollenberg, J. 343(1985)419
Picard, M., Olichon, D. and Gombert, J.
Determination of serotonin in plasma by LC with electrochemical detection 341(1985)445
Piccialli, G., see Cubellis, M.V. 329(1985)406
Piccoli, G., see Stocchi, V. 349(1985)77
Pickert, A., see Liebich, H.M. 338(1985)25
Pietrogrande, M.C., Blo, G. and Bighi, C.
HPLC determination of naphthols as 4-aminoantipyrine derivatives. Application to carbaryl 349(1985)63
Pietrogrande, M.C., see Barbaro, A.M. 347(1985)209
Pinatel, H., see Sabot, J.F. 339(1985)233
Pinchbeck, F.M., see Osman, M.A. 336(1985)329, 345(1985)430
Pinelli, A., see Zecca, L. 339(1985)203
Pingoud, V.
Homogenous [mono-^{125}I-Tyr^{1}]- and [mono-^{125}I-Tyr^{13}] glucagon 331(1985)125
Pingoud, V., see Sonnenberg, J. 331(1985)133
Pini, D., see Rosini, C. 348(1985)79
Pirkle, W.H. and Hyun, M.H.
Effect of interstrand distance upon chiral recognition by a chiral stationary phase 328(1985)1
Pirkle, W.H., Pochapsky, T.C., Mahler, G.S. and Field, R.E.
Chromatographic separation of the enantiomers of 2-carboalkoxyindolines and N-aryl-α-amino esters on chiral stationary phases derived from N-(3,5-dinitrobenzoyl)-α-amino acids 348(1985)89
Platt, D., see Brosche, T. 345(1985)219

Pizák, Z., see Weidenhoffer, Z. 350(1985)324
Pochapsky, T.C., see Pirkle, W.H. 348(1985)89
Podhradský, D., Toth, G., Suchár, G., Kristian, P. and Antalík, M.
Adsorption of SH-containing amino acids, peptides and proteins on the 2-nitrophenylhydrazonopropanedinitrile derivative of Spheron Ara 328(1985)402
Podojil, M., see Řezanka, T. 346(1985)453
Pok, P.R., Viala, A. and Durand, A.
Determination of citalopram, amitriptyline and clomipramine in plasma by reversed-phase HPLC 338(1985)171
Pokhilo, N.D., see Pokushalova, T.V. 329(1985)189
Poklis, A., see Hale, P.W. Jr. 336(1985)452
Pokushalova, T.V., Glebko, L.I., Pokhilo, N.D. and Uvarova, N.I.
Preparation and GLC of trimethylsilyl derivatives of dammarane-type triterpene triols and tetrols 329(1985)189
Politi, L., see Laganà, A. 345(1985)251
Poll, D., Harding, D.R.K., Hancock, W.S., Nye, E.R., Janus, E.D., Hannan, S.F. and Scott, P.J.
Use of reversed-phase HPLC to analyse the cholesterol ester profiles present in lipoproteins of patients exhibiting hyperalphalipoproteinaemia or hypoalphalipoproteinaemia (Tangier disease) 343(1985)149
Pommier, F., see Sioufi, A. 339(1985)117
Pons, J.-L., Rimbault, A., Darbord, J.C. and Leluan, G.
GC-MS analysis of volatile amines produced by several strains of *Clostridium* 337(1985)213
Poole, C.F., see Dhanesar, S.C. 349(1985)249
Poole, C.F., see Furton, K.G. 349(1985)235
Poon, B.T., see Brown, N.D. 345(1985)299
Poppe, H., see Van Vliet, H.P.M. 346(1985)149
Porsch, B., see Kotal, P. 333(1985)141
Porter, C., see Catlow, D.A. 328(1985)167
Portoukalian, J., see Bouchon, B. 342(1985)385
Postaire, E., see Le Hoang, M.D. 346(1985)382
Potaman, V.N., Ulmasov, T.N. and Demidov, V.V.
HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33
Pousset, J.L., see Bassene, E. 346(1985)428
Powis, G., Kroschel, D.M. and Moore, D.J.
Sensitive reversed-phase HPLC assay for the antitumor agent benzisoquinolinedione (Nafidimide) employing direct on-column injection of plasma and urine, and its use for pharmacokinetic studies in a dog 345(1985)402
Powis, G. and Moore, D.J.
HPLC assay for the antitumor glycoside [illegible] and its stability in plasma of several species 342(1985)129
Poy, F. and Cobelli, L.
Determination of permanent gases and light hydrocarbons by simultaneous operation on packed and capillary columns with thermal conductivity detection and flame-ionization detection on a single computer integrator 349(1985)17
Poziomek, E.J., see Munavalli, S. 346(1985)463
Pradeau, D., see Le Hoang, M.D. 346(1985)382
Praznik, W. and Beck, R.H.F.
Application of GPC systems to the determination of the molecular weight of inulin 348(1985)187
Preez, J.L. du, see Du Preez, J.L. 333(1985)249
Preston, M.R., see Varney, M.S. 348(1985)265
Prey, T., see Boos, R. 328(1985)233
Přikrylová, V., see Jizba, J.V. 329(1985)193
Principe, D. del, see Del Principe, D. 342(1985)285
Prinoth, M., see Spahn, H. 342(1985)458
Přistoupil, T.I., Kramlová, M., Fořtová, H. and Fričová, V.
Analytical isoelectric focusing of native and modified haemoglobins after treatment with 4-hydroxymercuribenzoate 346(1985)401
Prognon, P., see Le Hoang, M.D. 346(1985)382
Prókai, L.
Separation and identification of phenol-formaldehyde condensates by GC-MS. I. Acid-catalysed condensation products 329(1985)290
Prókai, L.
Separation and identification of phenol-formaldehyde condensates by GC-MS. II. Base-catalysed condensation products 331(1985)91
Prókai, L.
Separation and identification of phenol-formaldehyde condensates by GC-MS. III. Hemiformal isomers in resol 333(1985)161
Proksch, B., see Ehninger, G. 342(1985)119
Proust, S., see Cuisinaud, G. 341(1985)97
Provost, L.R., see D'Agostino, P.A. 331(1985)47, 347(1985)257
Przyjazny, A.
Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327
Przyjazny, A.
Evaluation of the suitability of selected porous polymers for preconcentration of organosulphur compounds from water 346(1985)61
Pucci, P., Borgia, R. and Marino, G.
Sequence determination of cyanogen bromide-peptides by combined use of HPLC and fast atom bombardment MS 331(1985)425
Puglisi, C.V., Pao, J., Ferrara, F.J. and De Silva, J.A.F.
Determination of midazolam (Versed) and its metabolites in plasma by HPLC 344(1985)199
Puglisi, C.V., see Strojny, N. 336(1985)301
Puigmacia, M., see Adzet, T. 348(1985)447
Puigserver, A., see Gaertner, H. 350(1985)279

Pullen, R.H. and Cox, J.W.
Determination of (15*R*)- and (15*S*)-15-methylprostaglandin E_2 in human plasma with picogram per milliliter sensitivity by column-switching HPLC 343(1985)271
Punziano, R., see Gambardella, P. 348(1985)229
Putnam, F.W., see Takahashi, N. 326(1985)373, 326(1985)407

Q

Quadro, G., see Marzo, A. 343(1985)77
Quarry, M.A., see Stadalius, M.A. 327(1985)93
Queen, W.D., see Phillips, T.M. 327(1985)205, 327(1985)213
Qui, Z.Y., see Rokushika, S. 332(1985)15
Quinn, M.E. and McGee, W.W.
Analysis technique for determining the levels of organic additives in an ethylene-propylene diene monomer rubber for use in underwater acoustic applications 350(1985)187
Quinn, M.J., see Wettenhall, R.E.H. 336(1985)51
Quintero, R., see De Guevara, O.L. 329(1985)428

R

Radecki, A., see Ośmiałowski, K. 346(1985)53
Räder, K., Wildfeuer, A., Schwedass, A. and Laufen, H.
Determination of josamycin in plasma, erythrocytes and leucocytes via HPLC by direct injection of plasma or homogenized blood cells, using alternating sample clean-up columns 344(1985)416
Radford, T., see Chamblee, T.S. 330(1985)141
Radivojša, P.N., see Janjić, T.J. 331(1985)273
Rae, I.D., see Browne, C.A. 336(1985)211
Rakhit, A., see Tipnis, V. 345(1985)396
Ramamonjinirina, E., see Gonnet, C. 330(1985)227
Ramus, T.L. and Thomas, L.C.
Temperature selection for chlorinated hydrocarbon reduction for the Hall electrolytic conductivity detector 328(1985)342
Ramus, T.L., see Dodo, G.H. 330(1985)172
Rantwijk, F. van, see Tjioe, T.T. 330(1985)412
Rapelli, S., see Marzo, A. 343(1985)77
Raponi, G., see Andreolini, F. 337(1985)379
Rapoport, S.I., see Sweeney, D.J. 339(1985)434
Rapp, U., see Van der Greef, J. 343(1985)397
Rapsomatiotis, A., see Voss, R.H. 346(1985)205
Rasmussen, K.E., see Lambrechts, M. 331(1985)339
Rassi, Z. el, see El Rassi, Z. 326(1985)79
Rathouská, Z., see Hajšlová, J. 348(1985)437
Raulin, F., see Bossard, A. 330(1985)400
Ravichandran, K. and Baldwin, R.P.
Determination of hydralazine and metabolites in urine by LC with electrochemical detection 343(1985)99
Ravinale, E. and Rosso, A.
Reversed-phase HPLC with isocratic elution of 6α-methyl-17α-acetoxyprogesterone (MAP) and its impurities 330(1985)415
Ravindranath, Y., see Stout, M. 342(1985)424
Ray, A., see Barnes, W.N. 347(1985)173
Ray, J.E. and Day, R.O.
HPLC determination of a new anti-inflammatory agent, nabumetone, and its major metabolite in plasma using fluorimetric detection 336(1985)234
Read, J., Mould, G. and Stevenson, D.
Simple HPLC method for the determination of medroxyprogesterone acetate in human plasma 341(1985)437
Reglero, G., Herraiz, M., Cabezudo, M.D., Fernández-Sánchez, E. and García-Domínguez, J.A.
Contribution to the study of micropacked columns in GC 348(1985)327
Regnier, F.E., see Geng, X. 332(1985)147
Reichert, A., see Toenges, V. 330(1985)429
Reid, J.L., see Howes, L.G. 338(1985)401
Reid, P.E., see Park, C.M. 339(1985)182
Reijenga, J.C., Verheggen, T.P.E.M. and Everaerts, F.M.
Theoretical and practical limitations of capillary isotachophoresis in trace analysis 328(1985)353
Reilly, P.J., see Selosse, E.J.-M. 328(1985)253
Reischer, R.J., see Anderson, D.J. 350(1985)313
Rejholec, V., see Kuchař, M. 333(1985)21, 347(1985)335
Rejholec, V., see Tomková, H. 329(1985)113
Rendi, M.A., Ellis, P.P. and Gal, J.
HPLC assay for pilocarpine in aqueous humor 336(1985)337
Renman, L., see Skarping, G. 346(1985)191
Renner, E., see Wahlländer, A. 338(1985)369
Retunsky, V.N., see Berezkin, V.G 330(1985)71
Reu, L. de, see De Schryver, E.P. 338(1985)389
Reusch, J., see Rühl, C. 328(1985)362
Reutter, D.J., see Bossle, P.C. 330(1985)388
Reynolds, E.S., see Bandi, Z.L. 329(1985)57
Řezanka, T. and Podojil, M.
Gradient separation of fatty acids (C_{14}-C_{30}) by reversed-phase HPLC 346(1985)453
Rhoda, M.-D., see Mahieu, J.-P. 327(1985)313
Rhys Williams, A.T.
Simultaneous determination of serum vitamin A and E by LC with fluorescence detection 341(1985)198

Ribick, M., see Jemal, M. 345(1985)299
Ricard, C.S. and Sturman, L.S.
Isolation of the subunits of the coronavirus envelope glycoprotein E2 by hydroxyapatite HPLC 326(1985)191
Richard, J., see Gaudry, D. 339(1985)404
Richfield-Fratz, N., Bailey, J.E. Jr. and Bailey, C.J.
Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109
Richter, B.E., see Cortes, H.J. 349(1985)55
Richter, W.O., see Schwandt, P. 345(1985)145
Rideau, M., see Montagu, M. 331(1985)437
Riekkola, M.-L., see Grob, K. Jr. 334(1985)129
Riemer, R.K. and Kuhn, R.W.
Rapid and simple purification of the ^{125}I-labeled α-1 adrenergic radioligand 2-[β-(4-hydroxylphenylethyl)aminomethyl] tetralone (BE 2254) using reversed-phase HPLC 338(1985)236
Riesner, D., see Hecker, R. 326(1985)251
Rietschel, E.T., see Shukla, A.K. 337(1985)231
Riffe, A., Delpech, M., Levy-Favatier, F., Boissel, J.-P. and Kruh, J.
Analysis and preparation of chromosomal high-mobility group proteins by ion-exchange HPLC 344(1985)332
Righetti, P.G. and Gianazza, E.
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71
Righetti, P.G., see Bosisio, A.B. 330(1985)299
Rijk, M.A.H., see Van der Greef, J. 343(1985)397
Riley, C.M., see Karnes, H.T. 338(1985)377
Rimbault, A., see Pons, J.-L. 337(1985)213
Rimböck, K.-H., Kastner, F. and Mannschreck, A.
LC on triacetylcellulose. Preparative separation of enantiomers on an axially compressed column 329(1985)307
Rinkus, S.J. and Legator, M.S.
Use and regeneration of Amicon ultrafiltration cones for deproteination of microsomal solutions before chromatographic analysis 342(1985)465
Ritchie, A.S.
Effect of copper(II) sulphate impregnation of Chromarods on the sensitivity of the Iatroscan detection system to lipids 346(1985)468
Ritchie, A.S. and Jee, M.H.
HPLC technique for the separation of lipid classes 329(1985)273
Rittenhouse, C.T., see Sisco, W.R. 348(1985)253
Rivard, G.E., see Lin, K.-T. 345(1985)162
Rivett, D.E., see Logan, R.I. 348(1985)421
Rizzi, A.
Adsorption from multi-component liquid mixtures on to heterogeneous solid surfaces: the role of adsorption of water on silica in liquid-solid adsorption chromatography 348(1985)1
Rizzi, A. and Lang, H.R.M.
Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33
Robards, K., see Patsalides, E. 331(1985)149, 350(1985)353
Robert, J.C., Soumarmon, A. and Lewin, M.J.M.
Determination of O-phosphothreonine, O-phosphoserine, O-phosphotyrosine and phosphate by HPLC 338(1985)315
Robert, T.A. and Hagardorn, A.N.
Plasma levels and kinetic disposition of 2,4-dinitrophenol and its metabolites 2-amino-4-nitrophenol and 4-amino-2-nitrophenol in the mouse 344(1985)177
Roberts, L.J. II, Aulsebrook, K.A. and Oates, J.A.
Comparative evaluation of the radioenzymatic method for the determination of urinary histamine with a MS assay 338(1985)41
Roberts, M.S., see Rumble, R.H. 342(1985)436
Roberts, R.F. and Fields, M.J.
Monitoring radioactive compounds in HPLC eluates: fraction collection versus on-line detection 342(1985)25
Robinson, J.S., see Owens, J.A. 338(1985)303
Robinson, Y., Gerry, N. and Vose, C.W.
Simultaneous HPLC assay for 5-(2-bromo-*E*-ethenyl)-2'-deoxyuridine and its metabolite, 5-(2-bromo-*E*-ethenyl)uracil, in plasma 338(1985)219
Roche, M., see Meinard, C. 349(1985)105
Rochemont, J., see Lazure, C. 326(1985)339
Rochette, J., see Bosisio, A.B. 330(1985)299
Rodick, C.L., see Edwards, G. 343(1985)224
Rodriguez, C., see Julien, C. 344(1985)51
Rodriguez, H.
Automated on-line identification of phenylthiohydantoin-amino acids from a vapor phase protein sequencer 350(1985)217
Rodriguez, P.A., Eddy, C.L. and Culbertson, C.R.
Temperature stabilization of a commercial photoionization detector. Performance evaluation 330(1985)365
Roemen, T.H.M. and Van der Vusse, G.J.
Application of silica gel CC in the assessment of non-esterified fatty acids and phosphoglycerides in myocardial tissue 344(1985)304
Roeraade, J.
Automated monitoring of organic trace components in water. I. Continuous flow extraction together with on-line capillary GC 330(1985)263
Roets, E., see Kibwage, I.O. 330(1985)275, 346(1985)309
Rogers, L.B., see McNally, M.E. 331(1985)23
Rogers, L.B., see Men, L.-C. 347(1985)39

Rogers, L.B., see Parekh, N.J. 330(1985)19
Rogers, L.B., see Shah, P. 329(1985)5
Rogers, M.E., Adlard, M.W., Saunders, G. and Holt, G.
High-performance liquid affinity chromatography of β-lactamase 326(1985)163
Rokushika, S., Huang, D.Y., Qui, Z.Y. and Hatano, H.
Polyallylamine-coated silica gel microbore column for LC 332(1985)15
Rokushika, S., see Naikwadi, K.P. 331(1985)69
Romano, G., see Musumarra, G. 350(1985)151
Romeo, G., see Bousquet, E. 344(1985)325
Rónai-Lukács, S., Kozma, M. and Vereczkey, L.
HPLC method for quantitative determination of Yutac, a new antiarrhythmic agent, in dog plasma 338(1985)428
Ronchetti, M., Cartoni, G. and Zoccolillo, L.
Determination of polynuclear aromatic hydrocarbons in petroleum oils by combination of TLC and GC-MS 348(1985)159
Rosa, J. de la, see Hirschberger, L.L. 343(1985)303
Rosati, F., see Cartoni, G.P. 330(1985)315
Rosini, C., Altemura, P., Pini, D., Bertucci, C., Zullino, G. and Salvadori, P.
Cinchona alkaloids for preparing new, easily accessible chiral stationary phases. II. Resolution of binaphthol derivatives on silica-supported quinine 348(1985)79
Rosseel, M.T., see Teirlynck, O.A. 342(1985)399
Rössle, C., see Kohse, K.P. 344(1985)319
Rosso, A., see Ravinale, E. 330(1985)415
Rossum, J. van, see Daenens, P. 342(1985)79
Roth, E., see Hamilton, G. 341(1985)411
Rotocki, P. and Drozdowicz, B.
Effects of contact potentials on the pulsed electron-capture detector 349(1985)155
Rouan, M.C.
Antibiotic monitoring in body fluids (Review) 340(1985)361
Roussel, P., see Houdret, N. 343(1985)437
Roussel, P., see Lhermitte, M. 342(1985)416
Rovei, V., see Vajta, S. 343(1985)239
Roy, S.K., McGregor, W.C. and Orichowskyj, S.T.
Automated high-performance immunosorbent assay for recombinant leukocyte A interferon 327(1985)189
Rubio, F. and Garland, W.A.
Determination of acetyltrimoprostil and its metabolite trimoprostil in human or dog plasma by GC-negative-ion chemical ionization MS 339(1985)313
Ruckmick, S.C. and Hurtubise, R.J.
Class separation of polycyclic aromatic hydrocarbons, nitrogen heterocycles and hydroxyl aromatics by LC 331(1985)55
Rühl, C. and Reusch, J.
Analysis of volatile nitrosamines by microbore HPLC and thermal energy analyser detection 328(1985)362
Rumble, R.H. and Roberts, M.S.
Determination of benzylpenicillin in plasma and urine by HPLC 342(1985)436
Rusconi, L. and Montecucchi, P.C.
Reversed-phase HPLC of natural and synthetic sauvagines 346(1985)390
Rusconi, L., Perseo, G., Franzoi, L. and Montecucchi, P.C.
Reversed-phase HPLC characterization of a novel proline-rich tryptophyllin 349(1985)117
Rushing, L.G., Hansen, E.B. Jr. and Thompson, H.C. Jr.
Analysis of D-penicillamine by GC utilizing nitrogen-phosphorus detection 337(1985)37
Russell, J.W., see Gammans, R.E. 339(1985)303
Ryan, P.J. and Honeyman, T.W.
Formation and separation of fluorescent derivatives of diacylglycerols 331(1985)177
Ryman, T., see Beck, O. 337(1985)402

S

Sabot, J.F., Deruaz, D., Dechaud, H., Bernard, P. and Pinatel, H.
Determination of plasma testosterone by mass fragmentography using [3,4-^{13}C]testosterone as an internal standard 339(1985)233
Šafařík, L., see Stránský, Z. 342(1985)167
Saffen, D., see Alpert, C.A. 326(1985)363
Sagara, K., Ito, Y., Oshima, T., Misaki, T., Murayama, H. and Itokawa, H.
Simultaneous determination of baicalein, wogonin, oroxylin-A and their glucuronides in Scutellariae Radix by ion-pair HPLC 328(1985)289
Saindrenan, P., Darakis, G. and Bompeix, G.
Determination of ethyl phosphite, phosphite and phosphate in plant tissues by anion-exchange HPLC and GC 347(1985)267
Saito, A., see Chung, T.G. 339(1985)67
Saito, A., see Niwa, T. 336(1985)345
Saito, M., see Hoshino, T. 332(1985)139
Saito, M., see Sugiyama, K. 332(1985)107
Saitoh, H. and Oikawa, K.
Simultaneous determination of iron(II) and -(III) by ion chromatography with post-column reaction 329(1985)247
Sakabe, Y., see Terada, H. 346(1985)333, 348(1985)379
Sakai, K., see Shijo, Y. 333(1985)133
Sakai, T., see Antoku, Y. 342(1985)359
Sakakibara, S., see Kumagaye, K.Y. 327(1985)327
Sakamoto, N., see Maeda, K. 345(1985)11
Sakamoto, N., see Shiraishi, S. 338(1985)51

Salama, A.I., see Lin, L. 341(1985)43
Sales, Y., see Necciari, J. 341(1985)202
Salkinoja-Salonen, M., see Pellinen, J. 328(1985)299
Saller, C.F., see Lin, L. 341(1985)43
Salvadori, P., see Rosini, C. 348(1985)79
Salzman, S.K., Sellers, M.S., Beckman, A.L., Naidu, R. and Macewen, G.D.
Determination of intraoperative plasma catecholamine concentrations using LC with electrochemical detection 343(1985)285
Samperi, R., see Andreolini, F. 337(1985)379
Sandlin, J.L. and Ito, Y.
Large-scale preparative countercurrent chromatography for separation of polar compounds 348(1985)131
Sane, R.T., Desai, S.V., Sonawne, K.K. and Nayak, V.G.
Quantitative GC determination of tamoxifen citrate in pharmaceuticals 331(1985)432
Sangó, C., see Skarping, G. 346(1985)191
Sannia, G., see Cubellis, M.V. 329(1985)406
Santa María, I., Carmi, J.D., Haddad, G.R. and Cifuentes, M.
Open-tubular columns support-coated *in situ* with cross-linked SE-54 stationary phase 329(1985)123
Santagostino, G., see Schinelli, S. 338(1985)396
Sappa, E., see Castiglioni, M. 349(1985)173
Sardessai, M., see Lim, H.K. 328(1985)378
Sarver, E.W., see Bossle, P.C. 330(1985)388
Sasabe, H., see Miyamoto, G. 338(1985)450
Sasaki, H., see Kuwata, K. 332(1985)29
Sasaki, H., see Sasaki, I. 332(1985)237
Sasaki, I., Makino, K., Takeuchi, T. and Sasaki, H.
Use of Toyopearl as a support for the immobilization of urease 332(1985)237
Sasaki, K., see Mikasa, H. 343(1985)179
Sasaki, M., see Chung, T.G. 339(1985)67
Sassard, J., see Cuisinaud, G. 341(1985)97
Sassard, J., see Julien, C. 344(1985)51
Sassen, W. von, see Von Sassen, W. 338(1985)113
Sato, A., see Shimada, K. 339(1985)331
Sato, H., see Tanaka, Y. 347(1985)275
Sato, K., see Yonezawa, K. 339(1985)219
Sato, M., see Hatake, K. 344(1985)339
Sato, T., see Kadofuku, T. 343(1985)51
Sato, T., see Yamada, K. 339(1985)210
Satoh, A., see Oyama, H. 338(1985)71
Saunders, G., see Rogers, M.E. 326(1985)163
Sauziere, J., see Le Hoang, M.D. 346(1985)382
Savaraj, N., see Lu, K. 345(1985)408
Sävenhed, R., Borén, H. and Grimvall, A.
Stripping analysis and chromatographic sniffing for the source identification of odorous compounds in drinking water 328(1985)219
Sävenhed, R., see Borén, H. 348(1985)67
Savolainen, H., see Pfäffli, P. 337(1985)146
Sawada, S., see Sogami, M. 332(1985)19
Saynor, D.A., see Colthup, P.V. 345(1985)111
Scalia, S. and Morgan, E.D.
Simultaneous determination of free and conjugated ecdysteroids by LC 346(1985)301
Scandurra, R., see Laganà, A. 345(1985)251
Scanlon, D., Haralambidis, J., Southwell, C., Turton, J. and Tregear, G.
Purification of synthetic oligodeoxyribonucleotides by ion-exchange HPLC 336(1985)189
Scarlata, G., see Musumarra, G. 350(1985)151
Scarpa, A., see Beers, M.F. 336(1985)380
Schaaf, L.J., Ferry, D.G., Hung, C.T., Perrier, D.G. and Edwards, I.R.
Analysis of 5'-deoxy-5-fluorouridine and 5-fluorouracil in human plasma and urine by HPLC 342(1985)303
Schade, U., see Shukla, A.K. 337(1985)231
Schafer-Mitchell, M., see Cook, L. 339(1985)293
Schallinger, L.E., Gray, J.E., Wagner, L.W., Knowlton, S. and Kirkland, J.J.
Preparative isolation of plasmid DNA with sedimentation field flow fractionation 342(1985)67
Schamp, N., see Van der Straeten, D. 331(1985)207
Schauer, R., see Shukla, A.K. 337(1985)231
Schaufelberger, D. and Hostettmann, K.
Analytical and preparative reversed-phase LC of secoiridoid glycosides 346(1985)396
Scheler, F., see Schneider, K. 345(1985)19
Schepper, P. de, see Daenens, P. 342(1985)79
Schilcher, R.B., Young, J.D. and Baker, L.H.
Reversed-phase HPLC determination of tricyclic nucleoside and tricyclic nucleoside 5'-phosphate in biological specimens 337(1985)55
Schiller, E., see Ehninger, G. 342(1985)119
Schilsky, R.L. and Ordway, F.S.
Simultaneous determination of cytosine arabinoside, its nucleotides and metabolites by ion-pair HPLC 337(1985)63
Schinelli, S., Santagostino, G., Frattini, P., Cucchi, M.L. and Corona, G.L.
Assay of 3-methoxy-4-hydroxyphenylglycol in human plasma using HPLC with amperometric detection 338(1985)396
Schladitz-Keil, G., Spahn, H. and Mutschler, E.
Fluorimetric determination of the quaternary compound trospium and its metabolite in biological material after derivatization with benoxaprofen chloride 345(1985)99
Schlimme, E., see Kemper, K. 336(1985)374
Schmerr, M.J.F., Goodwin, K.R., Lehmkuhl, H.D. and Cutlip, R.C.
Preparation of sheep and cattle immunoglobulins with antibody activity by HPLC 326(1985)225
Schmidt, A., see Krauss, F. 348(1985)296
Schmitz, F.P., Hilgers, H., Lorenschat, B. and Klesper, E.
Separation of oligomers with UV-absorbing side groups by supercritical fluid

chromatography using eluent gradients 346(1985)69
Schmuck, M.N., see Gooding, K.M. 327(1985)139
Schneider, K., Neupert, M., Spiteller, G., Henning, H.V., Matthaei, D. and Scheler, F. GC of amino acids in urine and haemofiltrate 345(1985)19
Schneider, M., see Staveris, S. 339(1985)97
Schoemaker, T.J., see Van Belle, S.J.P. 337(1985)73
Scholze, H. Determination of phenylthiocarbamyl amino acids by reversed-phase HPLC 350(1985)453
Schootstra, R., see Storey, G.C.A. 341(1985)223
Schouten, T.J., see Van Baal, J.M. 336(1985)422
Schrieck, P. van der, see Frenz, J. 330(1985)1
Schryver, E.P. de, see De Schryver, E.P. 338(1985)389
Schuller, E., see Allinquant, B. 326(1985)281
Schulman, S.G., see Karnes, H.T. 338(1985)377
Schultz, J.R., see DiBella, E.E. 328(1985)101
Schulz, A., Wiedemann, K. and Bitsch, I. Stabilization of 5-methyltetrahydrofolic acid and subsequent analysis by reversed-phase HPLC 328(1985)417
Schutter, J.A. de, see De Schutter, J.A. 350(1985)135
Schwadrohn, G., see Balansard, G. 348(1985)299
Schwandt, P., Richter, W.O., Heinemann, V. and Weisweiler, P. Characterization of human apolipoprotein A-I by reversed-phase HPLC 345(1985)145
Schwartz, B.D., see McCourt, D.W. 327(1985)9
Schwartz, J.B., see Abernethy, D.R. 342(1985)216
Schwarz, V., Deyl, Z. and Macek, K. Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401
Schwarz, W. HPLC determination of urocanic acid in stripped human stratum corneum 342(1985)247
Schwarzenbach, R. Adsorption chromatographic separations on buffered silica gel (Review) 334(1985)35
Schwedass, A., see Räder, K. 344(1985)416
Schweer, H., Kammer, J. and Seyberth, H.W. Simultaneous determination of prostanoids in plasma by GC-negative-ion chemical-ionization MS 338(1985)273
Scott, P.J., see Poll, D. 343(1985)149
Scrivens, J.H., see Catlow, D.A. 328(1985)167
Sculier, J.P., see Brassinne, C. 339(1985)445
Sebille, B., see Mahieu, J.-P. 327(1985)313
Šefer, I., see Petrović, S.M. 348(1985)49
Segelman, F.H., see Kucharczyk, N. 340(1985)243
Seidah, N.G., see Lazure, C. 326(1985)339
Seiffarth, K., see Döring, C.E. 348(1985)430
Seiler, N. and Knödgen, B. Determination of polyamines and related compounds by reversed-phase HPLC: improved separation systems 339(1985)45
Seiler, N. and Knodgen, B. Determination of amino acids by separation of their ion pairs with dodecyl sulphate 341(1985)11
Seki, T., Yamaguchi, Y., Noguchi, K. and Yanagihara, Y. Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9
Seki, T., Yamaguchi, Y., Noguchi, K. and Yanagihara, Y. Separation of ascorbic acid, dehydroascorbic acid, diketogulonic acid and glucose by isocratic elution from a column of a hydrophilic gel 332(1985)283
Sekine, Y., see Hayashi, K. 338(1985)161
Sekiya, C., see Taniguchi, N. 339(1985)394
Selan, F., Blair, N. and Evans, M.A. HPLC analysis for fluorescein and fluorescein monoglucuronide in plasma 338(1985)213
Sellergren, B., Ekberg, B. and Mosbach, K. Molecular imprinting of amino acid derivatives in macroporous polymers. Demonstration of substrate- and enantio-selectivity by chromatographic resolution of racemic mixtures of amino acid derivatives 347(1985)1
Sellergren, B., see Ekberg, B. 333(1985)211
Sellergren, B., see Glad, M. 347(1985)11
Sellers, M.S., see Salzman, S.K. 343(1985)285
Selosse, E.J.-M. and Reilly, P.J. Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253
Seltzer, J.L., Eschbach, M.L. and Eisen, A.Z. Purification of gelatin-specific neutral protease from human skin by conventional and high-performance LC 326(1985)147
Senda, M., see Hoshino, T. 332(1985)139
Senda, M., see Sugiyama, K. 332(1985)107
Senda, M., see Yoshioka, M. 344(1985)345
Senft, V. Simple determination of trichloroacetic acid in urine using head space GC: a suitable method for monitoring exposure to trichloroethylene 337(1985)126
Sentissi, A., O'Connell, K. and Giese, R.W. Relative precision in the determination of some structurally-similar O-pentafluorobenzyloximes by ultratrace GC 331(1985)396
Sepulveda, F.I., see Yang, C.-Y. 346(1985)413
Sequeira, R.P., see Chaiken, I.M. 336(1985)63
Serfontein, W.J., see Ubbink, J.B. 342(1985)277
Serpinet, J., see Claudy, P. 329(1985)331
Serpinet, J., see Gonnet, C. 350(1985)227
Sevenich, G.J. and Fritz, J.S. Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147
Seyberth, H.W., see Schweer, H. 338(1985)273
Seyer, J.M., see Heinemann, M.A. 329(1985)295

Smith, A. and Jones, B.
Impact of [illegible] on [illegible] in [illegible] and [illegible]
Lee, J. K. and Wong, L. M.
[illegible] study of [illegible] in [illegible] and [illegible]
Patel, R., Shah, M. and Desai, P. N.
Influence of [illegible] on the [illegible] of [illegible] in [illegible] [illegible]
Johnson, M. A., Brown, K. L., Davis, S. R. and Wilson, T. J.
Optimization of [illegible] (Fabaceae) in [illegible] [illegible] [illegible]
Garcia, O., van Dongen, R. G. S. [illegible]
Simonsen, J. K., van Leeuwen, M. C. [illegible]
Kumar, S. P., van Dijkstra, L. H. [illegible]
Andersen, B. O., van Berkum, J. J. [illegible]
Rodriguez, C. M., van Straaten, H. [illegible]
Schoonhoven, L., van Loon, J. [illegible]
Takahashi, Y., van Kampen, C. [illegible]
Hernandez, A., van Steenbergen, T. [illegible]
Bakker, J., van Dongen, M. [illegible]
Okoye, V. and Nwosu, K.
[illegible] of [illegible], [illegible], [illegible], [illegible] and [illegible] [illegible] with [illegible] [illegible]
Ivanova, E. and Petrov, G.
[illegible] of [illegible] on [illegible] and [illegible] of [illegible] [illegible] in [illegible] [illegible]
Kowalski, P., Wiśniewska, M. and Lis, T.
[illegible] of [illegible] on [illegible] [illegible] in the [illegible] [illegible]
Tanaka, Y. and Watanabe, K.
[illegible] [illegible] [illegible] [

Simmaco, M., Barra, D. and Bossa, F.
Separation of phenylthiohydantoin-amino acids by HPLC and some applications in dansyl Edman sequence analysis 349(1985)99
Simmonds, H.A., see Morris, G.S. 344(1985)101
Simmonds, R.J.
Determination of atracurium, laudanosine and related compounds in plasma by HPLC 343(1985)431
Simon, J. and Barla, F.
Application of capillary GC in the stability control of phthalate-ester-type poly(vinyl chloride) plasticisers treated with light and plasma 329(1985)389
Simon, M., see De Vries, J.X. 338(1985)325
Simon, P., see Diquet, B. 336(1985)415
Simpson, R., see Cunico, R.L. 336(1985)105
Singh, A.K. and Drewes, L.R.
Improved analysis of acetylcholine and choline in canine brain and blood samples by capillary GC-MS 339(1985)170
Sinha, D.K., see Morreal, C.E. 345(1985)380
Sinkule, J.A., see Choi, K.E. 345(1985)197
Sintov, A., see Yagen, B. 343(1985)67
Sioufi, A. and Pommier, F.
Quantitative determination of nitroglycerin in human plasma by capillary GC with electron-capture detection 339(1985)117
Sioufi, A., see Colussi, D. 339(1985)157
Sisco, W.R., Rittenhouse, C.T. and Everhart, L.A.
Simultaneous HPLC stability-indicating analysis of acetaminophen and codeine phosphate in tablets and capsules 348(1985)253
Siu, K.W.M., Gardner, G.J. and Berman, S.S.
Hypercoulometry in a pulsed conventional electron-capture detector 330(1985)87
Sivorinovsky, G., see Binder, S.R. 336(1985)173
Sjöberg, P. and Bondesson, U.
Determination of di(2-ethylhexyl) phthalate and four of its metabolites in blood plasma by GC-MS 344(1985)167
Sjöquist, B. and Ljungquist, C.
Identification and quantification of 1-carboxysalsolinol and salsolinol in biological samples by GC-MS 343(1985)1
Skalická, O., see Kuchař, M. 333(1985)21
Skarping, G., Renman, L., Sangö, C., Mathiasson, L. and Dalene, M.
Capillary GC method for the determination of complex mixture of isocyanates and amines 346(1985)191
Skarping, G., Smith, B.E.F. and Dalene, M.
Trace analysis of amines and isocyanates using glass capillary GC and selective detection. V. Direct determination of isocyanates using nitrogen-selective and electron-capture detection 331(1985)331
Skarping, G., see Dalene, M. 331(1985)321
Skee, D., see McCoy, L. 344(1985)211
Skelly, J.P., see Minagawa, K. 343(1985)231
Skidmore, L.F., see Colthup, P.V. 345(1985)111
Skrabalak, D.S., Cuddy, K.K. and Henion, J.D.
Quantitative determination of betamethasone and its major metabolite in equine urine by micro-LC-MS 341(1985)261
Skrlletz, U., see Gleispach, H. 344(1985)11
Slack, J.A. and Goddard, C.
Antitumour imidazotetrazines. VII. Quantitative analysis of mitozolomide in biological fluids by HPLC 337(1985)178
Slevin, M.L., see Harvey, V.J. 339(1985)419
Slobodzian, D.K., Hsieh, J.Y.-K. and Bayne, W.F.
HPLC determination of 3-hydroxymethyl-dibenzo[*b,f*]thiepin 5,5-dioxide and its acid metabolite in human plasma and urine 338(1985)253
Slobodzian, D.K., Hsieh, J.Y.-K. and Bayne, W.F.
Simultaneous determination of tramlast and metabolites in plasma and urine using HPLC 345(1985)345
Slopsema, K., see Welling, G.W. 326(1985)173
Smallwood, A.W., see Sollenberg, J. 343(1985)175
Smet, M. de, see De Smet, M. 345(1985)309
Smet, M. de, see Musch, G. 348(1985)97
Smith, A.L., see Weber, A.F. 337(1985)434
Smith, B.E.F., see Skarping, G. 331(1985)331
Smith, I.J., see Hartley, R. 342(1985)105
Smith, J.A. and O'Hare, M.J.
Ion-exchange HPLC of mouse epidermal growth factor and its congeners: mobile phase optimization with ion-pairing additives 345(1985)168
Smith, J.M., see Greene, S.V. 343(1985)160
Smith, L.A., Norman, H.A., Cho, S.H. and Thompson, G.A. Jr.
Isolation and quantitative analysis of phosphatidylglycerol and glycolipid molecular species using reversed-phase HPLC with flame ionization detection 346(1985)291
Smith, L.L., see Teng, J.I. 339(1985)35, 350(1985)445
Smith, R.J., Haken, J.K. and Wainwright, M.S.
A problem with Kováts' retention index equation 331(1985)389
Smith, R.J., Haken, J.K. and Wainwright, M.S.
Estimation of dead time and calculation of Kováts indices (Review) 334(1985)95
Smith, R.J., Haken, J.K., Wainwright, M.S. and Madden, B.G.
Comparison of mathematical methods for the calculation of Kováts indices 328(1985)11
Smith, R.L. and Walker, D.D.
HPLC determination of acyclovir in serum 343(1985)203
Smith, R.W., see Parker, C.E. 344(1985)378, 347(1985)61
Smith, S.L., Novotný, M. and Karmen, A.
Elevation of certain polyols in the cerebrospinal fluid of patients with multiple sclerosis 336(1985)351

Smithers, J.A., Lang, J.F. and Okerholm, R.A.
Quantitative analysis of vigabatrin in plasma and urine by reversed-phase HPLC 341(1985)232
Smolková-Keulemansová, E., see Krýsl, S. 349(1985)167
Smolková-Keulemansová, E., see Macek, J. 333(1985)309
Smolková-Keulemansová, E., see Tomková, H. 329(1985)113
Smyth, R.D., see Forgue, S.T. 342(1985)221
Smythe, G.A., see Duncan, M.W. 336(1985)199
Snoussi, M., see Bressolle, F. 343(1985)443
Snyder, L.R., see Banes, A.J. 326(1985)419
Snyder, L.R., see Stadalius, M.A. 327(1985)27, 327(1985)93
So, N., Chandra, D.P., Alexander, I.S., Webster, V.J. and O'Gorman Hughes, D.W.
Determination of serum methotrexate and 7-hydroxymethotrexate concentrations. Method evaluation showing advantages of HPLC 337(1985)81
Soffer, E.F., see Leelavathi, D.E. 339(1985)105
Sogami, M., Era, S., Nagaoka, S., Kuwata, K., Kida, K., Shigemi, J., Miura, K., Suzuki, E., Muto, Y., Tomita, E., Hayano, S., Sawada, S., Noguchi, K. and Miyata, S.
HPLC studies on non-mercapt mercapt conversion of human serum albumin II 332(1985)19
Sogami, M., see Kuwata, K. 332(1985)29
Soine, W.H., see Bhargava, V.O. 343(1985)219
Sollenberg, J., Phipps, F.C., Stringer, B. and Lowry, L.K.
Determination of methylhippuric acid in human urine by HPLC and by isotachophoresis 343(1985)419
Sollenberg, J., Smallwood, A.W. and Lowry, L.K.
Determination of mandelic and phenylglyoxylic acids in rat urine by HPLC and by isotachophoresis 343(1985)175
Soltés, L., see Marko, V. 339(1985)410
Sommer, W., see Jacob, K. 349(1985)283
Sonawne, K.K., see Sane, R.T. 331(1985)432
Sonnenberg, J. and Pingoud, V.
Purification of radioiodinated photoactivable glucagon by isocratic HPLC 331(1985)133
Sonnino, S., see Gazzotti, G. 348(1985)371
Sønsterud, B., see Stray, H. 349(1985)331
Soparkar, C.N.S., Malison, R.T. and Burton, J.
Development of a system to simplify identification of peptides and their catabolites in a HPLC effluent 337(1985)110
Sørensen, M.B., see Angelo, H.R. 345(1985)413
Sorrell, T.C., see Müller, M. 343(1985)213
Sotirhos, N., Thörngren, C. and Herslöf, B.
Reversed-phase HPLC separation and mass detection of individual phospholipid classes 331(1985)313
Soubeyrand, J., see Désage, M. 336(1985)285
Soumarmon, A., see Robert, J.C. 338(1985)315
Soun, A., see Désage, M. 336(1985)285
Southwell, C., see Scanlon, D. 336(1985)189
Spahn, H., Prinoth, M. and Mutschler, E.
Determination of pindolol in plasma and urine by TLC 342(1985)458
Spahn, H., see Schladitz-Keil, G. 345(1985)99
Spandau, D.J., see Gaffney, J.S. 347(1985)121
Spatola, A.F. and Benovitz, D.E.
Improved detection limits in the analysis of tyrosine-containing polypeptide hormones by using electrochemical detection 327(1985)165
Speedie, M.K., Wong, D.L. and Ciaranello, R.D.
Characterization of the subunit structure of dopamine β-hydroxylase by anion-exchange HPLC 327(1985)351
Spener, F., see Shukla, V.K.S. 348(1985)441
Spicer, W.J., see Huysmans, M.B. 337(1985)223
Spiteller, G., see Schneider, K. 345(1985)19
Spragg, B.P., see Littlewood, T.J. 336(1985)434
Spraul, M., see Albert, K. 346(1985)17
Spriggle, J.E., see Metz, P.A. 330(1985)307
Stacey, C., see Colgan, S.T. 333(1985)349
Stadalius, M.A., Gold, H.S. and Snyder, L.R.
Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27
Stadalius, M.A., Quarry, M.A. and Snyder, L.R.
Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Application to method development and the choice of column configuration 327(1985)93
Stan, H.-J. and Lipinski, J.
Microcomputer programming in BASIC for the evaluation of capillary GC in the analysis of pesticide residues. I. MATRIXCOMP: a program that facilitates the recognition of interfering peaks from the biological matrix 349(1985)49
Stancher, B., see Zonta, F. 329(1985)257
Stanley, M., see Novak, E.A. 339(1985)457
Staszewski, R., see Wardencki, W. 329(1985)128
Staveris, S., Blaise, P., Efthymiopoulos, C., Schneider, M., Jamet, G., Jung, L. and Koffel, J.C.
Quantitative determination of tertatolol in biological fluids by GC-MS 339(1985)97
Staveris, S., Jung, L., Jamet, G. and Koffel, J.C.
Quantitative analysis of veralipride in plasma and urine by GC-MS and GC with flame-ionization detection 338(1985)79
Stec, W.J., Zon, G. and Uznański, B.
Reversed-phase HPLC separation of diastereomeric phosphorothioate analogues of oligodeoxyribonucleotides and other backbone-modified congeners of DNA 326(1985)263
Steele, R.D., see Fell, D. 345(1985)150
Štefek, M., see Marko, V. 339(1985)410
Stehle, P. and Fürst, P.
Isotachophoretic control of peptide

synthesis and purification. A novel approach using UV detection at 206 nm 346(1985)271
Stein, S., see Lahm, H.-W. 326(1985)357
Stephany, R.W., see Jansen, E.H.J.M. 347(1985)375, 347(1985)379
Stephenson, G.P., see Breslin, F. 329(1985)434
Stern, M., see Beers, M.F. 336(1985)380
Stetson, P.L., Shukla, U.A., Amin, P.R. and Ensminger, W.D.
HPLC method for the determination of bromodeoxyuridine and its major metabolite, bromouracil, in biological fluids 341(1985)217
Stetson, P.L., Shukla, U.A. and Ensminger, W.D.
Sensitive HPLC method for the determination of 5-fluorouracil in plasma 344(1985)385
Stetson, P.L., see Shukla, U.A. 342(1985)406
Stevenson, D., see Read, J. 341(1985)437
Sticher, O., see Erdelmeier, C.A.J. 346(1985)456
Stierlin, H., see Francotte, E. 346(1985)321
Stikkelman, R.M., see Tjioe, T.T. 330(1985)412
Stipanuk, M.H., see Hirschberger, L.L. 343(1985)305
Stivaletta, L.A., see Goldstein, S. 339(1985)388
Stocchi, V., Cucchiarini, L., Piccoli, G. and Magnani, M.
Complete HPLC separation of 4-N,N-dimethylaminoazobenzene-4'-thiohydantoin and 4-dimethylaminoazobenzene-4'-sulphonyl chloride amino acids utilizing the same reversed-phase column at room temperature 349(1985)77
Stohs, S., see Cook, L. 339(1985)293
Storey, G.C.A., Schootstra, R. and Henry, J.A.
Measurement of meptazinol in plasma by HPLC with electrochemical detection 341(1985)223
Storme, G.A., see De Smet, M. 345(1985)309
Storshane, R.M., see Brown, R.R. 339(1985)440
Stout, M., Ravindranath, Y. and Kauffman, R.
HPLC assay for methotrexate utilizing a cold acetonitrile purification and separation of plasma or cerebrospinal fluid 342(1985)424
Stout, R.W. and DeStefano, J.J.
A new, stabilized, hydrophilic silica packing for the high-performance gel chromatography of macromolecules 326(1985)63
Stout, R.W., see Link, G.W. Jr. 331(1985)253
Straathof, A.J.J., see Tjioe, T.T. 330(1985)412
Strachan, M.G. and Johns, R.B.
Determination of molecular weights of coal derived liquids by GPC-HPLC 329(1985)65
Straeten, D. van der, see Van der Straeten, D. 331(1985)207
Stransky, Z., Chmela, Z., Peč, P. and Šafařík, L.
Determination of trimecaine metabolites in blood plasma by capillary isotachophoresis 342(1985)367
Stray, H., Mikalsen, A., Sønsterud, B. and Oehme, M.
Preseparation of polar, substituted, polycyclic aromatic hydrocarbons in aerosol extracts by HPLC on acidic- or basic-buffered silica 349(1985)331
Street, J.M., Trafford, D.J.H. and Makin, H.L.J.
Extraction and fractionation of bile acids and their conjugates using pre-packed microparticulate silica cartridges (Sep-Pak Sil and Bond-Elut C_{18}) 343(1985)259
Streinz, L., see Vaněk, T. 347(1985)188
Stringer, B., see Sollenberg, J. 343(1985)419
Strojny, N., Conzentino, P. and De Silva, J.A.F.
Determination of coumermycin A_1 in plasma by reversed-phase HPLC analysis 342(1985)145
Strojny, N. and De Silva, J.A.F.
Determination of diclofensine, an antidepressant agent, and its major metabolites in human plasma by HPLC with fluorometric detection 341(1985)313
Strojny, N., Puglisi, C.V. and De Silva, J.A.F.
Determination of the antiallergenic agent, N-[4-(1H-imidazol-1-yl)butyl]-2-(1-methylethyl)-11-oxo-11H-pyrido[2,1-*b*]quinazoline-8-carboxamide, in plasma by reversed-phase HPLC analysis using fluorometric detection 336(1985)301
Strolin Benedetti, M., see Vajta, S. 343(1985)239
Strom, J., see Angelo, H.R. 345(1985)413
Stuart, J.D., see Huang, J.-X. 330(1985)181
Sturm, U., see König, W.A. 328(1985)357
Sturman, L.S., see Ricard, C.S. 326(1985)191
Succari, M., Foglietti, M.-J. and Percheron, F.
Two-step purification of human α_1-acid glycoprotein 341(1985)457
Suchár, G., see Podhradsky, D. 328(1985)402
Suckow, R.F. and Cooper, T.B.
Determination of amoxapine and metabolites in plasma by LC with electrochemical detection 338(1985)225
Sugimoto, I., see Suzuki, H. 341(1985)341
Sugimoto, T., see Takasugi, N. 342(1985)349
Sugita, S., see Kobayashi, A. 336(1985)410
Sugiyama, K., Saito, M., Hondo, T. and Senda, M.
New double-stage separation analysis method. Directly coupled laboratory-scale supercritical fluid extraction-supercritical fluid chromatography, monitored with a multiwavelength-UV detector 332(1985)107
Sugiyama, K., see Furuta, H. 337(1985)103
Sumi, M., see Tetsumi, T. 348(1985)389
Sun, Y., see Wang, T. 330(1985)167
Suprynowicz, Z., Buda, W.M., Mardarowicz, M. and Patrykiejew, A.
Correlations between retention on liquid crystalline stationary phases and chemical structure. I. Dimethylnaphthalenes 333(1985)11
Surridge, D.H., see Owen, J.A. 342(1985)333

Susa, J.B., see Boylan, J.M. 338(1985)192
Suzaki, K., see Goto, J. 343(1985)241, 348(1985)181
Suzuki, E., see Sogami, M. 332(1985)19
Suzuki, H., Fujiwara, S., Kondo, S. and Sugimoto, I.
Determination of nifedipine in human plasma by HPLC with electrochemical detection 341(1985)331
Suzuki, H., Yata, J., Kojima, K. and Nagatsu, T.
Simple and sensitive assay of dopamine β-hydroxylase in human cerebrospinal fluid by HPLC with electrochemical detection 341(1985)108
Suzuki, S., see Hobo, T. 336(1985)131
Suzuki, T., see Tokumura, A. 343(1985)138
Svartling, N., see Pfäffli, P. 345(1985)326
Svendsen, J.S., Whist, J.E. and Sydnes, L.K.
Absolute configuration of 3-hydroxyadipic acid in human urine 337(1985)9
Svensson, J.-O., Muchtar, A. and Ericsson, O.
Ion-pair HPLC determination of isoniazid and acetylisoniazid in plasma and urine. Application for acetylator phenotyping 341(1985)158
Svetashev, V.I. and Zhukova, N.V.
Analysis of labelled fatty acid methyl esters by argentation and reversed-phase two-dimensional TLC 320(1985)396
Swagzdis, J.E., Gifford, R. and Mico, B.A.
Determination of 4-(2-di-*n*-propylaminoethyl)-7-hydroxy-2(3H)-indolone and N-(2'-hydroxy-5'-(N,N-di-*n*-propylaminoethyl)-phenyl)-methane sulfonamide in plasma by HPLC with electrochemical detection 345(1985)203
Swaisgood, H.E. and Chaiken, I.M.
Quantitative affinity HPLC of neuroendocrine polypeptides using porous and non-porous glass derivatives 327(1985)93
Swaisgood, H.E., see Chaiken, I.M. 336(1985)63
Swarin, S.J., see Olson, K.L. 332(1985)337
Sweeney, D.J., Greig, N.H. and Rapoport, S.I.
HPLC analysis of melphalan in plasma, brain and peripheral tissue by *o*-phthalaldehyde derivatization and fluorescence detection 339(1985)424
Swinkels, J., see Penz, A.M. 327(1985)359
Swobodnik, W., Klüppelberg, U., Wechsler, J.G., Volz, M., Normann, C. and Ditschuneit, H.
Rapid and accurate reversed-phase HPLC determination of conjugated bile acids in human bile for routine clinical applications. Therapeutic control during gallstone dissolution therapy 339(1985)263
Sybilska, D., see Dębowski, J. 329(1985)206
Sybilska, D., see Dębowski, J. 349(1985)131
Sybilska, D., see Kościelski, T. 349(1985)3
Sydnes, L.K., see Svendsen, J.S. 337(1985)9
Szabó, L.D., see Csató, E. 348(1985)230
Szafranek, J., see Kuśmierz, J. 331(1985)219
Szakács-Dobozi, M., Halász, A. and Vámos-Vigyázó, L.
Isoelectric focusing separation of *Gliocladium* enzyme components 361(1985)310
Szczepaniak, W., see Nawrocki, J. 339(1985)379
Szczepanik-van Leeuwen, P.A.
Improved GC-MS assay for haloperidol utilizing ammonia chemical ionization and selected-ion monitoring 339(1985)321
Szederkényi, F., Horváth, G., Ambrus, G., Tóth-Sarudy, É., Czira, G. and Tamás, J.
Selective trimethylsilylation for the determination of 16(*S*)-amino-$PGF_{2\alpha}$ methyl ester by GLC 333(1985)404
Szepesi, G., see Gazdag, M. 328(1985)279, 347(1985)284
Szogyi, M., see Cserháti, T. 349(1985)295
Szotyory, L., see Siklósi, A. 347(1985)382
Szymanowski, J., Voelkel, A., Beger, J. and Merkwitz, H.
Increments of some polarity parameters for polyoxyethylene glycol dialkyl ethers and for some of their sulphur analogues 336(1985)1.
Szymanowski, J., see Beger, J. 333(1985)319

T

Táborský, J., see Horna, A. 348(1985)141
Tabuchi, F., see Tatsuhara, T. 339(1985)149
Tadano, K., Yuhki, Y., Aoki, I., Miyazaki, K. and Arita, T.
HPLC determination of tranilast in plasma 341(1985)228
Tagliaro, F., Frigerio, A., Dorizzi, R., Lubli, G. and Marigo, M.
LC with pre-column dansyl derivatization and fluorimetric detection applied to the assay of morphine in biological samples 336(1985)323
Taira, H., Mizutani, S., Kurauchi, O., Narita, O. and Tomoda, Y.
Rapid purification of human placental angiotensin I converting enzyme by captopril affinity chromatography 339(1985)366
Tajima, K., see Mizutani, T. 333(1985)171
Takada, H. and Ishiwatari, R.
Quantitation of long-chain alkylbenzenes in environmental samples by silica gel CC and high-resolution GC 346(1985)281
Takagi, N., see Tsuchiya, H. 339(1985)59
Takahashi, M., see Komuro, C. 338(1985)206
Takahashi, N., Ishioka, N., Takahashi, Y. and Putnam, F.W.
Automated tandem HPLC system for

separation of extremely complex peptide mixtures [By: ...]
Takahashi, N., Takahashi, Y., Henry, M.E. and Putnam, F.W.
Purification of hemopexin and its domain fragments by affinity chromatography and HPLC [...]
Takahashi, N., see Ishii, I. 343(1985)194
Takamoto, T., see Chang, J.-J. 339(1985)35
Takahashi, Y., see Takahashi, N. 326(1985)375
Takai, M., see Kumagaye, K.Y. 327(1985)327
Takai, N., see Matsushima, Y. 332(1985)265
Takata, Y., see Fujioka, K. [illegible]
Takakusagi, K., see Matsushima, Y. [illegible]
Takamatsu, M., see Kida, N. 328(1985)304
Takamu, H., see Kawada, T. 329(1985)94
Takasugi, N., Mafune, E., Yokokawa, S., Teruyama, K., Tsuchiya, K. and Sugimoto, T.
Determination of malotilate and its metabolites in plasma and urine [...]
Takata, Y., Take, M., Ito, M. and Murao, T.
Low dead volume coulometric detector for LC 332(1985)117
Takauchi, K., see Tokumura, A. 343(1985)135
Takayanagi, H., Ogra, H., Fujimura, K. and Ando, T.
Ligand-exchange GC separation of aromatic sulphonates [...]
Takayanagi, H., Hashimoto, M., Fujimura, K. and Ando, T.
Ligand-exchange GC of cationic surfactants [...]
Takayanagi, H., see Fujimura, K. 350(1985)371
Takei, S., see Moriguchi, S. 350(1985)13
Takei, S., see Narita, K. 330(1985)293
Takemori, Y., see Hara, S. 344(1985)93
Takeuchi, M., see Yoshioka, S. 342(1985)69
Takeuchi, T., Asai, M., Hashimoto, Y., Watanabe, K. and Ishii, D.
Application of micro HPLC to the separation of chiral amino acids [...]
Takeuchi, T., see Ishii, D. 351(1985)8
Takeuchi, T., see Ito, Y. 346(1985)61
Takeuchi, T., see Otani, H. 353(1985)43
Takeuchi, T., see Sasaki, S. 352(1985)297
Taki, M., see Tanaka, Y. 332(1985)17
Tamás, J., see Sandermann, F. 338(1985)xxx
Tarresch, A.H.H., Hanna, M.H. and Somers, S.
GC determination of urea aromatics in formulations 328(1985)207
Tamura, Z., see Yoshioka, M. 344(1985)345
Tan, L., see Muto, N. 329(1985)137
Tanaka, H., see Iwasaki, S. 342(1985)152
Tanaka, H., see Nakagawa, T. 330(1985)43
Tanaka, M., Mio, T. and Shono, T.
Mercaptoalkylcelluloses as mobile phase additives in LC 330(1985)253
Tanaka, M., see Furuno, K. 346(1985)213
Tanaka, M., see Tetsumi, T. 346(1985)389
Tanaka, N., Kimoto, E., Araki, M. and Tsuda, T.
Simultaneous separation of chemically bonded saturated phase with monolayer surface coverage and high molecular basis and its application to polymer, macromolecular polymers 332(1985)91
Tanaka, T., Hasegawa, K., Matsuo, Y., Lee, C.-S. and Imai, Y.
Rapid and sensitive determination of amphetamine in urine by HPLC with fluorescence detection 338(1985)xxx
Tanaka, T., see Tsuruta, K. 338(1985)xxx
Tanaka, Y., Seki, H., Kaguei, A. and Kimura, I.
Separation of ganglioside-mixed isomers by HPLC and identification by ^{13}C nuclear magnetic resonance spectroscopy [...]
Tanaka, Y., see Nagashima, H. [illegible]
Tanaka, A.K., see Fujimura, S.H. [illegible]
Tanigawa, N., Yamashita, Y., Ono, H., Kawashima, K., Hirata, K. and Sekine, C.
Chromatography of hydrophilic proteolytic enzymes between 100-80 cells and human primary hepatocytes on Phenomenex reagent multi-suggestion [illegible] [illegible] 338(1985)xxx
Tanimoto, K., see Inoue, T. [illegible]
Tannenbaum, S.R., see Baker, I. 329(1985)202
Tanner, R.L., see Gaffney, J.S. 347(1985)221
Tarnowski, S.J., see Felix, A.M. 327(1985)334
Tarui, S., see Hayashi, M. 339(1985)245
Tarvin, T.L., see Hollingmeyer, K.A. 338(1985)xxx
Tas, A.C., see Van der Greef, J. 348(1985)387
Tassel, J.P., see Montay, G. 339(1985)254
Tatsuhara, T., Tabuchi, F., Umino, M., Okamura, Y., Shigemasa, C., Abe, K. and Mashiba, H.
Determination of thiamazole in serum by HPLC with electrochemical detection 338(1985)xxx
Tatsuno, M., see Tsuruta, K. 338(1985)xxx
Tay, E.T., see Cunningham, R. 344(1985)211
Taylor, A.J., see Anderson, D.J. 350(1985)213
Taylor, I.W., Taylor, T. and Chasseaud, L.F.
Determination of creatine in foreign plasma by GLC 348(1985)xxx
Taylor, M.D., see Andrews, A.T. 348(1985)277
Taylor, T., see Taylor, I.W. 348(1985)xxx
Tebbett, I.R.
Analysis of acetaminophen by HPLC 347(1985)431
Tebbett, I.R., Omole, C.I. and Umesh, B.
Determination of paracetamol, salicylic acid and acetyl salicylic acid in serum by HPLC 329(1985)196
Tenwick, D.A. and Rosser, M.T.
Determination of di- and mono-2-ethylhexyl phthalate in plasma by GC 348(1985)389
Ten Noever de Brauw, M.C., see Van der Greef, J. 348(1985)387
Tercyen, J., see Lemon, J. 340(1985)386

Tong, S.L. and Smith, L.L.
HPLC of carbadox, 224(1983)55

Tong, S.L. and Smith, L.L.
HPLC of linuron and hydroxymetabolites and other phenylurea herbicides, 310(1984)

Tanaka, K., Ozaki, H., Otsuka, K. and Arata, T
[illegible] 2-cyclodextrin as a moving stationary phase, 322(1985)211

Tanaka, K., see Otsuka, K. 322(1985)219, 350(1986)

Terada, H. and Takase, Y.
Studies on the analysis of food additives by HPLC. V. Simultaneous determination of preservatives and saccharin in foods by ion-pair chromatography, 239(1982)

Terada, H. and Takase, Y.
Studies on residual antibacterials in foods. IV. Simultaneous determination of penicillin G, penicillin V and ampicillin in milk by HPLC, 232(1982)219

Termonia, M. and Alaerts, G.
Trace analysis of organic pollutants in air by polymer adsorption and dual-[illegible] capillary GC, 282(1983)

Terrier, M., see Guesnard, G. 341(1985)61

Terrom, G., see Blanchin, J.-P. 326(1985)

Tebbi, Z.L., see Jangi, T.L. 331(1985)

Tetsuma, T., Sumi, M., Tanaka, M. and Ohmi, T.
Reversed-phase ion-exchange chromatography of sodium dialkyldithiocarbamates in the presence of tetraalkylammonium salts 242(1982)

Tett, B.K., Cutler, D.J. and Brown, K.F.
HPLC assay for hydroxychloroquine and metabolites in blood and plasma, using a stationary phase of polystyrene [illegible] and a mobile phase at pH 11, with fluorimetric detection 341(1985)

Thiéry, J.P., see Burke, J.T. 341(1985)

Thieme, R.E., Rochholtz, P.C.N. and [illegible]
Quantitative assay of G-methyl-4-[illegible]-phenylacetic acid (MHPA, a [illegible] in human urine by [illegible] and [illegible] 341(1985)

Thisted, B., see Angelo, H.R. 341(1985)

Thistle, J.L., see Tietz, P.S. 341(1985)

Thorne, J.G., see Buchanan, D.S. 341(1985)

Thomas, D., see Bastide, L. 341(1985)

Thomas, L.C., see Dodo, G.H. 322(1985)

Thomas, L.C., see Kanous, T.L. 322(1985)

Thomas, L.C., see Van Engelen, D.L. [illegible]

Thompson, A.K., see Pringle, T.W. [illegible]

Thompson, E.J., see Cox, P.J. 341(1985)

Thompson, G.A. Jr., see Lucas, L.A. 340(1985)

Thompson, H.C. Jr., see Hoskins, L.C. 341(1985)

Thompson, J.A., Norris, K.J. and Petersen, D.R.
Isolation and analysis of N-oxide [illegible] of urinary amino [illegible] of nicotine 1'-N-oxide formation in man, 341(1985)

Thompson, L., see Kabala, K.K. 341(1985)

Thompson, T.S., see Tong, H.Y. 341(1985)

Thomson, S.K., Tucker, D.J. and Knight, M.H.
Determination of homocysteine in urine, [illegible]

Thomson, W.
[illegible] of [illegible] [illegible] [illegible], 341(1985)

Thompson, [illegible], see [illegible] 341(1985)

Thiele, M., see Maly, P. 341(1985)

Thiry, F., see Hamilton, G. 341(1985)

Thor, P.J., Thom, J.L., Miller, J.C. and [illegible]
Development and validation of a method for measuring the glycine and taurine conjugates of bile acids in bile by HPLC, 341(1985)

Tipnis, V. and Rakhit, A.
[illegible] of [illegible] in [illegible] [illegible] and its active metabolite in urine 341(1985)

Tjaa, T.T., Van der Wiel, J.P., Stikkelman, K.M., Kraaijeveld, A.J.J. and Van Rantwijk, F.
HPLC determination of glucose-1-phosphate and glucose and a procedure for [illegible], 341(1985)

Tocco, D.J., see Musser, D.G. 341(1985)

Todd, K.L., see Abernethy, D.R. 341(1985)

Tonges, V. and Reichert, A.
Reversed-phase LC of biotin-labelled nucleotides. A new class of markers in [illegible], 341(1985)

Tolasetti, G., see Zecca, L. 341(1985)

Tokei, S., see Hoshino, T. 341(1985)

Tokuma, Y., Fujiwara, T. and Noguchi, H.
Determination of nitrendipine in human plasma by capillary column GC-negative-ion chemical-ionization MS 341(1985)

Tokuma, Y., Ohno, T., Tanaka, K. and Tsukatani, H.
GC-MS analysis of biologically active [illegible] and its 2-acetyl [illegible], 341(1985)

Tomas, F., see Barberan, F.A.T. 341(1985)

Tomassini, J.-L., Rue, R., Cousson, P., Aubert, C. and Cano, J.-P.
[illegible] of [illegible] in [illegible] [illegible] and their [illegible], [illegible] in plasma and urine by HPLC, 341(1985)

Tonne, P.P. de see Hermans, J. 341(1985)

Tonnis, B. see Bogen, W. 341(1985)

Tonnis, T. see Tanaka, Y. 341(1985)

Tonnis, W., Kuhnel, W., Peschken, T., Peschken, T. and

Smolková-Keulemansová, E.
Relationships between the chromatographic behaviour and structure of some substituted dibenzo[*b,f*]thiepins and their analogues 329(1985)113
Tomková, H., see Kuchař, M. 333(1985)21
Tomoda, Y., see Taira, H. 339(1985)399
Tonelli, D., see Gagliardi, L. 348(1985)321
Tong, H.Y., Thompson, T.S. and Karasek, F.W.
Quantitative responses of normal alkanes in GC-MS analysis 350(1985)27
Tong, H.Y., see Karasek, F.W. 332(1985)169
Tonkin, A.M., see Harrison, P.M. 339(1985)424, 339(1985)429, 343(1985)349
Torel, J., Cillard, J., Cillard, P. and Vie, M.
Determination of oxytetracycline, sulphamethazine and sulphamethoxypyridazine in feed premixes 330(1985)425
Toriyama, K., see Takasugi, N. 342(1985)349
Torok-Both, G.A., Coutts, R.T., Jamali, F., Pasutto, F.M. and Barrington, K.J.
Sensitive nitrogen-phosphorus capillary GC assay for doxapram in premature infants 344(1985)372
Tóth, G. and Zsadányi, J.
Adsorption chromatographic separation of ^{125}I-labelled cortisol-3-(O-carboxymethyl)oxime tyrosine methyl ester 329(1985)264
Toth, G., see Podhradský, D. 328(1985)402
Toth-Martinez, B.L., see Gál, S. 333(1985)239
Toth-Martinez, B.L., see Kiss, L. 333(1985)244
Tóth-Sarudy, É., see Szederkényi, F. 333(1985)404
Tracz, E. and Leboda, R.
Microscopic investigation of the surface of carbon-silica adsorbents. II. Relationships between the type of information obtainable about the surface and the microscopic techniques for its examination 346(1985)346
Trafford, D.J.H., see Coldwell, R.D. 338(1985)289
Trafford, D.J.H., see Street, J.M. 343(1985)259
Trautmann, K.-H., see Dell, D. 344(1985)125
Treffner, E., see Marzo, A. 343(1985)77, 345(1985)390
Tregear, G., see Scanlon, D. 336(1985)189
Tremblay, L., see Massé, R. 339(1985)11
Trivulzio, S., see Zecca, L. 339(1985)203
Trojánek, A. and Kopanica, M.
Thin-layer polarographic detector for the HPLC detection of thiourea derivatives 328(1985)127
Truskolaski, A., see Bertrand, O. 346(1985)111
Tsai, M.Y., Oliphant, C. and Josephson, M.W.
Identification of metabolites diagnostic for organic acidurias by simultaneous dual-column capillary GC 341(1985)1
Tsao, C.S. and Young, M.
Analysis of ascorbic acid derivatives by HPLC with electrochemical detection 330(1985)408
Tsementzis, S.A., see Elworthy, P.M. 343(1985)109
Tsuchiya, H., Hayashi, T., Tatsumi, M., Fukita, T. and Takagi, N.
Simplified procedure for specific determination of imino acids in human blood plasma by HPLC 339(1985)59
Tsuchiya, K., see Takasugi, N. 342(1985)349
Tsuda, T., see Tanaka, N. 332(1985)57
Tsuji, K. and Jenkins, K.M.
Derivatization of secondary amines with 2-naphthalenesulfonyl chloride for HPLC analysis of spectinomycin 333(1985)365
Tsukatani, H., see Tokumura, A. 343(1985)138
Tsukida, K., Ito, M., Tanaka, T. and Yagi, I.
HPLC and spectroscopic characterization of stereoisomeric retinaloximes. Improvements in resolution and implications of the method 331(1985)265
Tsutsumi, A., see Ageta, T. 343(1985)186
Tucker, D.J., see Thomson, S.B. 338(1985)201
Tucker, F.A., Minty, P.S.B. and McGregor, G.A.
Study of nifedipine photodecomposition in plasma and whole blood using capillary GLC 342(1985)193
Tucker, M.P., see Baker, J.O. 346(1985)93
Turci, S.M. and McDonald, M.J.
Isolation of normal and variant human hemoglobin subunits 343(1985)168
Turgeon, J., see Grech-Bélanger, O. 337(1985)172
Turner, C., see Cullen, M.P. 337(1985)29
Turner, R., see Cooper, E.H. 327(1985)269
Turner, R.L., see McDermott, B.J. 338(1985)335
Turton, J., see Scanlon, D. 336(1985)189

U

Ubbink, J.B., Serfontein, W.J. and De Villiers, L.S.
Stability of pyridoxal-5-phosphate semicarbazone: applications in plasma vitamin B_6 analysis and population surveys of Vitamin B_6 nutritional status 342(1985)277
Ube, S., see Shintani, H. 344(1985)145
Uche Nzekwe, E., see Kabra, P.M. 341(1985)383
Ueda, Y., see Kitada, Y. 347(1985)438
Ueno, Y., see Tanaka, T. 328(1985)271
Uetake, A., see Yoshida, T. 348(1985)425
Ul Zafar, N., see Jung, D. 339(1985)198
Ulmasov, T.N., see Potaman, V.N. 329(1985)33
Unate, M., see Tatsuhara, T. 339(1985)149
Uno, T., see Haginaka, J. 341(1985)115
Unruh, G.E. von, see Von Unruh, G.E. 345(1985)67
Uragg, H., see Lintz, W. 341(1985)65
Urano, C., see Higuchi, S. 341(1985)305
Urayama, T., see Yanagisawa, I. 345(1985)229

Urso, R., see Fong, M.H. 333(1985)191
Utamura, T., see Koizumi, K. 341(1985)31
Utley, D., see Maycock, R. 347(1985)429
Uvarova, N.I., see Pokushalova, T.V. 329(1985)189
Uznański, B., see Stec, W.J. 326(1985)263

V

Vaidya, R.A. and Hester, R.D.
Deconvolution of overlapping chromatograms. II. Analytical solution to the moments of the GEX function 333(1985)152
Vajda, M., see Juvancz, Z. 337(1985)121
Vajta, S., Le Moing, J.P., Rovei, V. and Strolin Benedetti, M.
Determination of low plasma concentrations of 3-methoxy-4-hydroxyphenylethylene glycol using GC-negative-ion chemical ionization MS 343(1985)239
Válka, I., Walterová, D. and Šimánek, V.
Determination of papaverine by capillary isotachophoresis 350(1985)336
Valterová, I., see Vaněk, T. 347(1985)188
Vámos-Vigyázó, L., see Szakács-Dobozi, M. 347(1985)310
Van Baal, J.M., Van Leeuwen, M.B., Schouten, T.J. and De Abreu, R.A.
Sensitive HPLC determination of 6-mercaptopurine, 6-thioguanine, 6-mercaptopurine riboside and 6-thioguanosine in biological fluids 336(1985)422
Van Belle, S.J.P., Schoemaker, T.J., Verwey, S.L., Paalman, A.C.A. and McVie, J.G.
Ion-paired HPLC determination of mitoxantrone in physiological fluids 337(1985)73
Van Belle, S.J.P., see De Smet, M. 345(1985)309
Van Blitterswijk, H., see Jansen, E.H.J.M. 347(1985)375, 347(1985)379
Van de Water, C., see Haagsma, N. 333(1985)256
Van den Berg, A.J.J. and Labadie, R.P.
HPLC separation and quantitative determination of 1,8-dihydroxyanthraquinones in plant cell cultures 329(1985)311
Van den Berg, G.A., Nagel, G.T. and Muskiet, F.A.J.
Mass fragmentographic identification of polyamine metabolites in the urine of normal persons and cancer patients, and its relevance to the use of polyamines as tumour markers 339(1985)223
Van den Bossche, W., see De Croo, F. 329(1985)422, 349(1985)301
Van den Bossche, W., see De Schutter, J.A. 350(1985)135
Van der Greef, J., Tas, A.C., Rijk, M.A.H., Ten Noever de Brauw, M.C., Höhn, M., Meyerhoff, G. and Rapp, U.
Determination of progesterone by LC-MS using a moving-belt interface and isotope dilution 343(1985)397
Van der Hem, G.K., see Koopman, B.J. 337(1985)259
Van der Kleyn, E., see Wuis, E.W. 337(1985)341
Van der Schrieck, P., see Frenz, J. 330(1985)1
Van der Straeten, D., Van Langenhove, H. and Schamp, N.
Comparison between theoretical and experimental sampling efficiencies on Tenax GC 331(1985)207
Van der Vliet, A., see Maris, F.A. 347(1985)75
Van der Vusse, G.J., see Roemen, T.H.M. 344(1985)304
Van der Weken, G., see De Schutter, J.A. 350(1985)135
Van der Wiel, J.P., see Tjioe, T.T. 330(1985)412
Van der Wiele, K., see Dijkgraaf, P.J.M. 329(1985)371
Van der Zee, R. and Welling, G.W.
Detection of Sendai virus protein by reversed-phase HPLC combined with immuno-chromatography 327(1985)377
Van Engelen, D.L., Adams, A.K. and Thomas, L.C.
Long pathlength UV absorbance detection for GC with concurrent fluorescence measurement 331(1985)77
Van Eyk, J., see Cachia, P.J. 343(1985)315
Van Gelder, W.M.J.
Determination of the total C_{27}-steroidal alkaloid composition of *Solanum* species by high resolution GC 331(1985)285
Van Gorp, L., see Marescau, B. 345(1985)215
Van Groeningen, C.J., see Kok, R.M. 343(1985)59
Van Harken, D.R., see Forgue, S.T. 342(1985)221
Van Hook, W.A.
Separation factors for the system 1,2-dibromoethane-1,2-dibromo[$^2H_4$]ethane 328(1985)333
Van Kempen, G.M.J., see Pennings, E.J.M. 341(1985)172
Van Kuijk, F.J.G.M., Handelman, G.J. and Dratz, E.A.
Rapid analysis of the major classes of retinoids by step gradient reversed-phase HPLC using retinal (O-ethyl) oxime derivatives 348(1985)241
Van Langenhove, H., see Van der Straeten, D. 331(1985)207
Van Leeuwen, M.B., see Van Baal, J.M. 336(1985)422
Van Loon, G.R., see Mousa, S. 347(1985)405
Van Rantwijk, F., see Tjioe, T.T. 330(1985)412
Van Rossum, J., see Daenens, P. 342(1985)79

Van Veelen, H., see Dikkeschei, L.D. 345(1985)1
Van Vliet, H.P.M. and Poppe, H. Performance of some cell designs for laser-induced fluorescence detection in open-tubular LC 346(1985)149
Vanderhaeghe, H., see Kibwage, I.O. 330(1985)275, 346(1985)309
Vaněk, T., Valterová, I. and Streinz, L. Preparative HPLC of (2*E*)- and (2*Z*)-farnesol 347(1985)188
Vanhaelen, M., see Duez, P. 329(1985)415
Vanhaelen-Fastré, R., see Duez, P. 329(1985)415
Vardanis, A. Ion-exchange columns in HPLC of proteins; a simple loading technique that improves resolution 350(1985)299
Varley, M.J., see Coldwell, R.D. 338(1985)289
Varney, M.S. and Preston, M.R. Measurement of trace aromatic amines in seawater using HPLC with electrochemical detection 348(1985)265
Varsányi-Riedl, K., see Gazdag, M. 328(1985)279, 347(1985)284
Vassanji, N.A., see Lum, J.T. 338(1985)242
Vaughan, G.T. and Milborrow, B.V. Resoltuion of *RS*-abscisic acid and the separation of abscisic acid metabolites from plant tissue by HPLC 336(1985)221
Vázquez, M.A., see García-Raso, A. 331(1985)406
Vedovelli, A., see Palmerini, C.A. 339(1985)285
Veelen, H. van, see Dikkeschei, L.D. 345(1985)1
Veening, H., see Metz, P.A. 330(1985)307
Végh, Z., see Gazdag, M. 328(1985)279
Ventre, E.R. Determination of the content of 2-chlorobenzenesulphonamide and bis(*p*-chlorophenyl) sulphone in 4-chlorobenzenesulphonamide 333(1985)253
Verebey, K., DePace, A. and Mulé, S.J. Quantitation of *l*-α-acetylmethadol and its metabolites in human serum by capillary GLC and nitrogen detection 343(1985)339
Vereczkey, L., see Rónai-Lukács, S. 338(1985)428
Verhaar, L.A.T., see Dijkgraaf, P.J.M. 329(1985)371
Verhagen, J.C.M., see Pennings, E.J.M. 341(1985)172
Verheggen, T.P.E.M., see Reijenga, J.C. 328(1985)353
Verotta, L., see Orsini, F. 349(1985)69
Verwey, S.L., see Van Belle, S.J.P. 337(1985)73
Verzele, M., Dewaele, C. and Duquet, D. Quality criteria and structure of silica gel column packing material 329(1985)351
Verzele, M., see Nevejans, F. 350(1985)145
Verzella, G., Bagnasco, G. and Mangia, A. Ion-pair HPLC analysis of aspartame and related products 349(1985)83
Verzella, G. and Mangia, A. HPLC analysis of Aspartame 346(1985)417
Vessman, J., see Ahnoff, M. 340(1985)73
Vessman, J., see Gyllenhaal, O. 350(1985)328
Viala, A., see Pok, P.R. 338(1985)171
Vidal, E., Guigues, M., Balansard, G. and Elias, R. Determination of ophthalmic therapeutic metipranolol and its degradation product by reversed-phase HPLC 348(1985)304
Vidal, E., see Balansard, G. 348(1985)299
Vie, M., see Torel, J. 330(1985)425
Vijayalakshmi, M.A., see Krishnan, S. 329(1985)165
Vilde, S., see Murel, A. 347(1985)325
Villiers, L.S. de, see Ubbink, J.B. 342(1985)277
Vimalasiri, P.A.D.T., see Burford, R.P. 329(1985)132
Vimalasiri, P.A.D.T., see Haken, J.K. 349(1985)347
Vincek, W.C., Constanzer, M.L., Hessey, G.A. II and Bayne, W.F. Analytical method for the quantification of famotidine, an H_2-receptor blocker, in plasma and urine 338(1985)438
Vincek, W.C., see Constanzer, M.L. 339(1985)414
Vinçon, G.A., see Demotes-Mainard, F.M. 336(1985)438, 342(1985)234
Vine, J., see Brown, A.S. 341(1985)139
Vine, J.H., see Coombe, R.G. 341(1985)146
Vioque, E., Maza, M.P. and Millán, F. HPLC of fatty acids as their *p*-phenylazophenacyl esters 331(1985)187
Vliet, A. van der, see Maris, F.A. 347(1985)75
Vliet, H.P.M. van, see Van Vliet, H.P.M. 346(1985)149
Vodička, L., see Kříž, J. 347(1985)297
Voelkel, A., see Beger, J. 333(1985)319
Voelkel, A., see Szymanowski, J. 330(1985)61
Vogelzang, N.J., see Kotake, A.N. 337(1985)194
Vogt, W., see Jacob, K. 349(1985)283
Volpe, P., see Castiglioni, M. 349(1985)173
Volz, M., see Swobodnik, W. 339(1985)263
Von Sassen, W., Castro-Parra, M., Musch, E. and Eichelbaum, M. Determination of isoniazid, actylisoniazid, acetylhydrazine and diacetylhydrazine in biological fluids by HPLC 338(1985)113
Von Unruh, G.E. and Paar, W.D. GC assay for oxcarbazepine and its main metabolites in plasma 345(1985)67
Vora, M.M., see Boothe, T. 333(1985)269
Vose, C.W., see Robinson, Y. 338(1985)219
Voss, R.H. and Rapsomatiotis, A. Improved solvent-extraction based procedure for the GC analysis of resin and fatty acids in pulp mill effluents 346(1985)205
Vostrowsky, O., see Brosche, T. 345(1985)219
Voyksner, R.D., Williams, F.P. and Hines, J.W. Analysis of candidate anticancer drugs by thermospray HPLC-MS and by high-resolution MS 347(1985)137
Vree, T.B., Baars, A.M. and Wijnands, W.J.A. HPLC and preliminary pharmacokinetics of enoxacin and its 4-oxo metabolite in

human plasma, urine and saliva 343(1985)449
Vree, T.B., see Wuis, E.W. 337(1985)341
Vries, J.X. de, see De Vries, J.X. 338(1985)325
Vusse, G.J. van der, see Roemen, T.H.M. 344(1985)304

W

Wad, N.
Rapid extraction method for ethosuximide and other antiepileptics in serum for determination by HPLC 338(1985)460
Wada, E., see Mizutani, H. 331(1985)359, 331(1985)371
Wada, H.
Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255
Wada, H., see Ozaki, H. 332(1985)243
Wada, H., see Yamamoto, T. 342(1985)261
Wada, H., see Yamatodani, A. 344(1985)115
Wagner, L.W., see Schallinger, L.E. 342(1985)67
Wagner, R.M.
Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399
Wahbi, A.M., see Malspeis, L. 343(1985)190
Wahl, R., Maasch, H.J. and Geissler, W.
HPTLC analysis of dialysable materials in allergen extracts 329(1985)153
Wahlländer, A., Renner, E. and Karlaganis, G.
HPLC determination of dimethylxanthine metabolites of caffeine in human plasma 338(1985)369
Wainer, I.W., Crabos, M. and Cloix, J.F.
Rapid large-scale isolation of biologically active molecules using reversed-phase "flash" chromatography: initial purification of endogenous Na^+,K^+-ATPase inhibitors from human urine 338(1985)417
Wainwright, M.S., see Smith, R.J. 328(1985)11, 331(1985)389, 334(1985)95
Wajcman, H., see Bosisio, A.B. 330(1985)299
Wakabayashi, H., see Shimada, K. 339(1985)331
Wakai, J., see Haginaka, J. 341(1985)115
Waldmann-Meyer, H.
Structure parameters of molecules and media evaluated by chromatographic partition. I. Controlled-pore glasses 350(1985)1
Walendziak, L. and Jadczak, J.
Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193
Walker, D.D., see Smith, R.L. 343(1985)203
Walle, T., Christ, D.D., Walle, U.K. and Wilson, M.J.
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 341(1985)213
Walle, U.K., see Walle, T. 341(1985)213
Wallisch, E., see Hamilton, G. 341(1985)411
Walmsley, L.M., Wilkinson, P.A., Brodie, R.R. and Chasseaud, L.F.
Determination of naftidrofuryl in human plasma by HPLC with fluorescence detection 338(1985)433
Walseth, F., see Hagen, B. 345(1985)173
Walstad, R.A., see Hagen, B. 345(1985)173
Walterová, D., see Válka, I. 350(1985)336
Walters, R.R., see Anderson, D.J. 331(1985)1
Walton, H.F.
Counter-ion effects in partition chromatography 332(1985)203
Wang, C.-T., see Felix, A.M. 327(1985)359
Wang, J., Bonakdar, M. and Deshmukh, B.K.
Measurement of labetalol by HPLC with electrochemical detection 344(1985)412
Wang, M.-Z., Wu, D.-F. and Yu, Y.-W.
HPLC with electrochemical detection of gossypol in human plasma 343(1985)387
Wang,T. and Sun, Y.
Correlation of retention indices obtained with two temperature programmes 330(1985)167
Wantiez, D., see Gaudry, D. 339(1985)404
Ward, S.A., see Edwards, G. 343(1985)224
Ward, S.A., see Mihaly, G.W. 337(1985)166
Wardencki, W. and Staszewski, R.
Sorption properties of some modified molecular sieves 13X towards thiophene and benzene 329(1985)128
Wardlow, M.L.
Rapid isocratic procedure for the separation of platelet-activating factor from phospholipids in human saliva by HPLC 342(1985)380
Warrell, D.A., see Mihaly, G.W. 337(1985)166
Watanabe, K. and Yoshizawa, I.
Clinical analysis of steroids. XXXI. Assay of oestradiol 17-sulphate 4-hydroxylase activity by HPLC with electrochemical detection 337(1985)114
Watanabe, K., see Ishii, D. 332(1985)3
Watanabe, K., see Takeuchi, T. 331(1985)99
Watanabe, N.
Amperometric detection of reducing carbohydrates in HPLC using an amino-bonded column and acetonitrile-water as the eluent 330(1985)333
Watanabe, T., see Kawada, T. 329(1985)99
Watanabe, T., see Yamatodani, A. 344(1985)115
Water, C. van de, see Haagsma, N. 333(1985)256
Watkinson, J.H., see Lauren, D.R. 348(1985)317
Watson, D.G., see Brooks, C.J.W. 347(1985)455
Watson, E., see Kapur, P.A. 337(1985)160
Watson, I.D.
Clavulanate-potentiated ticarcillin: HPLC assays for clavulanic acid and ticarcillin isomers in serum and urine 337(1985)301
Wattenberg, L.W., see Lam, L.K.T. 328(1985)422

Webb, J.R., see Cooper, E.H. 327(1985)179, 327(1985)269
Weber, A.F., Lee, D.W., Opheim, K. and Smith, A.L.
Quantitation of iothalamate in serum and urine by HPLC 337(1985)434
Weber, D.J., see Shah, J.A. 344(1985)41
Webster, S.G.
Catalysed derivatisation of trimethylsilyl ethers of ecdysterone. A preliminary study 333(1985)186
Webster, V.J., see So, N. 337(1985)81
Wechsler, J.G., see Swobodnik, W. 339(1985)263
Wee, E.L., see Norman, E.J. 337(1985)21
Wehinger, G., see Battista, H.J. 345(1985)77
Wehr, C.T., see Cunico, R.L. 336(1985)105
Weidenhoffer, Z., Plzák, Z. and Dolanský, J.
Trace analysis of organic compounds in volatile inorganic halides by the dynamic headspace technique and GC-MS 350(1985)324
Weidolf, L.
Bimodal column switching LC assay of six metabolites of [^{14}C]felodipine in rat urine 343(1985)85
Weir, D.E., see Minkler, P.E. 336(1985)271
Weissman, J.B., see DiBella, E.E. 328(1985)101
Weisweiler, P., see Schwandt, P. 345(1985)145
Weken, G. van der, see De Schutter, J.A. 350(1985)135
Welinder, B.S., Linde, S. and Hansen, B.
Reversed-phase HPLC of insulin and insulin derivatives. A comparative study 348(1985)347
Welling, G.W., Groen, G., Slopsema, K. and Welling-Wester, S.
Combined size-exclusion and reversed-phase HPLC of a detergent extract of Sendai virus 326(1985)173
Welling, G.W., see Van der Zee, R. 327(1985)377
Welling-Wester, S., see Welling, G.W. 326(1985)173
Wells, D.E., Cowan, A.A. and Christie, A.E.G.
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372
Wells, G.
Theory of anion contributions to non-linear electron-capture detector response 346(1985)1
Wells, P.G., see Lum, J.T. 338(1985)242
Wells, R.D.
HPLC of DNA (Review) 336(1985)3
Welscher, T.M., see Chang, S.F. 343(1985)119
Wenclawiak, B., Barkley, R.M., Williams, E.J. and Sievers, R.E.
GC and LC separation of geometrical isomers of tris(2,2,7-trimethyloctane-3,5-dionato)chromium(III) and cobalt(III) chelates and other Cr-β-diketonates 349(1985)469
Wendt, G., see Dell, D. 344(1985)125
Wenzel, T.J., see DiBella, E.E. 328(1985)101
Werner, B., see Del Principe, D. 342(1985)285
Westerlund, D., see Carlqvist, J. 344(1985)285
Westhead, D., see Elworthy, P.M. 343(1985)109
Wettenhall, R.E.H. and Quinn, M.J.
Separation of tryptic phosphopeptides of ribosomal origin by reversed-phase HPLC 336(1985)51
Wheals, B.B., White, P.C. and Paterson, M.D.
HPLC method utilising single or multi-wavelength detection for the comparison of disperse dyes extracted from polyester fibres 350(1985)205
Wheeler, W.B., see Bardalaye, P.C. 330(1985)403
Whelpton, R. and Moore, T.
Sensitive LC method for physostigmine in biological fluids using dual-electrode electrochemical detection 341(1985)361
Whist, J.E., see Svendsen, J.S. 337(1985)9
Whitaker, J.N., see Heinemann, M.A. 329(1985)295
White, C.J., see Morreal, C.E. 345(1985)380
White, P.C., see Bedford, K.R. 347(1985)398
White, P.C., see Finden, D.A.S. 342(1985)179
White, P.C., see Wheals, B.B. 350(1985)205
Whiton, R.S., Lau, P., Morgan, S.L., Gilbart, J. and Fox, A.
Modifications in the alditol acetate method for analysis of muramic acid and other neutral and amino sugars by capillary GC-MS with selected ion monitoring 347(1985)109
Widholm, J.M., see Brotherton, J.E. 350(1985)332
Wiechert, P., see Marescau, B. 345(1985)215
Wiedemann, K., see Schulz, A. 328(1985)417
Wiehle, R.D., see Boyle, D.M. 327(1985)369
Wiehle, R.D., see Hyder, S.M. 327(1985)237
Wiel, J.P. van der, see Tjioe, T.T. 330(1985)412
Wieland, H., see McConathy, W.J. 342(1985)47
Wiele, K. van der, see Dijkgraaf, P.J.M. 329(1985)371
Wigilius, B., see Borén, H. 348(1985)67
Wijnands, W.J.A., see Vree, T.B. 343(1985)449
Wildfeuer, A., see Räder, K. 344(1985)416
Wilke, C.R., see Arnold, F.H. 330(1985)159
Wilkinson, P.A., see Walmsley, L.M. 338(1985)433
Wiłkomirski, B.
TLC and GLC separation of trihydroxy pentacyclic triterpene alcohols from *Calendula officinalis* flowers 330(1985)177
Willems, J.L., see De Schryver, E.P. 338(1985)389
Willemse, P.H.B., see Dikkeschei, L.D. 345(1985)1
Williams, A.T. Rhys, see Rhys Williams, A.T. 341(1985)198
Williams, C.H., Dozier, S.E., Buzello, W., Gehrke, C.W., Wong, J.K. and Gerhardt, K.O.
Plasma levels of norepinephrine and epinephrine during malignant hyperthermia in susceptible pigs 344(1985)71
Williams, E.J., see Wenclawiak, B. 349(1985)469
Williams, F.P., see Voyksner, R.D. 347(1985)137

Willmott, N., see Cummings, J. 343(1985)208
Wilman, D.E.V., see McDermott, B.J. 338(1985)335
Wils, E.R.J. and Hulst, A.G.
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379
Wilson, D.C., see Maguire, J.H. 342(1985)323
Wilson, K., see Colthup, P.V. 345(1985)111
Wilson, M.F.
Rapid method for the separation and quantification of simple phenolic acids in plant material using HPLC 346(1985)440
Wilson, M.J., see Walle, T. 341(1985)213
Winkler, G., Heinz, F.X., Guirakhoo, F. and Kunz, C.
Separation of flavivirus membrane and capsid proteins by multistep HPLC optimized by immunological monitoring 326(1985)113
Winkler, G., Wolschann, P., Briza, P., Heinz, F.X. and Kunz, C.
Spectral properties of trifluoroacetic acid-acetonitrile gradient systems for separation of picomole quantities of peptides by reversed-phase HPLC 347(1985)83
Wittliff, J.L., see Boyle, D.M. 327(1985)369
Wittliff, J.L., see Hyder, S.M. 327(1985)237
Wohlrab, H., see Kolbe, H.V.J. 327(1985)1
Wolf, R.M., see Francotte, E. 347(1985)25
Wolff, G.A., see Gill, J.P. 350(1985)37
Wolschann, P., see Winkler, G. 347(1985)83
Wolthers, B.G., see Dikkeschei, L.D. 345(1985)1
Wolthers, B.G., see Koopman, B.J. 337(1985)259
Wong, D.L., see Speedie, M.K. 327(1985)351
Wong, J.K., see Williams, C.H. 344(1985)71
Wong, L.T., Xu, Z.J. and Hsia, C.J.C.
Simple purification procedure of rat α-fetoprotein by a combination of Cibacron Blue gel affinity chromatography and anion-exchange HPLC 338(1985)410
Wong, L.T., Xu, Z.J. and Hsia, C.J.C.
Fractionation of rat α-fetoprotein by HPLC 341(1985)452
Woodward, A.J., see Lewis, P.A. 327(1985)261
Wright, G.J., see Osman, M.A. 336(1985)329, 345(1985)430
Wu, B.-L., see Hjertén, S. 341(1985)295
Wu, D.-F., see Wang, M.-Z. 343(1985)387
Wu, H.-L., see Funazo, K. 346(1985)215
Wu, Q., see Chang, C.A. 347(1985)51
Wuis, E.W., Dirks, R.J.M., Vree, T.B. and Van der Kleyn, E.
HPLC analysis of baclofen in plasma and urine of man after precolumn extraction and derivatization with *o*-phthaldialdehyde 337(1985)341

X

Xu, Z.J., see Wong, L.T. 338(1985)410, 341(1985)452

Y

Yachetti, S.D., see Leelavathi, D.E. 339(1985)105
Yagen, B., Bialer, M. and Sintov, A.
GC assay with pharmacokinetic applications for monitoring T-2 and HT-2 toxins in plasma 343(1985)67
Yagi, I., see Tsukida, K. 331(1985)265
Yamada, J.-I., see Yamamura, K. 331(1985)383
Yamada, K., Yamato, C. and Sato, T.
Improved assay procedure for tripamide in blood 339(1985)210
Yamada, K., see Niwa, T. 336(1985)345, 337(1985)1
Yamagami, T., see Okamoto, T. 342(1985)35
Yamaguchi, K., see Ohta, K. 350(1985)292
Yamaguchi, K., see Parker, C.E. 344(1985)378
Yamaguchi, M., Hara, S., Matsunaga, R., Nakamura, M. and Ohkura, Y.
3-Bromomethyl-6,7-dimethoxy-1-methyl-2(1*H*)-quinoxalinone as a new fluorescence derivatization reagent for carboxylic acids in HPLC 346(1985)227
Yamaguchi, M., see Hara, S. 344(1985)33
Yamaguchi, M., see Iwata, T. 344(1985)351
Yamaguchi, Y., see Seki, T. 332(1985)9, 332(1985)283
Yamamoto, H., see Nakano, K. 332(1985)127, 343(1985)21
Yamamoto, K., see Kato, S. 333(1985)175
Yamamoto, K., see Mizutani, T. 333(1985)171
Yamamoto, T., Yamatodani, A., Nishimura, M. and Wada, H.
Determination of dopamine-3- and -4-O-sulphate in human plasma and urine by anion-exchange HPLC with fluorimetric detection 342(1985)261
Yamamoto, Y., see Mikasa, H. 343(1985)179
Yamamura, K., Yamada, J.-I. and Yotsuyanagi, T.
HPLC assay of anti-inflammatory drugs incorporated in gel ointments. Separation and stability testing 331(1985)383
Yamanaka, N., see Maeda, K. 345(1985)11
Yamanaka, N., see Shiraishi, S. 338(1985)51
Yamane, M., see Yanagisawa, I. 345(1985)229
Yamane, T., see Furuta, H. 337(1985)103
Yamato, C., see Yamada, K. 339(1985)210

Yamatodani, A., Fukuda, H., Wada, H., Iwaeda, T. and Watanabe, T.
HPLC determination of plasma and brain histamine without previous purification of biological samples: cation-exchange chromatography coupled with post-column derivatization fluorometry 344(1985)115
Yamatodani, A., see Yamamoto, T. 342(1985)261
Yanagihara, Y., see Hanai, T. 349(1985)323
Yanagihara, Y., see Seki, T. 332(1985)9, 332(1985)283
Yanagihara, Y., see Yasukawa, K. 332(1985)287
Yanagisawa, I., Yamane, M. and Urayama, T.
Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 345(1985)229
Yang, C.-Y. and Sepulveda, F.I.
Separation of phenylthiocarbamyl amino acids by HPLC on Spherisorb octadecylsilane columns 346(1985)413
Yasaka, T., see Nakano, K. 332(1985)127, 343(1985)21
Yasuda, H., see Haginaka, J. 341(1985)115
Yasuhara, A., Morita, M. and Fuwa, K.
Temperature-programmed retention indices of 221 halogenated organic compounds with 1-bromoalkanes as references 328(1985)35
Yasukawa, K., Kasai, M., Yanagihara, Y. and Noguchi, K.
HPLC analysis of peptides on an Asahipak GS-320 column packed with hydrophilic polymer gel 332(1985)287
Yata, J., see Suzuki, H. 341(1985)176
Yean, W.Q., see Lewis, N.G. 331(1985)419
Yee, C., see Lam, L.K.T. 328(1985)422
Yee, G.C., see Gmur, D.J. 344(1985)422
Yeh, K.C., see Musson, D.G. 337(1985)363
Yi, C., Fasching, J.L. and Brown, P.R.
Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 339(1985)75
Yinon, J. and Hwang, D.-G.
Metabolic studies of explosives. II. HPLC-MS of metabolites of 2,4,6-trinitrotoluene 339(1985)127
Yip, H., see Coombe, R.G. 341(1985)146
Yip, K.F. and Albarella, J.P.
Use of HPLC in the preparation of flavin adenine dinucleotide analyte conjugates 326(1985)301
Yokokawa, S., see Takasugi, N. 342(1985)349
Yokosawa, N., see Taniguchi, N. 339(1985)394
Yoneda, M., see Oyama, H. 338(1985)71
Yonezawa, K., Sato, K. and Kobayashi, A.
HPLC of a new β-blocker, 4-[3-(*tert.*-butylamino)-2-hydroxypropoxyl-N-methylisocarbostyril hydrochloride, in plasma using fluorometric detection 339(1985)219
Yoo, B., see Lee, M. 348(1985)407
Yoshida, A., Ichihashi, Y. and Yoshioka, M.
Determination of homovanillic acid in human plasma by HPLC with electrochemical detection 343(1985)155
Yoshida, T., Uetake, A., Murayama, H., Nimura, N. and Kinoshita, T.
Fluorescent labelling of amino acids with 9-anthryldiazomethane and its applications to HPLC 348(1985)425
Yoshihira, K., see Inoue, T. 346(1985)450
Yoshioka, M., Tamura, Z., Senda, M. and Miyazaki, T.
Analyser of adenine nucleotides 344(1985)345
Yoshioka, M., see Yoshida, A. 343(1985)155
Yoshizawa, I. and Kameyama, M.
Clinical analysis of steroids. XXXII. New method for the determination of hapten number of the antigen prepared for the radioimmunoassay of steroidal sulphates using ion chromatography 338(1985)404
Yoshizawa, I., see Watanabe, K. 337(1985)114
Yost, R.L. and Derendorf, H.
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131
Yotsuyanagi, T., see Yamamura, K. 331(1985)383
Young, J.D., see Schilcher, R.B. 337(1985)55
Young, M., see Tsao, C.S. 330(1985)408
Young, N.M. and Jackson, G.E.D.
Anomalous behavior of lectins in size-exclusion HPLC and gel electrophoresis 336(1985)397
Yousefnejad, D., see Cone, E.J. 337(1985)291
Yoza, N., see Baba, Y. 348(1985)27, 350(1985)119, 350(1985)461
Yu, Y.-W., see Wang, M.-Z. 343(1985)387
Yuen, S. and Lau-Cam, C.A.
TLC screening procedure for undeclared synthetic drugs in Chinese herbal preparations 329(1985)107
Yuhki, Y., see Tadano, K. 341(1985)228
Yuki, H., see Okamoto, Y. 350(1985)127

Z

Zafar, N.U., see Jung, D. 339(1985)198
Zambonin, P.G., see Palmisano, F. 344(1985)249
Zamudio, F., see Ladrón de Guevara, O. 349(1985)91
Zecca, L. and Ferrario, P.
Determination of neurotransmitter amino acids by HPLC with fluorescence detection 337(1985)391
Zecca, L., Trivulzio, S., Pinelli, A., Colombo, R. and Tofanetti, O.
Determination of glibenclamide, chlorpropamide and tolbutamide in plasma by HPLC with UV detection 339(1985)203
Zee, R. van der, see Van der Zee, R. 327(1985)377

Zeeuw, R.A. de, see Bogusz, M. 342(1985)241
Zeeuw, R.A. de, see Feitsma, K.G. 333(1985)59
Zenkevich, I.G., see Ioffe, B.V. 349(1985)385
Zhu, A.
Calculation of retention indices in temperature-programmed capillary GC 331(1985)229
Zhu, M.-D., see Hjertén, S. 327(1985)157, 346(1985)265
Zhukova, N.V., see Svetashev, V.I. 330(1985)396
Zielke, H.R.
Determination of amino acids in the brain by HPLC with isocratic elution and electrochemical detection 347(1985)320
Zierman, L., see Gleispach, H. 344(1985)11
Zimmerman, E.F., see Norman, E.J. 337(1985)21
Zimmerman, R., see De Vries, J.X. 338(1985)325
Zimmermann, B., see Kuhnz, W. 344(1985)309
Zipser, R.D., Morrison, A., Laffi, G. and Duke, R.
Assay methods for 6-keto-prostaglandin $F_{1\alpha}$ in human urine. Comparison of chromatographic techniques with radioimmunoassay and GC-negative-ion chemical-ionization MS 339(1985)1
Zoccolillo, L., see Ronchetti, M. 348(1985)159
Zon, G., see Stec, W.J. 326(1985)263
Zonta, F. and Stancher, B.
Quantitative analysis of phylloquinone (Vitamin K_1) in soy bean oils by HPLC 329(1985)257
Zoontjes, P.W., see Jansen, E.H.J.M. 347(1985)375, 347(1985)379
Zsadányi, J., see Tóth, G. 329(1985)264
Zschunke, M., see Brendel, E. 339(1985)359
Zuder, G.F.X., see Gamerith, G. 328(1985)241
Żukowski, J., see Debowski, J. 329(1985)206
Zullino, G., see Rosini, C. 348(1985)79

SUBJECT INDEX

VOLUMES 326–350

A

Abbé refractometer
The Abbé refractometer as a LC detector using an image sensor for boundary location 332(1985)195

Abscisic acid
Simultaneous determination of abscisic acid and jasmonic acid in plant extracts using HPLC 330(1985)347

Resoltuion of *RS*-abscisic acid and the separation of abscisic acid metabolites from plant tissue by HPLC 336(1985)221

7-Acetamidonitrazepam
Determination of nitrazepam and its main metabolites in urine by TLC and direct densitometry 339(1985)163

Acetaminophenols
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Acetone
Adsorption of mixtures of acetone and benzene vapors on graphitized carbon black 328(1985)63

The re-injection method for recovery of components adsorbed during syringe gas sampling: a statistical evaluation 329(1985)119

Gaseous acidity and basicity scales as guides to chemical ionization in reversed-phase LC-MS with direct liquid introduction 347(1985)61

Acetonitrile
Spectral properties of trifluoroacetic acid-acetonitrile gradient systems for separation of picomole quantities of peptides by reversed-phase HPLC 347(1985)83

N-Acetylamino acids
Urinary excretion of N-acetylamino acids 338(1985)187

2-Acetylaminofluorene
Analysis of 2-acetylaminofluorene and its metabolites by GC with electron-capture detection 350(1985)307

Acetylator phenotype
Ion-pair HPLC determination of isoniazid and acetylisoniazid in plasma and urine. Application for acetylator phenotyping 341(1985)193

Acetylcholine
Improved analysis of acetylcholine and choline in canine brain and blood samples by capillary GC-MS 339(1985)170

Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 341(1985)23

Highly sensitive assay for acetylcholinesterase activity by HPLC with electrochemical detection 344(1985)93

Acetylcholinesterase
Highly sensitive assay for acetylcholinesterase activity by HPLC with electrochemical detection 344(1985)93

N-Acetylcysteine
Improved method for the determination of N-acetylcysteine in human plasma by HPLC 327(1985)261

Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193

6-Acetyldihydrohomopterin
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

6-Acetyl-7,8-dihydropterin
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

Acetylhydrazine
Determination of isoniazid, actylisoniazid, acetylhydrazine and diacetylhydrazine in biological fluids by HPLC 338(1985)113

Acetylisoniazid
Determination of isoniazid, actylisoniazid, acetylhydrazine and diacetylhydrazine in biological fluids by HPLC 338(1985)113

Ion-pair HPLC determination of isoniazid and acetylisoniazid in plasma and urine. Application for acetylator phenotyping 341(1985)193

***l*-α-Acetylmethadol**
Quantitation of *l*-α-acetylmethadol and its metabolites in human serum by capillary GLC and nitrogen detection 343(1985)339

***sn*-2-Acetyl phospholipids**
GC-MS analysis of biologically active phospholipids having and *sn*-2-acetyl group 343(1985)138

Acetylsalicylic acid
Determination of paracetamol, salicylic acid and acetyl salicylic acid in serum by HPLC 329(1985)196

HPLC of antipyretics on chemically modified porous glass 332(1985)265

Analysis of antipyretics by semimicro LC 332(1985)269

1-(D-3-Acetylthio-2-methylpropanoyl)-L-propyl-L-phenylalanine
Determination of captopril and its mixed disulphides in plasma and urine by HPLC 338(1985)161

Acetyltrimoprostil
Determination of acetyltrimoprostil and its metabolite trimoprostil in human or dog plasma by GC-negative-ion chemical ionization MS 339(1985)313

N-Acetyl-DL-tryptophan
Chromatographic determination of N-acetyl-DL-tryptophan and octanoic acid in human albumin solutions 333(1985)381

N-Acetyl-L-valine-*tert.*-butylamide
Chiral recognition mechanisms in the enantioselectivity of chiral hydrogen bonding additives in LSC 349(1985)143

α_1-Acid glycoprotein
Two-step purification of human α_1-acid glycoprotein 341(1985)457

Acidities
Gaseous acidity and basicity scales as guides to chemical ionization in reversed-phase LC-MS with direct liquid introduction 347(1985)61

Acid protease
Purification of an acid protease and a serine carboxypeptidase from *Aspergillus niger* using metal-chelate affinity chromatography 329(1985)165

Acrinol
Application of thin-layer stick chromatographic identification test methods to drugs contained in preparations in the Pharmacopoeia of Japan 333(1985)115

Acrylamide
Direct determination of acrylamide in tissue culture solution by LC using column switching 346(1985)376

Acrylates
Chromatography of monomers. III. TLC separation of C_6-C_{18} straight-chain acrylates from corresponding methacrylates and alcohols 330(1985)439

Actinobacillus
Differentiation between major species of the *Actinobacillus-Haemophilis-Pasteurella* group by GC of trifluoroacetic acid anhydride derivatives from whole-cell methanolysates 342(1985)13

***Actinomadura* R39**
Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary GC-MS 341(1985)139

Activity coefficients
Adsorption from multi-component liquid mixtures on to heterogeneous solid surfaces: the role of adsorption of water on silica in liquid-solid adsorption chromatography 348(1985)1

Utilization of the functional group contribution concept in LC on chemically bonded reversed phases 348(1985)49

Measurement of activity coefficients by reversed-flow GC 349(1985)369

Possibility and consequences of using different concentration scales in the study of solution thermodynamics by GLC 350(1985)339

Acyclovir
HPLC determination of acyclovir in serum 343(1985)203

Adenine nucleotides
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides 337(1985)201

Rapid automated HPLC analysis of cyclic adenosine 3',5'-monophosphate. Synthesis in brain tissues 341(1985)43

Analyser of adenine nucleotides 344(1985)345

Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column LC under isocratic conditions 345(1985)267

Simple separation of adenosine-5'-phosphosulphate, adenosine-5'-phosphoramidate and cyclic adenosine-5'-monophosphate nucleotides involved in adenosine-5'-phosphosulphate degradation 348(1985)296

Adenosine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 345(1985)150

Adenosine 5'-monophosphate
Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255

Adenosine 3':5'-monophosphate
Rapid automated HPLC analysis of cyclic adenosine 3',5'-monophosphate. Synthesis in brain tissues 341(1985)43

Adenosine 5'-monophosphate
Simple separation of adenosine-5'-phosphosulphate, adenosine-5'-phosphoramidate and cyclic adenosine-5'-monophosphate nucleotides involved in adenosine-5'-phosphosulphate degradation 348(1985)296

Adenosine 3':5'-monophosphate
Simple separation of adenosine-5'-phosphosulphate, adenosine-5'-phosphoramidate and cyclic adenosine-5'-monophosphate nucleotides involved in adenosine-5'-phosphosulphate degradation 348(1985)296

Adenosine 3':5'-monophosphate derivatives
Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. II. Quantitative determination of hydrophobicity 350(1985)105

Adenosine 5'-phosphosulphate
Simple separation of adenosine-5'-phosphosulphate, adenosine-5'-phosphoramidate and cyclic adenosine-5'-monophosphate nucleotides involved in adenosine-5'-phosphosulphate degradation 348(1985)296

S-Adenosylethionine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 345(1985)150

S-Adenosylhomocysteine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 345(1985)150

S-Adenosylmethionine
Determination of adenosine and S-adenosyl derivatives of sulfur amino acids in rat liver by HPLC 345(1985)150

Adrenaline
Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Direct clean-up and analysis of urinary catecholamines 336(1985)374

Simple and fast analysis of adrenaline and noradrenaline in plasma on microbore HPLC columns using fluorimetric detection 337(1985)384

Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Estimation of adrenal catecholamines by elevated-temperature LC with amperometric detection 342(1985)363

HPLC assay of free norepinephrine, epinephrine, dopamine, vanillylmandelic acid and homovanillic acid 343(1985)9

Determination of intraoperative plasma catecholamine concentrations using LC with electrochemical detection 343(1985)285

Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61

Plasma levels of norepinephrine and epinephrine during malignant hyperthermia in susceptible pigs 344(1985)71

Rapid and reliable estimation of urinary free catecholamines in patients with pheochromocytoma: comparison with plasma catecholamines and vanillylmandelic acid excretion 344(1985)313

Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

Determination of phenylethanolamine N-methyltransferase by HPLC with fluorescence detection 348(1985)407

β-Adrenoceptor antagonists
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Adriamycin
New HPLC assay for plasma doxorubicin 337(1985)194

Adriamycin-loaded albumin microspheres: qualitative assessment of drug incorporation and in vitro release by HPLC and high-speed multi-diode array spectrophotometric detection 343(1985)208

Adsorbents
Adsorption of mixtures of acetone and benzene vapors on graphitized carbon black 328(1985)63

Correlation between chemical and physical characteristics and adsorption properties of some styrene copolymers 330(1985)55

Use of bonded-phase silica sorbents for rapid sampling of impurities in illicit amphetamine for HPLC analyses 331(1985)339

Application of interferometry in studies of transport processes near ionites and adsorbents 333(1985)198

Microscopic investigation of the surface of carbon-silica adsorbents. II. Relationships between the type of information obtainable about the surface and the microscopic techniques for its examination 346(1985)346

Pre-concentration of organic pollutants. Potential interference from the use of styrene copolymer adsorbents 349(1985)181

Adsorption
Sorption properties of some modified molecular sieves 13X towards thiophene and benzene 329(1985)128

Adsorption effects in GLC. Solute retention in the hydrocarbon solute-polar stationary phase (polypropylene glycol 2000) system 330(1985)193

Elution and washing in fixed bed affinity chromatography 330(1985)356

Adsorption from multi-component liquid mixtures on to heterogeneous solid surfaces: the role of adsorption of water on silica in liquid-solid adsorption chromatography 348(1985)1

Adsorption isotherms
Adsorption of mixtures of acetone and benzene vapors on graphitized carbon black 328(1985)63

Affinity adsorbents
High-performance liquid affinity chromatography of β-lactamase 326(1985)163

High-performance affinity chromatography of trypsins on Asahipak GS-gel coupled with *p*-aminobenzamidine 333(1985)107

Comparative study of affinity chromatography of components of the hyaluronate-proteoglycan complex to immobilized hyaluronate 350(1985)227

Affinity chromatography-filtration
Combined recycling affinity column-filtration technique 346(1985)247

Affinity partitioning
Affinity partitioning of biopolymers and membranes in Ficoll-dextran aqueous two-phase systems 331(1985)11

Aflatoxins
TLC separations using a temperature gradient 333(1985)69

TLC of mycotoxins (Review) 334(1985)211

Agarose
Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33

Column lifetime of a new agarose medium for high-performance gel filtration chromatography at basic pH 330(1985)360

Air pollutants
Trace analysis of organic pollutants in air by polymer adsorption and dual flame ionization detection-electron-capture detection capillary GC 328(1985)367

Precolumn sample enrichment device for analysis of ambient volatile organics by GC-MS 329(1985)399

Albumin
Purification of human serum gamma globulins by hydrophobic interaction HPLC 326(1985)235

Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

HPLC studies on non-mercapt mercapt conversion of human serum albumin II 332(1985)19

Ion-exchange HPLC studies on sulphydryl-catalysed structural alterations of bovine mercaptalbumin 332(1985)29

Chromatographic determination of N-acetyl-DL-tryptophan and octanoic acid in human albumin solutions 333(1985)381

Isotachophoresis as a useful tool for monitoring neurological complications of acute leukaemia in children 342(1985)285

Albumin microspheres
Adriamycin-loaded albumin microspheres: qualitative assessment of drug incorporation and in vitro release by HPLC and high-speed multi-diode array spectrophotometric detection 343(1985)208

Albuterol
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Alcoholic strength
Determination of alcoholic strength in alcoholic beverages by GLC. Precision and accuracy 333(1985)389

Alcohols
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. II. Influence of organic co-solvents on retention at 25°C 333(1985)41

HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Aldaric acids
High-speed LC of sugar acids on anion-exchange resins 329(1985)371

Aldehyde dimethyl acetals
Methyl enol ethers as artefacts in capillary GC profiles of aldehyde dimethyl acetals 345(1985)219

Aldehydes
Determination of low concentrations of low-molecularweight aldehydes and ketones in aqueous samples 329(1985)81

Determination of aldehydes and ketones by derivatization and LC-MS 333(1985)337

Alditol acetates
Modifications in the alditol acetate method for analysis of muramic acid and other neutral and amino sugars by capillary GC-MS with selected ion monitoring 347(1985)109

Capillary GC of partially methylated alditol acetates on a SP-2100 wall-coated open-tubular column 350(1985)304

Aldonic acids
High-speed LC of sugar acids on anion-exchange resins 329(1985)371

Alfentanil
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Aliphatic acids
HPLC of microbial acid metabolites 336(1985)125

Alkaloids
Separation of carbazole alkaloids by GLC 329(1985)178

Comparison between TLC-densitometry and HPLC for the determination of hyoscyamine and hyoscyne in leaves, fruits and seeds of *Datura* (*Datura* spp.) 329(1985)415

Retention indices of some ergopeptines in reversed-phase HPLC 331(1985)245

Determination of the total C_{27}-steroidal alkaloid composition of *Solanum* species by high resolution GC 331(1985)285

HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Application of thin-layer stick chromatographic identification test methods to drugs contained in preparations in the Pharmacopoeia of Japan 333(1985)115

Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

HPLC assay for pilocarpine in aqueous humor 336(1985)337

Drug level monitoring: cytostatics (Review) 340(1985)31

Sub-nanogram analysis of yohimbine and related compounds by HPLC 342(1985)333

Capillary GC determination of vincamine in plasma 344(1985)367

HPLC determination of vinca-alkaloids in plasma and urine 345(1985)309

HPLC system for the separation of ergot alkaloids with applicability to the analysis of illicit lysergide (LSD) 346(1985)423

Cinchona alkaloids for preparing new, easily accessible chiral stationary phases. II. Resolution of binaphthol derivatives on silica-supported quinine 348(1985)79

Determination of pyrrolizidine alkaloids in *Senecio inaequidens* D.C. by capillary GC 349(1985)23

Determination of papaverine by capillary isotachophoresis 350(1985)336

Alkanes
Group separation of oil residues by supercritical fluid chromatography 349(1985)439

Quantitative responses of normal alkanes in GC-MS analysis 350(1985)27

***n*-Alkyl acrylates**
Chromatography of monomers. IV. GLC studies of C_1-C_6 *n*-alkyl and C_3-C_6 isoalkyl acrylates and their hydrogen halide and halogen addition derivatives 348(1985)141

Alkylbenzenes
Quantitation of long-chain alkylbenzenes in environmental samples by silica gel CC and high-resolution GC 346(1985)281

Alkyl benzoates
GLC analyses. XLII. Retention behaviour of C_1-C_{12} *n*-alkyl esters of benzoic and pentafluorobenzoic acids on SE-30 and OV-351 capillary columns 328(1985)325

Alkyl boronates
Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370

Alkyl disulphides
?In situ *generation of alkyl sulphides and alkyl disulphides* for GC identification 330(1985)131

Alkyl ether lysophospholipids
Reversed-phase HPLC separation of molecular species of alkyl ether, vinyl ether, and monoacyl lyosphospholipids 338(1985)61

Alkyl grafts
Long-chain alkyl grafts and mixed alkyl-alkane layers at the surface of macroporous silicas. Their GC properties below and above the phase transition 329(1985)331

Alkyl halides
Derivatization of alkyl halides and epoxides with picric acid salts for improved HPLC detection 333(1985)349

S-Alkylisothiourea
?In situ *generation of alkyl sulphides and alkyl disulphides* for GC identification 330(1985)131

Alkyl pentafluorobenzoates
GLC analyses. XLII. Retention behaviour of C_1-C_{12} *n*-alkyl esters of benzoic and pentafluorobenzoic acids on SE-30 and OV-351 capillary columns 328(1985)325

Alkylphosphines
Quantitative GC analysis of low-molecular-weight alkylphosphines in the presence of phosphine and hydrocarbons 330(1985)400

Alkyl sulphides
?In situ *generation of alkyl sulphides and alkyl disulphides* for GC identification 330(1985)131

Allergens
HPTLC analysis of dialysable materials in allergen extracts 329(1985)153

Alminoprofen
Rapid determination of alminoprofen in plasma by HPLC 343(1985)455

Aloe-emodin
HPLC separation and quantitative determination of 1,8-dihydroxyanthraquinones in plant cell cultures 329(1985)311

Alprenolol
Direct derivatization of alprenolol and its 4-hydroxy metabolite in urine with phosgene and methanol prior to analysis by capillary column GC 349(1985)447

Althesin
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Althiazide
Simultaneous determination of althiazide and spironolactone in tablets by HPLC 329(1985)422

Ametantrone
Ion-paired HPLC determination of mitoxantrone in physiological fluids 337(1985)73

Amikacin
Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Amines
Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109

Trace analysis of amines and isocyanates using glass capillary GC and selective detection. IV. Determination of free aromatic amines using nitrogen-selective detection 331(1985)321

Derivatization of secondary amines with 2-naphthalenesulfonyl chloride for HPLC analysis of spectinomycin 333(1985)365

GC-MS analysis of volatile amines produced by several strains of *Clostridium* 337(1985)213

Frequency-pulsed electron-capture GLC studies of chemical changes in sera of patients with schistosomiasis 339(1985)243

Isolation and analysis of N-oxide metabolites of tertiary amines: quantitation of nicotine-1'-N-oxide formation in mice 341(1985)349

Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370

Quantum chemical parameters in correlation analysis of GLC retention indices of amines 346(1985)53

Capillary GC method for the determination of complex mixture of isocyanates and amines 346(1985)191

Analytical separation of enantiomeric polygodials by GC of pyrrole derivatives 347(1985)455

Measurement of trace aromatic amines in seawater using HPLC with electrochemical detection 348(1985)265

L-Amino acid decarboxylase
Assay of aromatic L-amino acid decarboxylase by HPLC with radiochemical detection 342(1985)175

D-Amino acid oxidase apoenzyme
Purification of D-amino acid oxidase apoenzyme by affinity chromatography on Cibacron Blue Sepharose 347(1985)316

Amino acids
Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399

Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

Adsorption of SH-containing amino acids, peptides and proteins on the 2-nitrophenylhydrazonopropanedinitrile derivative of Spheron Ara 328(1985)402

Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-

cyclodextrin as mobile phase component 329(1985)206

Rapid separation of enantiomers by TLC on a chiral stationary phase 330(1985)375

Separation of free radicals by HPLC with electron spin resonance detection 332(1985)71

Counter-ion effects in partition chromatography 332(1985)203

Single-column separation of aminoethylcysteine and other amino acids 332(1985)275

HPLC analysis of peptides on an Asahipak GS-320 column packed with hydrophilic polymer gel 332(1985)287

Recent developments in isotachophoresis (Review) 334(1985)157

Plasma amino acid analysis by GC. Removal of glucose interference 339(1985)366

Evaluation of combined electrochemical and UV detection in the HPLC analysis of plasma for low-molecular-weight constituents 345(1985)125

Cooling plate for cellulose thin-layer electrophoresis and its application to amino acid analysis 347(1985)199

Amino acids, N-acetyl-O-*tert.*-butyl esters
Chiral recognition mechanisms in the enantioselectivity of chiral hydrogen bonding additives in LSC 349(1985)143

Amino acids, 9-anthryldiazomethane derivatives
Fluorescent labelling of amino acids with 9-anthryldiazomethane and its applications to HPLC 348(1985)425

Amino acids, DABS derivatives
Complete HPLC separation of 4-N,N-dimethylaminoazobenzene-4'-thiohydantoin and 4-dimethylaminoazobenzene-4'-sulphonyl chloride amino acids utilizing the same reversed-phase column at room temperature 349(1985)77

Amino acids, DABTH derivatives
Complete HPLC separation of 4-N,N-dimethylaminoazobenzene-4'-thiohydantoin and 4-dimethylaminoazobenzene-4'-sulphonyl chloride amino acids utilizing the same reversed-phase column at room temperature 349(1985)77

Amino acids, Dns derivatives
Application of micro HPLC to the separation of chiral amino acids 331(1985)99

TLC separations using a temperature gradient 333(1985)69

Determination of neurotransmitter amino acids by HPLC with fluorescence detection 337(1985)391

Amino acids, 9-fluorenylmethylchloroformate derivatives
Selective determination of secondary amino acids using precolumn derivatization with 9-fluorenylmethylchloroformate and reversed-phase HPLC 348(1985)213

Amino acids, heptafluorobutyryl butyl esters
GC of amino acids in urine and haemofiltrate 345(1985)19

Amino acids, N-heptafluorobutyryl isobutyl esters
GLC of amino acids. Variable arginine response 347(1985)179

Amino acids, OPA derivatives
High-speed LC analysis of amino acids by post-column sodium hypochlorite-*o*-phthalaldehyde reaction 327(1985)17

Determination of amino acids by separation of their ion pairs with dodecyl sulphate 341(1985)11

HPLC determination of plasma free trimethyllysine in humans 344(1985)319

Amino acids, OPA-*tert.*-butyl thiol derivatives
Determination of amino acids in the brain by HPLC with isocratic elution and electrochemical detection 347(1985)320

Amino acids, phenylthiocarbamyl derivatives
Rapid analysis of amino acids using pre-column derivatization 336(1985)93

Determination of phenylthiocarbamyl amino acids by reversed-phase HPLC 350(1985)453

Amino acids, PTC derivatives
Separation of phenylthiocarbamyl amino acids by HPLC on Spherisorb octadecylsilane columns 346(1985)413

Amino acids, PTH derivatives
Microscale isocratic separation of phenylthiohydantoin amino acid derivatives 326(1985)321

Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1

Electrokinetic chromatography with micellar solutions. Separation of phenylthiohydantoin-amino acids 332(1985)219

High-sensitivity phenylthiohydantoin amino acid analysis using conventional and microbore chromatography 336(1985)105

HPLC support for the baseline separation of all phenylthiohydantoin amino acids 342(1985)376

Separation of phenylthiohydantoin-amino acids by HPLC and some applications in dansyl Edman sequence analysis 349(1985)99

Automated on-line identification of phenylthiohydantoin-amino acids from a vapor phase protein sequencer 350(1985)217

Amino acids, N-TFA *n*-butyl esters
Oligo-oxaalkanoyl tetraamides derived from L-phenylalanine as stationary phases in capillary GC resolution of D,L-amino acids 349(1985)9

Amino acids, N(O,S)-trifluoroacetyl *n*-propyl esters
GLC determination of free amino acids in fermentation broth during growth of *Clostridium oncolyticum* M 55 328(1985)241

Aminoanthraquinones
Drug level monitoring: cytostatics (Review) 340(1985)31

4-Aminoantipyrine
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

HPLC determination of naphthols as 4-aminoantipyrine derivatives. Application to carbaryl 349(1985)63

4-Aminoazobenzene
Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109

4-Aminobiphenyl
Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109

γ-Aminobutyric acid
Reversed-phase HPLC method for determination of brain glutamate decarboxylase suitable for use in kinetic studies 336(1985)385

Rapid GC-MS quantitation of γ-aminobutyric acid in biological specimens 337(1985)21

Determination of amino acids by separation of their ion pairs with dodecyl sulphate 341(1985)11

Amino compounds
LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

Aminoethylcysteine
Single-column separation of aminoethylcysteine and other amino acids 332(1985)275

4-Amino-5-ethyl-3-thiophenecarboxylic acid methyl ester
Determination of 4-amino-5-ethyl-3-thiophenecarboxylic acid methyl ester and its acid metabolite in plasma and urine by HPLC 345(1985)333

Aminoglycosides
Analysis of aminoglycoside antibiotics as benzoyl derivatives by HPLC and its application to the quantitation of neomycin in the perilymph 337(1985)187

Antibiotic monitoring in body fluids (Review) 340(1985)361

Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Aminoisocyanates
Capillary GC method for the determination of complex mixture of isocyanates and amines 346(1985)191

7-Aminonitrazepam
Determination of nitrazepam and its main metabolites in urine by TLC and direct densitometry 339(1985)163

2-Amino-5-nitrobenzophenone
Determination of nitrazepam and its main metabolites in urine by TLC and direct densitometry 339(1985)163

2-Amino-4-nitrophenol
Plasma levels and kinetic disposition of 2,4-dinitrophenol and its metabolites 2-amino-4-nitrophenol and 4-amino-2-nitrophenol in the mouse 344(1985)177

4-Amino-2-nitrophenol
Plasma levels and kinetic disposition of 2,4-dinitrophenol and its metabolites 2-amino-4-nitrophenol and 4-amino-2-nitrophenol in the mouse 344(1985)177

Aminopeptidase M
Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399

Aminophenols
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

N-(3-Aminopropyl)-N^1-(2-carboxyethyl)-1,4-diaminobutane
Mass fragmentographic identification of polyamine metabolites in the urine of normal persons and cancer patients, and its relevance to the use of polyamines as tumour markers 339(1985)223

16(*S*)-Aminoprostaglandin $F_{2\alpha}$ methyl ester

Selective trimethylsilylation for the determination of 16(*S*)-amino-$PGF_{2\alpha}$ methyl ester by GLC 333(1985)404

p-Aminosalicylic acid
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Amino steroids
Chromatography of bis-quaternary amino steroids. I. Separation on silica by TLC and HPLC 328(1985)279

Chromatography of bis-quaternary amino steroids. II. Separation on chemically bonded silica phases 347(1985)284

Amino sugars
Modifications in the alditol acetate method for analysis of muramic acid and other neutral and amino sugars by capillary GC-MS with selected ion monitoring 347(1985)109

β-Aminothiols
Off-line LC-MS studies of fluorescent β-aminothiol-*o*-phthalaldehyde derivatives 330(1985)307

δ-Aminovaleric acid
Mass fragmentographic identification of polyamine metabolites in the urine of normal persons and cancer patients, and its relevance to the use of polyamines as tumour markers 339(1985)223

Amitriptyline
Determination of citalopram, amitriptyline and clomipramine in plasma by reversed-phase HPLC 338(1985)171

Amodiaquine
HPLC analysis of amodiaquine in human plasma 337(1985)166

Amoxapine
Determination of amoxapine and metabolites in plasma by LC with electrochemical detection 338(1985)225

Amoxycillin
Automated determination of amoxycillin in biological fluids by column switching in ion-pair reversed-phase LC systems with post-column derivatization 344(1985)285

cAMP-dependent kinase
Separation of tryptic phosphopeptides of ribosomal origin by reversed-phase HPLC 336(1985)51

cAMP-dependent protein kinase
Combined recycling affinity column-filtration technique 346(1985)247

Amperometric detector
Expert system for pharmaceutical analysis. I. Selection of the detection system in HPLC analysis: UV *versus* amperometric detection 348(1985)97

Amperometric detector for use with small-bore LC columns 350(1985)169

Amphetamine
Use of bonded-phase silica sorbents for rapid sampling of impurities in illicit amphetamine for HPLC analyses 331(1985)339

Amphotericin B
Reproducible measurement of amphotericin B in serum by HPLC in alkaline buffer 337(1985)423

Ampicillin
Studies on residual antibacterials in foods. IV. Simultaneous determination of penicillin G, penicillin V and ampicillin in milk by HPLC 348(1985)379

Amsacrine
Drug level monitoring: cytostatics (Review) 340(1985)31

α-Amylase
Ion-exchange chromatography of fluorogenic derivatives of maltooligosaccharides for preparation of α-amylase substrates 336(1985)368

Anabolic steroids
Selectivity of a diol phase HPLC system in trace analysis of anabolic compounds 347(1985)375

Anaesthetic agents
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Androsterone
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 341(1985)426

Angelicin
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Angiotensin I converting enzyme
Rapid purification of human placental angiotensin I converting enzyme by captopril affinity chromatography 339(1985)399

Angiotensins
Weak anion-exchange HPLC of peptides (Review) 334(1985)49

Separation of biologically important angiotensin peptides by HPLC on a weak cationic exchange bonded phase 336(1985)392

Anhydrochymotrypsin
High-performance affintiy chromatography of anhydrochymotrypsin on a hydrophilic vinyl-polymer gel coupled with tryptophan 350(1985)265

Aniline
Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109

Measurement of trace aromatic amines in seawater using HPLC with electrochemical detection 348(1985)265

2-Anilinoethanol
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

Anion exchangers
Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Performance characteristics of some commercially available low-capacity anion-exchange columns suitable for non-suppressed ion chromatography 346(1985)139

Preparation of low-capacity anion-exchange resins for ion chromatography on a methacrylic copolymer matrix 350(1985)296

Anions
Chromatography of radiolabelled anions using reversed-phase LC columns 333(1985)269

Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Occurrence and origin of system peaks in non-suppressed ion chromatography of inorganic anions with indirect UV absorption detection 346(1985)125

Performance characteristics of some commercially available low-capacity anion-exchange columns suitable for non-suppressed ion chromatography 346(1985)139

Pentafluorobenzyl *p*-toluenesulphonate as a new derivatizing reagent for GC determination of anions 346(1985)215

Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Trace analysis of anions by use of a back-flush method and large injection volumes in ion chromatography 350(1985)317

Anthelmintics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Anthracycline glycosides
Preparative chromatography of epimers and anomers of daunomycin derivatives on Sephadex LH-20 329(1985)193

Anthracyclines
New HPLC assay for plasma doxorubicin 337(1985)194

Anthracyclinones
Preparative chromatography of epimers and anomers of daunomycin derivatives on Sephadex LH-20 329(1985)193

Anthralin
Analysis of anthralin in dermatological products by reversed-phase HPLC 328(1985)317

Anthranilate synthase
HPLC of anthranilate synthase using gel filtration and a post-column reactor 350(1985)332

Anthraquinone glycosides
Drug level monitoring: cytostatics (Review) 340(1985)31

Anthraquinones
Determination of anthraquinones in pulping materials by HPLC using online post-column derivatization 328(1985)309

9-Anthryldiazomethane
Fluorescent labelling of amino acids with 9-anthryldiazomethane and its applications to HPLC 348(1985)425

Anti-arrhythmic agents
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Anti-arteriosclerotics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Anti-asthmatic drugs
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Antibacterials
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antibiotics
Preparative chromatography of epimers and anomers of daunomycin derivatives on Sephadex LH-20 329(1985)193

Separation of erythromycin and related substances by HPLC on poly(styrene-divinylbenzene) packing materials 330(1985)275

Derivatization of secondary amines with 2-naphthalenesulfonyl chloride for HPLC analysis of spectinomycin 333(1985)365

Micromethod for the determination of cefotaxime and desacetylcefotaxime in plasma and urine by HPLC 336(1985)438

Modification of a new HPLC method for bleomycin to separate epi-, iso-, desamido-, and unmodified analogues 337(1985)182

Analysis of aminoglycoside antibiotics as benzoyl derivatives by HPLC and its application to the quantitation of neomycin in the perilymph 337(1985)187

New HPLC assay for plasma doxorubicin 337(1985)194

Determination of the antibiotic fludalanine in plasma and urine by HPLC using a packed-bed, post-column reactor with *o*-phthalaldehyde and 2-mercaptoethanol 338(1985)357

HPLC determination of cefonicid in human plasma and urine 339(1985)359

Drug level monitoring: cytostatics (Review) 340(1985)31

Antibiotic monitoring in body fluids (Review) 340(1985)361

HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 341(1985)115

Determination of coumermycin A_1 in plasma by reversed-phase HPLC analysis 342(1985)145

HPLC determination of apalcillin in plasma and urine 342(1985)234

Determination of benzylpenicillin in plasma and urine by HPLC 342(1985)436

Determination of streptomycin in serum by HPLC 343(1985)379

Automated HPLC determination of spiramycin by direct injection of plasma, using column-switching for sample clean-up 344(1985)275

Automated determination of amoxycillin in biological fluids by column switching in ion-pair reversed-phase LC systems with post-column derivatization 344(1985)285

Isolation of erythromycins and related substances from fermentation residues of *Streptomyces erythreus* by HPLC on silica gel 346(1985)309

Reversed-phase HPLC method for the assay of oxytetracycline 347(1985)173

Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Studies on residual antibacterials in foods. IV. Simultaneous determination of penicillin G, penicillin V and ampicillin in milk by HPLC 348(1985)379

Anticholinergics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Anticoagulants
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antidepressant drugs
Analysis of tricyclic antidepressant drugs in plasma and serum by chromatographic techniques (Review) 340(1985)173

Antidiabetics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Anti-emetics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Anti-epileptic drugs
Determination of antiepileptic drugs (Review) 340(1985)199

Antihistamines
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antihypertensive agents
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Antimalarials
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antimycotics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antiparasitics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antiparkinsonics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antiprotozoal drugs
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antipyretics
HPLC of antipyretics on chemically modified porous glass 332(1985)265

Analysis of antipyretics by semimicro LC 332(1985)269

Antipyrine
HPLC method for the simultaneous determination of cimetidine and antipyrine in plasma 345(1985)192

Antirheumatic drugs
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77

Antisympathetic drugs
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

α_1-Antitrypsin
Isolation and characterization of two minor fractions of α_1-antitrypsin by HPLC chromatofocusing 327(1985)173

Antituberculosis drugs
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Antitussives
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antiulcer drugs
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Antiviral compounds
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Apalcillin
HPLC determination of apalcillin in plasma and urine 342(1985)234

Apohaemopexin
Purification of hemopexin and its domain fragments by affinity chromatography and HPLC 326(1985)373

Apolipoproteins
HPLC of apolipoproteins in serum high-density lipoproteins 336(1985)151

Evaluation of immunoaffinity chromatography for isolating human lipoproteins containing apolipoprotein B 342(1985)47

Characterization of human apolipoprotein A-I by reversed-phase HPLC 345(1985)145

Apomorphine
Determination of apomorphine in rat plasma and brain by HPLC with electrochemical detection 338(1985)230

Apparent density
Quality criteria and structure of silica gel column packing material 329(1985)351

Appetite depressants
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Arabic gum
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Arbaprostil
Determination of (15*R*)- and (15*S*)-15-methylprostaglandin E_2 in human plasma with picogram per milliliter sensitivity by column-switching HPLC 343(1985)271

Arginine
GLC of amino acids. Variable arginine response 347(1985)179

Arginine-vasopressin
Quantitative affinity HPLC of neuroendocrine polypeptides using porous and non-porous glass derivatives 327(1985)193

Aroclors
Vapor pressure estimates of individual polychlorinated biphenyls and commercial fluids using GC retention data 330(1985)203

Recent advances in the analysis of polychlorinated biphenyls in environmental and biological media (Review) 334(1985)277

Aromatic acids
Energy effects in the retention of aromatic acids in LC 332(1985)189

Aromatic compounds
GC determination of total aromatics in kerosene 328(1985)207

Long pathlength UV absorbance detection for GC with concurrent fluorescence measurement 331(1985)77

Quantitative analysis of hydrocarbons in gasolines by capillary GLC. II. Isothermal and temperature-programmed analyses 346(1985)177

Group separation of oil residues by supercritical fluid chromatography 349(1985)439

Arsenate
Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Arylacetic acids
TLC and HPLC in the evaluation of the lipophilicity of aryloxoalkanoic and arylhydroxyalkanoic acids 347(1985)335

Arylaliphatic acids
Relationships between GLC behaviour and structure of arylaliphatic acids. 333(1985)21

Aryl alkyl sulphides
Ligand-exchange GC separation of aromatic sulphides 350(1985)75

N-Aryl-α-amino esters
Chromatographic separation of the enantiomers of 2-carboalkoxyindolines and N-aryl-α-amino esters on chiral stationary phases derived from N-(3,5-dinitrobenzoyl)-α-amino acids 348(1985)89

Arylhydroxyalkanoic acids
TLC and HPLC in the evaluation of the lipophilicity of aryloxoalkanoic and arylhydroxyalkanoic acids 347(1985)335

Aryloxoalkanoic acids
TLC and HPLC in the evaluation of the lipophilicity of aryloxoalkanoic and arylhydroxyalkanoic acids 347(1985)335

1-Arylpiperazines
Ionization constants and lipophilicity of 1-arylpiperazines investigated by reversed-phase HPLC 333(1985)191

Asahipak GS-gel
High-performance affinity chromatography of trypsins on Asahipak GS-gel coupled with *p*-aminobenzamidine 333(1985)107

β-Asarone
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

GC and MS studies of the constituents of the rhizome of calamus. II. The volatile constituents of alcoholic extracts 328(1985)195

Ascorbic acid
Analysis of ascorbic acid derivatives by HPLC with electrochemical detection 330(1985)408

Separation of ascorbic acid, dehydroascorbic acid, diketogulonic acid and glucose by isocratic elution from a column of a hydrophilic gel 332(1985)283

Determination of total ascorbic acid in human serum by HPLC with fluorescence detection 344(1985)351

Aspartame
HPLC analysis of Aspartame 346(1985)417

Ion-pair HPLC analysis of aspartame and related products 349(1985)83

Aspartate aminotransferase
Sequence determination of cyanogen bromide-peptides by combined use of HPLC and fast atom bombardment MS 331(1985)425

Aspochalasins
TLC of mycotoxins (Review) 334(1985)211

Atenolol
HPLC determination of sotalol in biological fluids 336(1985)313

Simple and rapid analysis of atenolol and metoprolol in plasma using solid-phase extraction and HPLC 339(1985)429

Na^+,K^+-ATPase inhibitors
Rapid large-scale isolation of biologically active molecules using reversed-phase "flash" chromatography: initial purification of endogenous Na^+,K^+-ATPase inhibitors from human urine 338(1985)417

Atracurium
Determination of atracurium, laudanosine and related compounds in plasma by HPLC 343(1985)431

Atrazine
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Atrial natriuretic factor
Novel approach to rapid and sensitive localization of protein disulfide bridges by HPLC and electrochemical detection 326(1985)339

Atropine
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Automation
Automation of analytical isotachophoresis (Review) 334(1985)83

Aza-arenes
Preseparation of polar, substituted, polycyclic aromatic hydrocarbons in aerosol extracts by HPLC on acidic- or basic-buffered silica 349(1985)331

6-Azacytidine
Reversed-phase ion-paired HPLC determination of 6-azacytidine in blood 345(1985)157

5-Aza-2'-deoxycytidine
Sample preparation for the determination of 5-aza-2'-deoxycytidine in plasma by HPLC 345(1985)162

Aztreonam
Antibiotic monitoring in body fluids (Review) 340(1985)361

B

Back-flush method
Trace analysis of anions by use of a back-flush method and large injection volumes in ion chromatography 350(1985)317

Baclofen
Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59

HPLC analysis of baclofen in plasma and urine of man after precolumn extraction and derivatization with *o*-phthaldialdehyde 337(1985)341

Determination of 4-amino-3-(*p*-chlorophenyl)butyric acid (baclofen) in plasma by HPLC 339(1985)424

Bacterial growth
Assessment of head-space GLC for the rapid detection of growth in blood cultures 337(1985)223

Bacteriorhodopsin
Isolation of detergent-solubilized monomers of bacteriorhodopsin by size-exclusion HPLC 326(1985)243

Rapid delipidation of the membrane protein bacteriorhodopsin by HPLC 350(1985)427

Baicalein
Simultaneous determination of baicalein, wogonin, oroxylin-A and their glucuronides in Scutellariae Radix by ion-pair HPLC 328(1985)289

Baicalin
Simultaneous determination of baicalein, wogonin, oroxylin-A and their glucuronides in Scutellariae Radix by ion-pair HPLC 328(1985)289

Balfourodinium
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Bamethan
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Bamifylline
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Band broadening
Investigation of the mechanism of peak broadening observed in the HPLC analysis of cyclosporine 333(1985)231

Band width
Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27

HPLC theory for the practitioner (Review) 334(1985)197

Barbiturates
Improved method for the preparation of immobilised pyrimidonesilicones as polar stationary phases used for the capillary column GC separation of underivatised barbiturates 329(1985)213

Nitrogen-sensitive thermionic detection in microcolumn LC 333(1985)291

Drug level monitoring: Sedative hypnotics (Review) 340(1985)139

Determination of antiepileptic drugs (Review) 340(1985)199

Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Baseline drift
Spectral properties of trifluoroacetic acid-acetonitrile gradient systems for separation of picomole quantities of peptides by reversed-phase HPLC 347(1985)83

Base strength
Effects of various strengths of bases on the electrophoretic behavior of weak acids in a non-aqueous solvent 330(1985)19

Bathophenanthrolinedisulphonic acid disodium salt
Simultaneous determination of iron(II) and -(III) by ion chromatography with post-column reaction 329(1985)247

Bencyclane
GLC evaluation of bencyclane in biological samples for pharmacokinetic and bioavailability investigations: comparison of two analytical methods 343(1985)77

Benomyl
Simplified method for the clean-up and reversed-phase HPLC determination of benomyl in mangoes 330(1985)403

Benoxaprofen chloride
Fluorimetric determination of the quaternary compound trospium and its metabolite in biological material after derivatization with benoxaprofen chloride 345(1985)99

Bentiromide
Analysis of N-benzoyl-L-tyrosyl-*p*-aminobenzoic acid (bentiromide) metabolites in urine by ion-pair HPLC 338(1985)377

Benzene
Adsorption of mixtures of acetone and benzene vapors on graphitized carbon black 328(1985)63

Sorption properties of some modified molecular sieves 13X towards thiophene and benzene 329(1985)128

Comparison between theoretical and experimental sampling efficiencies on Tenax GC 331(1985)207

Benzenes
Methylated cyclodextrins as mobile phase additives in LC 330(1985)253

Benzenesulphonamides
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Benzidine
Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109

Benzisoquinolinedione
Sensitive reversed-phase HPLC assay for the antitumor agent benzisoquinolinedione (Nafidimide) employing direct on-column injection of plasma and urine, and its use for pharmacokinetic studies in a dog 345(1985)402

HPLC separation and quantitation of benzisoquinolinedione (Nafidimide) from biological fluids 345(1985)408

Benzodiazepines
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Benzoic acid
Studies on the analysis of food additives by HPLC. V. Simultaneous determination of preservatives and saccharin in foods by ion-pair chromatography 346(1985)333

***p*-Benzoquinones**
HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. II. Retention behavior, quantitative structure-retention relationships and octanol-water partition coefficients 330(1985)181

Benzotriazoles
Rapid chromatographic determination of benzotiazoles in automotive cooling waters and cooling water formulations 331(1985)149

S-Benzoyl captopril
Determination of S-benzoyl captopril in human urine by capillary GC with electron-capture detection 342(1985)186

Benzoyl esters
GLC analyses. XLIV. Benzoyl and pentafluorobenzoyl esters of lower saturated branched-chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 329(1985)359

Benzylideneamines
Retention indices of dimethylbenzamidines and benzylideneamines on a non-polar column 330(1985)79

Benzylpenicillin
Determination of benzylpenicillin in plasma and urine by HPLC 342(1985)436

Bergapten
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Bestatin
Determination of *p*-hydroxybestatin in human serum by HPLC using fluorescence detection 344(1985)267

Betaine
Determination of carnitine, butyrobetaine, and betaine as 4'-bromophenacyl ester derivatives by HPLC 336(1985)271

Betamethasone
Quantitative determination of betamethasone and its major metabolite in equine urine by micro-LC-MS 341(1985)261

Betaxolol
Determination of betaxolol in plasma by HPLC with fluorescence detection 342(1985)212

Bezafibrate
Determination of bezafibrate in human plasma and urine by HPLC 344(1985)259

Bicyclo[*n*.4.0]alken-2-ones
GC of bicyclo[*n*.4.0]alken-2-ones 331(1985)406

Bile acid 3-glucuronides
Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 345(1985)241

Studies on steroids. CCXVI. Separation of bile acid 3-glucuronides by HPLC 348(1985)151

Bile acids
Bile acids as an example to demonstrate the methodological aspects of the quantitative HPTLC analysis of fluorescent and non-fluorescent compounds (in German) 333(1985)276

Development and validation of a method for measuring the glycine and taurine conjugates of bile acids in bile by HPLC 336(1985)249

HPLC analysis of bile acids in hamster bile 337(1985)91

Determination of bile acids in rat bile by HPLC 338(1985)195

Rapid and accurate reversed-phase HPLC determination of conjugated bile acids in human bile for routine clinical applications. Therapeutic control during gallstone dissolution therapy 339(1985)263

Extraction and fractionation of bile acids and their conjugates using pre-packed microparticulate silica cartridges (Sep-Pak Sil and Bond-Elut C_{18}) 343(1985)259

Bile acids. LXXVI. Analyses of bile acids and sterols by HPLC with a microbore column 347(1985)155

Binaphthol derivatives
Cinchona alkaloids for preparing new, easily accessible chiral stationary phases. II. Resolution of binaphthol derivatives on silica-supported quinine 348(1985)79

Bio-Gel TSK-Phenyl-5-PW
Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

Biomer
Chromatographic analysis of thermoplastic, medical-grade, polyether-based polyurethane elastomers 349(1985)347

Biopolymers
Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33

Affinity partitioning of biopolymers and membranes in Ficoll-dextran aqueous two-phase systems 331(1985)11

Biotin
HPLC of biotin and analogues 330(1985)153

Biotin-labelled nucleotides
Reversed-phase LC of biotin-labelled nucleotides. A new class of markers in molecular biology 330(1985)429

4,4'-Bipyridyl quaternary salts
Separation of organic quaternary salts by reversed-phase TLC 346(1985)463

Bisbenzimidazole
HPLC assay and preclinical pharmacological studies of pibenzimol (bisbenzimidazole) 341(1985)89

N,N^1-Bis(2-carboxyethyl)-1,4-diaminobutane
Mass fragmentographic identification of polyamine metabolites in the urine of normal persons and cancer patients, and its relevance to the use of polyamines as tumour markers 339(1985)223

Bis(*p*-chlorophenyl) sulphone
Determination of the content of 2-chlorobenzenesulphonamide and bis(*p*-chlorophenyl) sulphone in 4-chlorobenzenesulphonamide 333(1985)253

Bis-quaternary amino steroids
Chromatography of bis-quaternary amino steroids. I. Separation on silica by TLC and HPLC 328(1985)279

Chromatography of bis-quaternary amino steroids. II. Separation on chemically bonded silica phases 347(1985)284

Bitertanol
Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181

Biuret
Determination of biuret and urea by HPLC 331(1985)161

Bleomycin
Modification of a new HPLC method for bleomycin to separate epi-, iso-, desamido-, and unmodified analogues 337(1985)182

Drug level monitoring: cytostatics (Review) 340(1985)31

Blue Dextran
Heterogeneities of blue dextran and differences among samples 350(1985)292

Boiling point
Capillary GC of chloro derivatives of 1,4-dimethylbenzene. Separation, identification and prediction of boiling points 331(1985)237

Bracketing
Bracketing, a simple loading technique that increases sample recovery, improves resolution, and increases sensitivity in HPLC 348(1985)123

Bradykinins
Weak anion-exchange HPLC of peptides (Review) 334(1985)49

Breakthrough curves
Comparison between theoretical and experimental sampling efficiencies on Tenax GC 331(1985)207

Breakthrough volumes
Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Evaluation of the suitability of selected porous polymers for preconcentration of organosulphur compounds from water 346(1985)61

Bromide
LC determination of bromide in human milk and plasma 344(1985)157

1-Bromoalkanes
Temperature-programmed retention indices of 221 halogenated organic compounds with 1-bromoalkanes as references 328(1985)35

Bromodeoxyuridine
HPLC method for the determination of bromodeoxyuridine and its major metabolite, bromouracil, in biological fluids 341(1985)217

5-(2-Bromo-*E*-ethenyl)-2'-deoxyuridine
Simultaneous HPLC assay for 5-(2-bromo-*E*-ethenyl)-2'-deoxyuridine and its metabolite, 5-(2-bromo-*E*-ethenyl)uracil, in plasma 338(1985)219

5-(2-Bromo-*E*-ethenyl)uracil
Simultaneous HPLC assay for 5-(2-bromo-*E*-ethenyl)-2'-deoxyuridine and its metabolite, 5-(2-bromo-*E*-ethenyl)uracil, in plasma 338(1985)219

3-Bromomethyl-6,7-dimethoxy-1-methyl-2(1*H*)-quinoxalinone
3-Bromomethyl-6,7-dimethoxy-1-methyl-2(1*H*)-quinoxalinone as a new fluorescence derivatization reagent for carboxylic acids in HPLC 346(1985)227

4-Bromomethyl-7-methoxycoumarin
4-methyl-7-methoxycoumarin as a fluorescent label for HPLC analysis of dicarboxylic acids 328(1985)111

2-(5-Bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol

HPLC of copper, cobalt, nickel, chromium and vanadium chelates with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol 333(1985)133

Bromouracil
HPLC method for the determination of bromodeoxyuridine and its major metabolite, bromouracil, in biological fluids 341(1985)217

Brush-type grafted monolayers
Insertion of various long alkyl chain molecules in brush-type grafted monolayers. Chromatographic study of the resulting materials 330(1985)227

Buffered silica gel
Adsorption chromatographic separations on buffered silica gel (Review) 334(1985)35

Buffer-memory technique
Supercritical fluid chromatography-IR spectroscopy of oligomers: use of buffer-memory technique 332(1985)47

Buffers
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Bulk solution partition
Contributions to hydrocarbon retention in gas-liquid-solid chromatography with polyethylene glycol 6000 as stationary phase 350(1985)15

Buprenorphine
^{63}Ni electron-capture GC assay for buprenorphine and metabolites in human urine and feces 337(1985)291

Analysis of buprenorphine and its N-dealkylated metabolite in plasma and urine by selected-ion monitoring 338(1985)89

Analysis of buprenorphine by HPLC 347(1985)411

Buspirone
Capillary GC-MS determination of buspirone in plasma 345(1985)285

Busulfan
Drug level monitoring: cytostatics (Review) 340(1985)31

Butanone
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

Butenolide
TLC of mycotoxins (Review) 334(1985)211

4-[3-(*tert.*-Butylamino)-2-hydroxypropoxy]-N-methylisocarbostyril
HPLC of a new β-blocker, 4-[3-(*tert.*-butylamino)-2-hydroxypropoxy]-N-methylisocarbostyril hydrochloride, in plasma using fluorometric detection 339(1985)219

***tert.*-Butylated cresols**
Analysis of *tert.*-butylated cresol mixtures by capillary GC and capillary GC-MS 348(1985)430

Butylated hydroxytoluene
Derivatization of butylated hydroxytoluene for GC with electron-capture detection and HPLC with UV detection 333(1985)171

Butyl flufenamate
HPLC assay of anti-inflammatory drugs incorporated in gel ointments. Separation and stability testing 331(1985)383

Butyrobetaine
Determination of carnitine, butyrobetaine, and betaine as 4'-bromophenacyl ester derivatives by HPLC 336(1985)271

C

Cadaverine
Separation of cadaverine from putrescine, histamine and polyamines in rat kidney by phosphocellulose chromatography 337(1985)103

Frequency-pulsed electron-capture GLC studies of chemical changes in sera of patients with schistosomiasis 339(1985)243

Caeruloplasmin
Automated tandem HPLC system for separation of extremely complex peptide mixtures 326(1985)407

Cafestol
Use of silver nitrate impregnated silica cartridges in the separation of kahweol and cafestol esters by preparative LC 328(1985)422

Caffeine
New double-stage separation analysis method. Directly coupled laboratory-scale supercritical fluid extraction-supercritical fluid chromatography, monitored with a multiwavelength-UV detector 332(1985)107

HPLC of antipyretics on chemically modified porous glass 332(1985)265

Analysis of antipyretics by semimicro LC 332(1985)269

Improved micro-method for the HPLC determination of caffeine and paraxanthine in biological fluids 336(1985)293

HPLC determination of dimethylxanthine metabolites of caffeine in human plasma 338(1985)369

Improved HPLC method for the simultaneous determination of caffeine and its N-

demethylated metabolites in plasma using solid-phase extraction 342(1985)105

Caffeoylquinic acids
HPLC of caffeoylquinic acid derivatives of *Cynara scolymus* L. leaves 348(1985)447

Calamus
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

GC and MS studies of the constituents of the rhizome of calamus. II. The volatile constituents of alcoholic extracts 328(1985)195

Calcium
Improvement of ion chromatography with UV photometric detection and comparison with conductivity detection for the determination of serum cations 341(1985)53

Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251

Calcium antagonists
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Calmodulin
Purification and peptide mapping of calmodulin and its chemically modified derivatives by reversed-phase HPLC 326(1985)387

CAMP-factor
Purification and characterization of CAMP-factor from *Streptococcus agalactiae* by hydrophobic interaction chromatography and chromatofocusing 348(1985)363

Cannabinoids
Antibody-mediated extraction of the main tetrahydrocannabinol metabolite, 11-nor-Δ^9-tetrahydrocannabinol-9-carboxylic acid, from human urine and its identification by GC-MS in the sub-nanogram range 342(1985)393

Canrenone
Determination of the serum concentration of spironolactone and its metabolites by HPLC 341(1985)279

Cantharidin
Quantitative determination of cantharidin in biological materials using capillary GC with flame ionization detection 342(1985)411

Capacity ratio
A problem of optimization in capillary GC 348(1985)416

Capillary columns
see also Columns and microbore columns

Preparation of glass and fused-silica capillary columns coated with immobilized XE-60 346(1985)341

New approach to capillary columns for GC? Condensation of hydroxyl-terminated stationary phases 347(1985)351

Capillary connections
Connecting capillaries by fusion 330(1985)217

Capillary isotachophoresis
Theoretical and practical limitations of capillary isotachophoresis in trace analysis 328(1985)353

Capillary tube length
HPLC theory for the practitioner (Review) 334(1985)197

Capreomycin
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Capsaicin
Formation and metabolism of pungent principle of *Capsicum* fruits. XV. Microdetermination of capsaicin by HPLC with electrochemical detection 329(1985)99

Captopril
Determination of captopril and its mixed disulphides in plasma and urine by HPLC 338(1985)161

Rapid purification of human placental angiotensin I converting enzyme by captopril affinity chromatography 339(1985)399

Determination of S-benzoyl captopril in human urine by capillary GC with electron-capture detection 342(1985)186

Carbamates
Reversed-phase ion-interaction chromatography of sodium dialkyldithiocarbamates in the presence of tetraalkylammonium salts 348(1985)389

Carbamazepine
Concurrent determination of valproic acid with other antiepileptic drugs by HPLC 338(1985)131

Determination of antiepileptic drugs (Review) 340(1985)199

Simultaneous determination of carbamazepine and its epoxide metabolite in plasma and urine by HPLC 342(1985)341

Simultaneous determination of anticonvulsant drugs and metabolites in plasma by HPLC 342(1985)469

Carbaryl
HPLC determination of naphthols as 4-aminoantipyrine derivatives. Application to carbaryl 349(1985)63

Carbazole alkaloids
Separation of carbazole alkaloids by GLC 329(1985)178

Carbazoles
Preseparation of polar, substituted, polycyclic aromatic hydrocarbons in aerosol extracts by HPLC on acidic- or basic-buffered silica 349(1985)331

2-Carboalkoxyindolines
Chromatographic separation of the enantiomers of 2-carboalkoxyindolines and N-aryl-α-amino esters on chiral stationary phases derived from N-(3,5-dinitrobenzoyl)-α-amino acids 348(1985)89

Carbohydrates
see also Sugars and saccharides

Amperometric detection of reducing carbohydrates in HPLC using an amino-bonded column and acetonitrile-water as the eluent 330(1985)333

Radioactivity detection system with a CaF_2 (Eu) scintillator for HPLC 333(1985)83

Isolation and quantitation of carbohydrates in sheep plasma by HPLC 338(1985)303

Purification of the peracetylated glycosphingolipids of the Gala series (galactosyl- and galabiosylceramides) 342(1985)385

HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Carbon dioxide
On the possibility of increasing the effectiveness of chromatographic columns filled with zeolites 331(1985)402

Carbon disulphide
GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

Carbon monoxide
Measurement of mass and radioactivity of ^{14}CO using a reduction detector 329(1985)137

Carbon-silica adsorbents
Microscopic investigation of the surface of carbon-silica adsorbents. II. Relationships between the type of information obtainable about the surface and the microscopic techniques for its examination 346(1985)346

Carbon tetrachloride
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233

Carbonyl compounds
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

Determination of low concentrations of low-molecularweight aldehydes and ketones in aqueous samples 329(1985)81

Carbonyl sulphide
On-line analysis of sulphur compounds in natural gas with flame photometric detection 349(1985)465

Carbosils
Microscopic investigation of the surface of carbon-silica adsorbents. II. Relationships between the type of information obtainable about the surface and the microscopic techniques for its examination 346(1985)346

1-(4'-Carboxybutyl)-3,7-dimethylxanthine
Determination of 1-(3'-carboxypropyl)-3,7-dimethylxanthine and 1-(4'-carboxybutyl)-3,7-dimethylxanthine, two major metabolites of oxpentifylline, in urine by HPLC 344(1985)397

N-(2-Carboxyethyl)-4-amino-*n*-butyric acid
Mass fragmentographic identification of polyamine metabolites in the urine of normal persons and cancer patients, and its relevance to the use of polyamines as tumour markers 339(1985)223

Carboxylic acids
see also Organic acids

4-methyl-7-methoxycoumarin as a fluorescent label for HPLC analysis of dicarboxylic acids 328(1985)111

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. II. Influence of organic co-solvents on retention at 25°C 333(1985)41

Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59

Reversed-phase ion-pair chromatography of some biologically important carboxylic acids as 2-nitrophenylhydrazides 333(1985)215

HPLC of microbial acid metabolites 336(1985)125

Rapid and selective derivatization method for the nitrogen-selective detection of carboxylic acids in biological fluids prior to GC analysis 336(1985)241

Frequency-pulsed electron-capture GLC studies of chemical changes in sera of patients with schistosomiasis 339(1985)243

GC-MS of carboxylic acids in tissues as their *tert.*-butyldimethylsilyl derivatives 344(1985)1

Photodiode array UV spectrophotometric profiling of carboxylic acids in physiological fluids 344(1985)23

Pentafluorobenzyl *p*-toluenesulphonate as a new derivatizing reagent for GC determination of anions 346(1985)215

3-Bromomethyl-6,7-dimethoxy-1-methyl-2(1*H*)-quinoxalinone as a new fluorescence derivatization reagent for carboxylic acids in HPLC 346(1985)227

HPLC separation of aromatic carboxylic acids with β-cyclodextrin-bonded stationary phases 350(1985)371

2-O-Carboxymethyl-β-cyclodextrin
Electrokinetic chromatography with 2-O-carboxymethyl-β-cyclodextrin as a moving "stationary" phase 332(1985)211

S-Carboxymethylcysteine
Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1

Carboxypeptidase Y
Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399

1-(3'-Carboxypropyl)-3,7-dimethylxanthine
Determination of 1-(3'-carboxypropyl)-3,7-dimethylxanthine and 1-(4'-carboxybutyl)-3,7-dimethylxanthine, two major metabolites of oxpentifylline, in urine by HPLC 344(1985)397

1-Carboxysalsolinol
Identification and quantification of 1-carboxysalsolinol and salsolinol in biological samples by GC-MS 343(1985)1

Cardiolipin
HPLC of cardiolipin 339(1985)35

Cardiothane
Chromatographic analysis of thermoplastic, medical-grade, polyether-based polyurethane elastomers 349(1985)347

Cardiovascular drugs
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Carnitine
Determination of carnitine, butyrobetaine, and betaine as 4'-bromophenacyl ester derivatives by HPLC 336(1985)271

Carob gum
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

β-Carotene
Serum and plasma β-carotene levels measured with an improved method of HPLC 339(1985)273

ε,ε-Carotene-3,3'-diol
Direct diastereomeric resolution of carotenoids. II. All ten stereoisomers of tunaxanthin (ε,ε-carotene-3,3'-diol) 328(1985)387

Carotenoids
Direct diastereomeric resolution of carotenoids. II. All ten stereoisomers of tunaxanthin (ε,ε-carotene-3,3'-diol) 328(1985)387

Decapreno-β-carotene as an internal standard for the quantification of the hydrocarbon carotenoids by HPLC 346(1985)237

Caseins
Rapid analysis of bovine milk proteins by fast protein LC 348(1985)177

Catalase
Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Catastrophe theory
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Catechol
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Catecholamines
Determination of catecholamines by HPLC with catalytic photometric detection 329(1985)147

Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Direct clean-up and analysis of urinary catecholamines 336(1985)374

Simultaneous assay of 3,4-dihydroxyphenylalanine, catecholamines and O-methylated metabolites in human plasma using HPLC 337(1985)239

Simple and fast analysis of adrenaline and noradrenaline in plasma on microbore HPLC columns using fluorimetric detection 337(1985)384

Determination of adult sheep plasma catecholamines using [^{3}H]norepinephrine as the internal standard 338(1985)192

Estimation of adrenal catecholamines by elevated-temperature LC with amperometric detection 342(1985)363

Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370

Determination of intraoperative plasma catecholamine concentrations using LC with electrochemical detection 343(1985)285

Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61

Rapid and reliable estimation of urinary free catecholamines in patients with pheochromocytoma: comparison with plasma

catecholamines and vanillylmandelic acid excretion 344(1985)313

Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 345(1985)140

Cation exchangers
Separation of peptides by strong cation-exchange HPLC 327(1985)147

Cations
Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. I. Cation binding in the mobile phase 329(1985)321

Improvement of ion chromatography with UV photometric detection and comparison with conductivity detection for the determination of serum cations 341(1985)53

Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147

Cavity size
LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

Cefonicid
HPLC determination of cefonicid in human plasma and urine 339(1985)359

Cefoperazone
HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 341(1985)115

Cefotaxime
Micromethod for the determination of cefotaxime and desacetylcefotaxime in plasma and urine by HPLC 336(1985)438

Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131

Cefoxitin
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131

Ceftazidime
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131

3-O-β-Cellobiosyl-D-glucose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Cellotriose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Cellular proteins
Detection of the changes in cellular proteins in regenerating rat liver by high-resolution two-dimensional electrophoresis 343(1985)51

Cellulase
Separation and partial characterization of *Trichoderma reesei* cellulase by fast chromatofocusing 329(1985)379

Isoelectric focusing separation of *Gliocladium* enzyme components 347(1985)310

Cellulose triacetate
Chromatographic resolution of racemates on chiral stationary phases. I. Influence of the supramolecular structure of cellulose triacetate 347(1985)25

Cembrene A
Preparative HPLC of (2*E*)- and (2*Z*)-farnesol 347(1985)188

Cephalexin
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131

Cephalosporins
Micromethod for the determination of cefotaxime and desacetylcefotaxime in plasma and urine by HPLC 336(1985)438

HPLC determination of cefonicid in human plasma and urine 339(1985)359

Antibiotic monitoring in body fluids (Review) 340(1985)361

Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131

Cephazolin
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131

Ceramides
Purification of the peracetylated glycosphingolipids of the Gala series (galactosyl- and galabiosylceramides) 342(1985)385

Cetiedil
Automated capillary GC assay using nitrogen-phosphorus detection for the determination of cetiedil in plasma 345(1985)178

Chaetoglobosins
TLC of mycotoxins (Review) 334(1985)211

Charcoal
Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251

Chaulmoogric oils
HPLC of triglycerides of Flacourtiaceae seed oils containing cyclopentenyl fatty acids (chaulmoogric oils) 348(1985)441

Chemical analyzer
Automation of analytical isotachophoresis (Review) 334(1985)83

Chemical ionization mass spectrometry
Gaseous acidity and basicity scales as guides to chemical ionization in reversed-phase LC-MS with direct liquid introduction 347(1985)61

Chemical warfare agents
GC retention indices of chemical warfare agents and simulants 331(1985)47

Chemiluminescence detection
GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

Chenodeoxycholic acid
Determination of bile acids in rat bile by HPLC 338(1985)195

Rapid and accurate reversed-phase HPLC determination of conjugated bile acids in human bile for routine clinical applications. Therapeutic control during gallstone dissolution therapy 339(1985)263

Chinese herbal preparations
TLC screening procedure for undeclared synthetic drugs in Chinese herbal preparations 329(1985)107

Chiral additives
Application of micro HPLC to the separation of chiral amino acids 331(1985)99

Chiral recognition mechanisms in the enantioselectivity of chiral hydrogen bonding additives in LSC 349(1985)143

Chiral stationary phases
see also Stationary phases

Chiral stationary phases, GC
Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181

Oligo-oxaalkanoyl tetraamides derived from L-phenylalanine as stationary phases in capillary GC resolution of D,L-amino acids 349(1985)9

Chiral stationary phases, LC
Effect of interstrand distance upon chiral recognition by a chiral stationary phase 328(1985)1

Determination of (*R*)- and (*S*)-disopyramide in human plasma using a chiral α_1-acid glycoprotein column 336(1985)321

Molecular imprinting of amino acid derivatives in macroporous polymers. Demonstration of substrate- and enantio-selectivity by chromatographic resolution of racemic mixtures of amino acid derivatives 347(1985)1

Use of silane monomers for molecular imprinting and enzyme entrapment in polysiloxane-coated porous silica 347(1985)11

Chromatographic resolution of racemates on chiral stationary phases. I. Influence of the supramolecular structure of cellulose triacetate 347(1985)25

Cinchona alkaloids for preparing new, easily accessible chiral stationary phases. II. Resolution of binaphthol derivatives on silica-supported quinine 348(1985)79

Chromatographic separation of the enantiomers of 2-carboalkoxyindolines and N-aryl-α-amino esters on chiral stationary phases derived from N-(3,5-dinitrobenzoyl)-α-amino acids 348(1985)89

Resolution of enantiomeric amino acid derivatives by HPLC on chiral stationary phases 348(1985)275

Chromatographic resolution. IX. HPLC resolution of enantiomers on optically active poly(triphenylmethyl methacrylate) 350(1985)127

Application of chromatographic chiral stationary phases to pharmaceutical analysis. Enantiomeric purity of D-penicillamine 350(1985)179

Chiral stationary phases, TLC
Rapid separation of enantiomers by TLC on a chiral stationary phase 330(1985)375

Chlorambucil
Drug level monitoring: cytostatics (Review) 340(1985)31

Determination of chlorambucil in plasma using reversed-phase HPLC 342(1985)447

Chloramphenicol
Antibiotic monitoring in body fluids (Review) 340(1985)361

Chlordanes
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Evaluation of tetraalkylammonium tetrafluoroborate salts as high-temperature stationary phases for packed and open-tubular column GC 349(1985)249

Chlorhexidine
Determination of preservatives in cosmetic products by ion-pair reversed-phase HPLC. III 348(1985)321

Chloride
Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Chlorinated 1,4-dimethylbenzene
Capillary GC of chloro derivatives of 1,4-dimethylbenzene. Separation, identification and prediction of boiling points 331(1985)237

Chlorinated hydrocarbons
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233

Temperature selection for chlorinated hydrocarbon reduction for the Hall electrolytic conductivity detector 328(1985)342

Catalytic hydrogenation and exchange reactions on GC columns, in the presence of supported $(\eta^5\text{-}C_5H_5)NiOs_3(\mu\text{-}H)_3(CO)_9$ 349(1985)173

ω-Chloroacetophenone
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379

7-Chloro-4-aminoquinoline
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 345(1985)209

Chlorobenzenes
Determination of trace chlorinated benzenes in fuel oil by on-line multidimensional chromatography using packed-capillary LC and capillary GC 349(1985)55

2-Chlorobenzenesulphonamide
Determination of the content of 2-chlorobenzenesulphonamide and bis(*p*-chlorophenyl) sulphone in 4-chlorobenzenesulphonamide 333(1985)253

4-Chlorobenzenesulphonamide
Determination of the content of 2-chlorobenzenesulphonamide and bis(*p*-chlorophenyl) sulphone in 4-chlorobenzenesulphonamide 333(1985)253

***o*-Chlorobenzylidenemalononitrile**
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379

2-Chloroethyl esters
GLC analyses. XLIII. Retention increments for 2-chloro-, 2,2-dichloro- and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns 329(1985)43

Chloroform
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233

Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

3-Chloroformyl-7-methoxycoumarin
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 341(1985)426

***R*-(+)-7-Chloro-8-hydroxy-1-phenyl-2,3,4,5-tetrahydro-1H-3-methyl-3-benzazepine**
Quantification of *R*-(+)-7-chloro-8-hydroxy-1-phenyl-2,3,4,5-tetrahydro-1H-3-methyl-3-benzazepine in brain and blood by use of reversed-phase HPLC with electrochemical detection 342(1985)452

7-Chloro-4-hydroxyquinoline
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 345(1985)209

Chlorophenols
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Synthesis and GC-MS determination of polychlorinated dibenzo-*p*-dioxins and related compounds in the technical chlorophenol formulation Ky-5 331(1985)410

Electrokinetic chromatography with micellar solutions. Retention behaviour and separation of chlorinated phenols 348(1985)39

Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

Chloroquine
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 345(1985)209

Chlorosulphonamido diphenyl ethers
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Chlorpropamide
Determination of glibenclamide, chlorpropamide and tolbutamide in plasma by HPLC with UV detection 339(1985)203

Chlorprothixene
Determination of chlorprothixene and its sulfoxide metabolite in plasma by HPLC with UV and amperometric detection 337(1985)351

Cholecystokinin
Rapid, small-scale preparation of gastrointestinal hormones by HPLC on a C_{18} column 326(1985)121

Cholestanol
Simultaneous determination of cholestanol and cholesterol in human serum by HPLC with fluorescence detection 341(1985)432

Cholesterol
Rapid separation of plasma lipoproteins by GPC on agarose gel Superose 6B 341(1985)154

Simultaneous determination of cholestanol and cholesterol in human serum by HPLC with fluorescence detection 341(1985)432

Bile acids. LXXVI. Analyses of bile acids and sterols by HPLC with a microbore column 347(1985)155

Cholesteryl esters
Assay of lipids in dog myocardium using capillary GC and derivatization with boron trifluoride and methanol 339(1985)25

Cholesteryl ester quantitation: non-equivalent rates of transesterification of molecular species in the presence of silica gel 343(1985)143

Use of reversed-phase HPLC to analyse the cholesterol ester profiles present in lipoproteins of patients exhibiting hyperalphalipoproteinaemia or hypoalphalipoproteinaemia (Tangier disease) 343(1985)149

Cholic acid
Determination of bile acids in rat bile by HPLC 338(1985)195

Rapid and accurate reversed-phase HPLC determination of conjugated bile acids in human bile for routine clinical applications. Therapeutic control during gallstone dissolution therapy 339(1985)263

Choline
Improved analysis of acetylcholine and choline in canine brain and blood samples by capillary GC-MS 339(1985)170

Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 341(1985)23

Determination of free choline in human semen using an isotachophoretic analyser 343(1985)186

Highly sensitive assay for acetylcholinesterase activity by HPLC with electrochemical detection 344(1985)93

Use of radio GC for monitoring the *in vivo* labelling of postmorem [^{3}H]choline production 347(1985)387

HPLC method for the separation and estimation of choline, glycinebetaine aldehyde and related compounds 350(1985)245

Choline acetyltransferase
Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 341(1985)23

Cholinergics
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Cholinesterase
Affinity electrophoresis in polyacrylamide gel. Influence of the gel concentration on the apparent affinity of cholinesterase for an anionic site ligand (in French) 328(1985)135

Chorismic acid
HPLC of shikimate pathway intermediates 329(1985)268

Chromarods
Effect of copper(II) sulphate impregnation of Chromarods on the sensitivity of the Iatroscan detection system to lipids 346(1985)468

Chromatofocusing
Use of chromatofocusing for separation of β-lactamases. V. Inducible chromosomally mediated β-lactamase of the *Enterobacter cloacae* 53 strain 333(1985)239

Use of chromatofocusing for separation of β-lactamases. VI. Comparative studies on two chromatofocusing polybuffer exchanger matrices with the R46 plasmid coded β-lactamase carried by the *Escherichia coli* K12 J5-3 strain 333(1985)244

Chromatophoresis: a new approach to the theroy and practice of chromatofocusing. I. General principles 347(1985)325

Chromatogram abnormalities
Computer simulation of abnormal high-performance liquid chromatograms caused by solvents 329(1985)13

Chromatogram simulation
Computer-aided mobile phase optimization and chromatogram simulation in HPLC (Review) 338(1985)1

Chromatographic dwell
HPLC of amino acids, peptides and proteins. LXVI Investigations on the effects of chromatographic dwell in the reversed-phase HPLC separation of proteins 327(1985)47

Chromatographic-mass spectrometric data
New methods of interpretation of chromatographic-MS data 349(1985)385

Chromatographic sniffing
Stripping analysis and chromatographic sniffing for the source identification of odorous compounds in drinking water 328(1985)219

Chromatographic theory
HPLC theory for the practitioner (Review) 334(1985)197

Chromatophoresis
Chromatophoresis: a new approach to the theroy and practice of chromatofocusing. I. General principles 347(1985)325

Chromium chelates
HPLC of copper, cobalt, nickel, chromium and vanadium chelates with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol 333(1985)133

GC and LC separation of geometrical isomers of tris(2,2,7-trimethyloctane-3,5-dionato)chromium(III) and cobalt(III) chelates and other Cr-β-diketonates 349(1985)469

Chromium diphenylphosphinate
Coordination polymers as stationary phases and adsorbents for GC. VIII. An attempt to bind poly[chromium(III) diphenylphosphinate] to silica 330(1985)370

Chromosorb 103
Correlation between chemical and physical characteristics and adsorption properties of some styrene copolymers 330(1985)55

Chrysophanol
HPLC separation and quantitative determination of 1,8-dihydroxyanthraquinones in plant cell cultures 329(1985)311

Chymopapain
HPLC investigations on some enzymes of papaya latex 326(1985)103

Chymotrypsin
High-performance affintiy chromatography of anhydrochymotrypsin on a hydrophilic vinyl-polymer gel coupled with tryptophan 350(1985)265

Cianopramine
Determination of cianopramine in human plasma by HPLC and GLC with UV, fluorescence and electron-capture detection 342(1985)97

Cibenzoline
Quantitation of cibenzoline in human plasma by GC-negative-ion chemical-ionization MS 336(1985)403

Cicletanide
HPLC determination of cicletanide, a new diuretic, in plasma, red blood cells, urine and saliva 341(1985)97

Ciclopirox
Quantification of ciclopirox by HPLC after pre-column derivatization. An example of efficient clean-up using silica-bonded cyano phases 339(1985)451

Cilostazol
HPLC procedure for the determination of a new antithrombotic and vasodilating agent, cilostazol, in human plasma 338(1985)456

Cimetidine
Determination of cimetidine and metabolites in plasma by reversed-phase HPLC radial compression technique 339(1985)139

HPLC method for the simultaneous determination of cimetidine and antipyrine in plasma 345(1985)192

Cinchona alkaloids
Cinchona alkaloids for preparing new, easily accessible chiral stationary phases. II. Resolution of binaphthol derivatives on silica-supported quinine 348(1985)79

Cinnamic anhydride
Cinnamoylation of the sugar hydroxyls of 3-methylthymidine. Formation of a relatively stable ester derivative 347(1985)99

Ciprofloxacin
HPLC method for determination of ciprofloxacin in biological fluids 339(1985)347

Citalopram
Determination of citalopram, amitriptyline and clomipramine in plasma by reversed-phase HPLC 338(1985)171

Citrazinic acid
Detection of citrazinic acid by PC and TLC 331(1985)202

Citreoviridin
TLC of mycotoxins (Review) 334(1985)211

Citrinin
TLC of mycotoxins (Review) 334(1985)211

Clavulanic acid
Clavulanate-potentiated ticarcillin: HPLC assays for clavulanic acid and ticarcillin isomers in serum and urine 337(1985)301

Clenbuterol
Determination of clenbuterol in the high nanogram range in plasma of mice by HPLC with amperometric detection 336(1985)415

Clobazam
Determination of clobazam, N-desmethylclobazam and their hydroxy metabolites in plasma and urine by HPLC 343(1985)369

Clomipramine
Determination of citalopram, amitriptyline and clomipramine in plasma by reversed-phase HPLC 338(1985)171

Clonazepam
Simple HPLC assay for the routine monitoring of clonazepam in plasma 336(1985)229

Determination of antiepileptic drugs (Review) 340(1985)199

LC analysis of clonazepam in human serum with solid-phase (Bond-Elut) extraction 341(1985)383

Clonidine
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Clostridium
GC-MS analysis of volatile amines produced by several strains of *Clostridium* 337(1985)213

CN
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379

Coal derived liquids
Determination of molecular weights of coal derived liquids by GPC-HPLC 329(1985)65

Cobalt chelates
HPLC of copper, cobalt, nickel, chromium and vanadium chelates with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol 333(1985)133

GC and LC separation of geometrical isomers of tris(2,2,7-trimethyloctane-3,5-dionato)chromium(III) and cobalt(III) chelates and other Cr-β-diketonates 349(1985)469

Cobalt complexes
Phenol effect in PC of metal complexes 331(1985)273

Cobalt phthalocyanine
Determination of cysteine and glutathione in plasma and blood by LC with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 345(1985)43

Codeine
Determination of drugs in untreated body fluids by micellar chromatography with fluorescence detection 342(1985)293

Improved method for the simultaneous determination of morphine, codeine and dihydrocodeine in blood by HPLC with electrochemical detection 347(1985)398

Simultaneous HPLC stability-indicating analysis of acetaminophen and codeine phosphate in tablets and capsules 348(1985)253

Coenzyme Q
HPLC of coenzyme Q-related compounds and its application to biological materials 342(1985)35

Colchicine
Colchicine quantitation by HPLC in human plasma and urine 342(1985)416

Collagen
Comparison of reversed-phase columns for the separation of tryptic peptides by gradient elution. Correlation of experimental results and model prediction 326(1985)419

Collagenase
Highly sensitive assay for PZ-peptidase activity by HPLC 348(1985)205

Colony-stimulating factor
Purification of human urinary colony-stimulating factor by HPLC 344(1985)339

Column configuration
Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Application to method development and the choice of column configuration 327(1985)93

Column diameter
Contribution to the study of micropacked columns in GC 348(1985)327

Column efficiency
Re-packing reversed-phase HPLC columns as means of regenerating column efficiency and prolonging packing life 328(1985)347

Contribution to the study of micropacked columns in GC 348(1985)327

Effect of the mesh-size distribution of porous polymer beads on efficiency and selectivity of stationary phases 349(1985)189

Effects of density and temperature on efficiency in capillary supercritical fluid chromatography 349(1985)305

Column life-time
Column lifetime of a new agarose medium for high-performance gel filtration chromatography at basic pH 330(1985)360

Shelf-life of unused HPLC columns 346(1985)461

Column preparation
Open-tubular columns support-coated *in situ* with cross-linked SE-54 stationary phase 329(1985)123

Column re-packing
Re-packing reversed-phase HPLC columns as means of regenerating column efficiency and prolonging packing life 328(1985)347

Columns
see also Capillary columns and microbore columns

Shelf-life of unused HPLC columns 346(1985)461

Contribution to the study of micropacked columns in GC 348(1985)327

Column storage
Effects of solutions used for storage of size-exclusion columns on subsequent chromatography of peptides and proteins 331(1985)253

Column temperature
Effect of column temperature on HPLC behaviour of inorganic polyphosphates. I.

Isocratic ion-exchange chromatography 348(1985)27

Effect of column temperature on HPLC behaviour of inorganic polyphosphates. II. Gradient ion-exchange chromatography 350(1985)119

Combustion tube
Application of radioisotope tracer techniques to analytical GC: determination of GC peak yield 341(1985)251

Commodore 64
Versatile HPLC computer system for solvent delivery, gradient control and data acquisition using the Commodore 64 personal computer 333(1985)301

Complement component C4b-binding protein
Application of DNA-cellulose chromatography in the isolation of immunoglobulin M and complement component C4b-binding protein from human serum 347(1985)129

Complexing agents
Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147

Computer-assisted chromatography
Versatile HPLC computer system for solvent delivery, gradient control and data acquisition using the Commodore 64 personal computer 333(1985)301

Computer-aided mobile phase optimization and chromatogram simulation in HPLC (Review) 338(1985)1

Microcomputer programming in BASIC for the evaluation of capillary GC in the analysis of pesticide residues. I. MATRIXCOMP: a program that facilitates the recognition of interfering peaks from the biological matrix 349(1985)49

Computer-assisted prediction of retention times for inorganic polyphosphates in gradient ion-exchange chromatography 350(1985)461

HPLC determination of contaminants in salicylic acids after computer-assisted optimization of the separation parameters (in German) 350(1985)471

Computer simulation
Computer simulation of abnormal high-performance liquid chromatograms caused by solvents 329(1985)13

Concentration scales
Possibility and consequences of using different concentration scales in the study of solution thermodynamics by GLC 350(1985)339

Connectivity indices
Relationships between GLC behaviour and structure of arylaliphatic acids 333(1985)21

Contact potentials
Effects of contact potentials on the pulsed electron-capture detector 349(1985)155

Continuous flow extraction
Automated monitoring of organic trace components in water. I. Continuous flow extraction together with on-line capillary GC 330(1985)263

Continuous-flow nuclear magnetic resonance
Continuous-flow nuclear magnetic resonance 346(1985)17

Controlled-pore glasses
Structure parameters of molecules and media evaluated by chromatographic partition. I. Controlled-pore glasses 350(1985)1

Cooling plate
Cooling plate for cellulose thin-layer electrophoresis and its application to amino acid analysis 347(1985)199

Coordination polymers
Coordination polymers as stationary phases and adsorbents for GC. VIII. An attempt to bind poly[chromium(III) diphenylphosphinate] to silica 330(1985)370

Copolyamides
GC analysis of isocyanate-based copolyamide resins 329(1985)132

Copper chelates
HPLC of copper, cobalt, nickel, chromium and vanadium chelates with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol 333(1985)133

Copper sulphate
Effect of copper(II) sulphate impregnation of Chromarods on the sensitivity of the Iatroscan detection system to lipids 346(1985)468

Coproporphyrin
Ion-pair HPLC separation of porphyrin isomers 349(1985)283

Coronavirus glycoproteins
Isolation of the subunits of the coronavirus envelope glycoprotein E2 by hydroxyapatite HPLC 326(1985)191

Corticosteroids
Efficient chromatographic fractionation of steroids in human serum through regulation of liquid-liquid distribution ratios 339(1985)253

Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Quantitative determination of betamethasone and its major metabolite in equine urine by micro-LC-MS 341(1985)261

Cortisol
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Cortisol-3-(O-carboxymethyl)oxime tyrosine methyl ester
Adsorption chromatographic separation of ^{125}I-labelled cortisol-3-(O-carboxymethyl)oxime tyrosine methyl ester 329(1985)264

Corynanthine
Sub-nanogram analysis of yohimbine and related compounds by HPLC 342(1985)333

Cotinine
Determination of cotinine in biological fluids by capillary GC-MS selected-ion monitoring 342(1985)79

Rapid GC method for the determination of nicotine and cotinine in urine (in French) 343(1985)424

Simple HPLC method for rapid determination of nicotine and cotinine in urine 344(1985)391

Coulometric detector
Low dead volume coulometric detector for LC 332(1985)117

Coumermycin A_1
Determination of coumermycin A_1 in plasma by reversed-phase HPLC analysis 342(1985)145

Counter-current chromatography
Large-scale preparative countercurrent chromatography for separation of polar compounds 348(1985)131

Counter-current distribution
Direct chiral resolution in an aqueous two-phase system using the counter-current distribution principle 333(1985)211

Counter-ion effects
Counter-ion effects in partition chromatography 332(1985)203

C-peptide
New application of HPLC for analysis of urinary C-peptide 338(1985)71

CR
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379

Creatine phosphate
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides 337(1985)201

HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides. Reply to E. Harmsen et al 337(1985)203

Creatinine
Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251

Crepenynic acid
Semi-preparative isolation of crepenynic acid, a potential inhibitor of essential fatty acid metabolism 346(1985)431

***p*-Cresol**
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

Cresols
α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131

Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

Cromoglycate
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Crown ethers
LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

CS
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379

Cuticular hydrocarbons
Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. I. Introduction and characterization of the cuticular hydrocarbon patterns of *Solenopsis invicta* and *S. richteri* 349(1985)31

Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. II. Comparison of the cuticular hydrocarbon patterns between different colonies of *Solenopsis richteri* 349(1985)39

Cyanogen bromide digests
General method for the separation of cyanogen bromide digests of proteins by HPLC. Rabbit skeletal troponin I 326(1985)349

Cyclic nucleotide-metal ion complexes
Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. I. Cation binding in the mobile phase 329(1985)321

Cyclic nucleotides
Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. I. Cation binding in the mobile phase 329(1985)321

Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. II. Quantitative determination of hydrophobicity 350(1985)105

Cyclobenzaprine
Determination of cyclobenzaprine in plasma and urine using capillary GC with nitrogen-selective detection 339(1985)414

Cyclodextrins
Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

Methylated cyclodextrins as mobile phase additives in LC 330(1985)253

Electrokinetic chromatography with 2-O-carboxymethyl-β-cyclodextrin as a moving "stationary" phase 332(1985)211

Microanalyses of β-cyclodextrin in plasma by HPLC 341(1985)31

LC retention behavior of organometallic compounds and ligands with amine-, octadecylsilica- and β-cyclodextrin-bonded phase columns 347(1985)51

Resolution of *cis* - and *trans* - dimethylcyclohexanes by partition GC through cyclodextrin complexes 349(1985)3

α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131

Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

HPLC separation of aromatic carboxylic acids with β-cyclodextrin-bonded stationary phases 350(1985)37

Cyclohexylalkanes
Kováts retention indices of high-molecular-weight monomethyl-, cyclopentyl-, cyclohexyl- and phenylalkanes 331(1985)219

Cyclohexylphenylacetic acid
Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59

Cyclohexylphenylglycolic acid
Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59

Cyclopentenyl fatty acids
HPLC of triglycerides of Flacourtiaceae seed oils containing cyclopentenyl fatty acids (chaulmoogric oils) 348(1985)441

Cyclopentylalkanes
Kováts retention indices of high-molecular-weight monomethyl-, cyclopentyl-, cyclohexyl- and phenylalkanes 331(1985)219

Cyclophosphamide
Drug level monitoring: cytostatics (Review) 340(1985)31

α-Cyclopiazonic acid
TLC of mycotoxins (Review) 334(1985)211

Cycloserine
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Cyclosporins
Investigation of the mechanism of peak broadening observed in the HPLC analysis of cyclosporine 333(1985)231

Simplified LC analysis for cyclosporin A in blood and plasma with use of rapid extraction 337(1985)416

Semi-automated HPLC determination of cyclosporine A in whole blood using one-step sample purification and column-switching 341(1985)411

Modified column-switching HPLC method for the measurement of cyclosporine in serum 344(1985)422

Measurement of cyclosporin by HPLC following charcoal adsorption from whole blood 344(1985)423

Cyproheptadine
HPLC method for quantification of cyproheptadine in serum or plasma 339(1985)457

Cyproterone acetate
15-Hydroxycyproterone acetate and cyproterone acetate levels in plasma and urine 338(1985)174

Cysteic acid
Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1

Cysteine
Rapid and simple method for quantitative determination of non-protein sulphydryls in mouse liver by reversed-phase HPLC 338(1985)209

Determination of cysteine and glutathione in plasma and blood by LC with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 345(1985)43

Cysteinesulphinate
Determination of cysteinesulfinate, hypotaurine and taurine in physiological samples by reversed-phase HPLC 343(1985)303

Cytochalasans
TLC of mycotoxins (Review) 334(1985)211

Cytochalasins
TLC of mycotoxins (Review) 334(1985)211

Cytochrome *c*
Comparison of reversed-phase columns for the separation of tryptic peptides by gradient elution. Correlation of experimental results and model prediction 326(1985)419

Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

Cytochrome P-450
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Cytokines
HPLC separation of distinct epidermal cell-derived cytokines 326(1985)129

Cytosine
HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33

Cytosine arabinoside
Simultaneous determination of cytosine arabinoside, its nucleotides and metabolites by ion-pair HPLC 337(1985)63

Cytosine arabinosine
Drug level monitoring: cytostatics (Review) 340(1985)31

Cytostatics
Drug level monitoring: cytostatics (Review) 340(1985)31

D

Daidzein
Analysis of isoflavones in *Puerariae radix* by HPLC with amperometric detection 347(1985)438

Daidzin
Analysis of isoflavones in *Puerariae radix* by HPLC with amperometric detection 347(1985)438

Dammarane-type triterpene alcohols
Preparation and GLC of trimethylsilyl derivatives of dammarane-type triterpene triols and tetrols 329(1985)189

Dansylamides
Properties of six octadecylsilane stationary phases for the separation of dansylamides 330(1985)243

Danthron
Analysis of anthralin in dermatological products by reversed-phase HPLC 328(1985)317

Dapsone
Rapid and sensitive LC method for the determination of dapsone and monoacetyldapsone in plasma and urine 345(1985)447

Daunomycin
Preparative chromatography of epimers and anomers of daunomycin derivatives on Sephadex LH-20 329(1985)193

DDT
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Dead time
Estimation of dead time and calculation of Kováts indices (Review) 334(1985)95

Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 345(1985)457

Dead volume
HPLC theory for the practitioner (Review) 334(1985)197

Theory of solvent disturbance peaks and experimental determination of thermodynamic dead-volume in column LC 349(1985)211

3-Deazauridine
Analysis of candidate anticancer drugs by thermospray HPLC-MS and by high-resolution MS 347(1985)137

Decapreno-β-carotene
Decapreno-β-carotene as an internal standard for the quantification of the hydrocarbon carotenoids by HPLC 346(1985)237

Deconvolution methods
Quantitative deconvolution of heavily fused chromatographic peaks of biological components using a multi-wavelength UV detector 332(1985)139

Deconvolution of overlapping chromatograms. II. Analytical solution to the moments of the GEX function 333(1985)152

Dehydroascorbic acid
Separation of ascorbic acid, dehydroascorbic acid, diketogulonic acid and glucose by isocratic elution from a column of a hydrophilic gel 332(1985)283

Dehydroepiandrosterone
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 341(1985)426

3-Dehydroquinic acid
HPLC of shikimate pathway intermediates 329(1985)268

3-Dehydroshikimic acid
HPLC of shikimate pathway intermediates 329(1985)268

Deltamethrin
High-performance liquid chromatograph coupled with two detectors: a UV spectrometer and a polarimeter. Example in the field of pyrethroids: identification of enantiomers 349(1985)109

Demethoxybuprenorphine
^{63}Ni electron-capture GC assay for buprenorphine and metabolites in human urine and feces 337(1985)291

Denaturation
Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

Density effects
Effects of density and temperature on efficiency in capillary supercritical fluid chromatography 349(1985)305

Density programming
Effects of density and temperature on efficiency in capillary supercritical fluid chromatography 349(1985)305

Deoxaphomin
TLC of mycotoxins (Review) 334(1985)211

6-Deoxyallitol
Identification of 6-deoxyallitol and 6-deoxygulitol in human urine. Electron-impact mass spectra of eight isomers of 6-deoxyhexitol 336(1985)345

Deoxycholic acid
Determination of bile acids in rat bile by HPLC 338(1985)195

Rapid and accurate reversed-phase HPLC determination of conjugated bile acids in human bile for routine clinical applications. Therapeutic control during gallstone dissolution therapy 339(1985)263

2-Deoxy-2,3-didehydro-N-acetylneuraminic acid
Structural analysis of underivatized sialic acids by combined HPLC-MS 337(1985)231

4'-Deoxydoxorubicin
Method for the determination of 4'-deoxydoxorubicin, 4'-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 341(1985)401

4'-Deoxydoxorubicinol
Method for the determination of 4'-deoxydoxorubicin, 4'-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 341(1985)401

5'-Deoxy-5-fluorouridine
Analysis of 5'-deoxy-5-fluorouridine and 5-fluorouracil in human plasma and urine by HPLC 342(1985)303

α-2-Deoxy-D-glucose
Determination of α-2-deoxy-D-glucose in topical formulations by HPLC with UV detection 331(1985)183

6-Deoxygulitol
Identification of 6-deoxyallitol and 6-deoxygulitol in human urine. Electron-impact mass spectra of eight isomers of 6-deoxyhexitol 336(1985)345

6-Deoxyhexitol
Identification of 6-deoxyallitol and 6-deoxygulitol in human urine. Electron-impact mass spectra of eight isomers of 6-deoxyhexitol 336(1985)345

11-Deoxyjervine
Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

Deoxynivalenol
Rapid and sensitive determination of zearalenone in cereals by HPLC with fluorescence detection 328(1985)271

Analysis of some trichothecene mycotoxins by LC 347(1985)429

Deoxypyrimidine 5'-nucleotidase
Assay of human erythrocyte pyrimidine and deoxypyrimidine 5'-nucleotidase by isocratic reversed-phase HPLC 339(1985)293

Deoxyribonucleosides
Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33

HPLC method for the analysis of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in biological fluids 339(1985)380

Deoxysepiapterin
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

Derivative spectroscopy
Application of derivative spectroscopy to the determination of chromatographic peak purity 347(1985)219

Derivatization
Determination of thiamine and its phosphate esters in human and rat blood by HPLC with post-column derivatization 332(1985)181

Derivatization, electropheresis
Paper electrophoretic characterization of ethylenic compounds as dimethylsulphoniomethylthio derivatives 328(1985)436

Derivatization, GC
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Determination of optical purity by enantioselective capillary GC: panthenol and related compounds 328(1985)357

Preparation and GLC of trimethylsilyl derivatives of dammarane-type triterpene triols and tetrols 329(1985)189

Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Evaluation of a rapid method for preparation of fatty acid methyl esters for analysis by GLC 331(1985)141

Catalysed derivatisation of trimethylsilyl ethers of ecdysterone. A preliminary study 333(1985)186

Rapid and selective derivatization method for the nitrogen-selective detection of carboxylic acids in biological fluids prior to GC analysis 336(1985)241

Automated theophylline assay using GC and a mass-selective detector 336(1985)285

Quantitation of cibenzoline in human plasma by GC-negative-ion chemical-ionization MS 336(1985)403

Absolute configuration of 3-hydroxyadipic acid in human urine 337(1985)9

Rapid GC-MS quantitation of γ-aminobutyric acid in biological specimens 337(1985)21

GC determination of mefloquine in human and dog plasma using electron-capture detection 337(1985)47

One-step extraction procedure for GC determination of viloxazine as its acetyl derivative in human plasma 337(1985)142

^{63}Ni electron-capture GC assay for buprenorphine and metabolites in human urine and feces 337(1985)291

N-Methylation and N,N-dimethylation of amino acids. An artifact production in the analysis of organic acids using diazomethane as derivatizing agent 338(1985)33

Quantitative analysis of veralipride in plasma and urine by GC-MS and GC with flame-ionization detection 338(1985)79

Analysis of buprenorphine and its N-dealkylated metabolite in plasma and urine by selected-ion monitoring 338(1985)89

GC determination of serum branched-chain α-keto acids derivatized by extractive alkylation 338(1985)265

GC-MS of epimeric 19-norandrostan-3-ol-17-ones as the trimethylsilyl ether, methyloxime-trimethylsilyl ether and trimethylsilyl-enol trimethylsilyl ether derivatives 339(1985)11

Assay of lipids in dog myocardium using capillary GC and derivatization with boron trifluoride and methanol 339(1985)25

Quantitative determination of tertatolol in biological fluids by GC-MS 339(1985)97

Analysis of epoxyeicosatrienoic acids by GC-MS using chlorohydrin adducts 339(1985)175

Determination of plasma testosterone by mass fragmentography using [3,4-^{13}C]testosterone as an internal standard 339(1985)233

Frequency-pulsed electron-capture GLC studies of chemical changes in sera of patients with schistosomiasis 339(1985)243

Determination of acetyltrimoprostil and its metabolite trimoprostil in human or dog plasma by GC-negative-ion chemical ionization MS 339(1985)313

GC determination of *m*- and *p*-hydroxyphenytoin in the urine of epileptic patients 339(1985)331

Electron-capture GC of taurine as its N-pentafluorobenzoyl di-*n*-butylamide derivative 339(1985)370

Simultaneous determination of metoprolol and deuterium-labelled metoprolol in human plasma by GC-negative-ion MS 339(1985)404

Drug level monitoring: Sedative hypnotics (Review) 340(1985)139

Determination of antiepileptic drugs (Review) 340(1985)199

Antibiotic monitoring in body fluids (Review) 340(1985)361

Identification of metabolites diagnostic for organic acidurias by simultaneous dual-column capillary GC 341(1985)1

Determination of meprobamate as an *n*-butylboronate ester derivative in serum by GLC 341(1985)462

Differentiation between major species of the *Actinobacillus-Haemophilis-Pasteurella* group by GC of trifluoroacetic acid anhydride derivatives from whole-cell methanolysates 342(1985)13

Improved direct injection method and extractive methylation method for determination of valproic acid in serum by GC 342(1985)89

Identification and quantification of 1-carboxysalsolinol and salsolinol in biological samples by GC-MS 343(1985)1

Highly sensitive determination of 5-fluorouracil in human plasma by capillary GC and negative ion chemical ionization MS 343(1985)59

Cholesteryl ester quantitation: non-equivalent rates of transesterification of molecular

species in the presence of silica gel 343(1985)143

GC determination of pyrazoloimidazole 343(1985)190

Determination of dexamethasone in human plasma and urine by electron-impact MS 343(1985)231

Determination of low plasma concentrations of 3-methoxy-4-hydroxyphenylethylene glycol using GC-negative-ion chemical ionization MS 343(1985)239

Measurement of imidazoleacetic acid in urine by GC-MS 343(1985)249

GC-MS analysis of methylphenidate and *p*-hydroxymethylphenidate using deuterated internal standards 343(1985)329

GC-MS of carboxylic acids in tissues as their *tert.*-butyldimethylsilyl derivatives 344(1985)1

Determination of di(2-ethylhexyl) phthalate and four of its metabolites in blood plasma by GC-MS 344(1985)167

Determination of oxpentifylline and four metabolites in plasma and urine by automated capillary GC using nitrogen-selective detection 344(1985)187

Specific and sensitive determination of medroxyprogesterone acetate in human serum by GC-MS 345(1985)1

Simultaneous determination of metapramine and its dimethylated metabolites in plasma by GC-MS 345(1985)59

GC assay for oxcarbazepine and its main metabolites in plasma 345(1985)67

Determination of SQ 27,519, the active phosphinic acid-carboxylic acid of the prodrug SQ 28,555, in human serum by capillary GC with nitrogen-phosphorus detection after a two-step derivatization 345(1985)299

GLC determination of isosorbide-5-mononitrate derivatized with perfluorinated anhydrides for pharmacokinetic and bioavailability investigations 345(1985)390

Improved solvent-extraction based procedure for the GC analysis of resin and fatty acids in pulp mill effluents 346(1985)205

Pentafluorobenzyl *p*-toluenesulphonate as a new derivatizing reagent for GC determination of anions 346(1985)215

Modifications in the alditol acetate method for analysis of muramic acid and other neutral and amino sugars by capillary GC-MS with selected ion monitoring 347(1985)109

Determination of ethyl phosphite, phosphite and phosphate in plant tissues by anion-exchange HPLC and GC 347(1985)267

Analytical separation of enantiomeric polygodials by GC of pyrrole derivatives 347(1985)455

Silica capillary GC of prostaglandins with electron-capture detection and its application to the forensic investigation of sexual offences 348(1985)111

Preparation of trimethylsilyl derivatives of ribonucleosides for GC 348(1985)280

Direct derivatization of alprenolol and its 4-hydroxy metabolite in urine with phosgene and methanol prior to analysis by capillary column GC 349(1985)447

GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Capillary GC of partially methylated alditol acetates on a SP-2100 wall-coated open-tubular column 350(1985)304

Enantiomer separation of metoprolol and its analogues and metabolites by capillary column GC after derivatization with phosgene 350(1985)328

Derivatization, LC

Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1

Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

High-speed LC analysis of amino acids by post-column sodium hypochlorite-*o*-phthalaldehyde reaction 327(1985)17

Improved method for the determination of N-acetylcysteine in human plasma by HPLC 327(1985)261

4-methyl-7-methoxycoumarin as a fluorescent label for HPLC analysis of dicarboxylic acids 328(1985)111

GLC determination of free amino acids in fermentation broth during growth of *Clostridium oncolyticum* M 55 328(1985)241

High-performance size-exclusion chromatography of lignin and its derivatives 328(1985)299

Determination of anthraquinones in pulping materials by HPLC using online post-column derivatization 328(1985)309

Direct diastereomeric resolution of carotenoids. II. All ten stereoisomers of tunaxanthin (ε,ε-carotene-3,3'-diol) 328(1985)387

Adsorption and reversed-phase HPLC of *p*-nitrobenzyl esters of monohydroxy fatty acids 329(1985)57

Determination of low concentrations of low-molecularweight aldehydes and ketones in aqueous samples 329(1985)81

Determination of catecholamines by HPLC with catalytic photometric detection 329(1985)147

A method for the determination of S-methylcysteine by HPLC: application to the study of carcinogenic methylating agents 329(1985)202

Simultaneous determination of iron(II) and -(III) by ion chromatography with post-column reaction 329(1985)247

Off-line LC-MS studies of fluorescent β-aminothiol-*o*-phthalaldehyde derivatives 330(1985)307

LC with pre-column dansyl derivatization and fluorimetric detection applied to the assay of morphine in biological samples 330(1985)323

Simultaneous determination of abscisic acid and jasmonic acid in plant extracts using HPLC 330(1985)347

Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109

Formation and separation of fluorescent derivatives of diacylglycerols 331(1985)177

HPLC of fatty acids as their *p*-phenylazophenacyl esters 331(1985)187

Determination of the enantiomeric composition of chiral epoxides using chiral derivatization and LC 331(1985)349

Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Measurement of neutral sugars in glycoproteins as dansyl derivatives by automated HPLC 333(1985)123

Reversed-phase ion-pair chromatography of some biologically important carboxylic acids as 2-nitrophenylhydrazides 333(1985)215

Ion-pair HPLC determination of dansylated aliphatic polyamines with fluorescence and UV detection 333(1985)220

Determination of formaldehyde as its lutidine derivative in the atmospheric liquid phase by HPLC 333(1985)262

Determination of aldehydes and ketones by derivatization and LC-MS 333(1985)337

Derivatization of alkyl halides and epoxides with picric acid salts for improved HPLC detection 333(1985)349

Derivatization of secondary amines with 2-naphthalenesulfonyl chloride for HPLC analysis of spectinomycin 333(1985)365

Rapid analysis of amino acids using pre-column derivatization 336(1985)93

High-sensitivity phenylthiohydantoin amino acid analysis using conventional and microbore chromatography 336(1985)105

HPLC separation of fatty acids as pentafluorobenzyl esters 336(1985)115

Determination of carnitine, butyrobetaine, and betaine as 4'-bromophenacyl ester derivatives by HPLC 336(1985)271

HPLC assay for pilocarpine in aqueous humor 336(1985)337

Reversed-phase HPLC method for determination of brain glutamate decarboxylase suitable for use in kinetic studies 336(1985)385

Quantification of branched-chain α-keto acids as quinoxalinols: importance of excluding oxygen during derivatization 337(1985)98

HPLC assay for mexiletine enantiomers in human plasma 337(1985)172

Analysis of aminoglycoside antibiotics as benzoyl derivatives by HPLC and its application to the quantitation of neomycin in the perilymph 337(1985)187

HPLC analysis of baclofen in plasma and urine of man after precolumn extraction and derivatization with *o*-phthaldialdehyde 337(1985)341

Determination of neurotransmitter amino acids by HPLC with fluorescence detection 337(1985)391

Determination of isoniazid, actylisoniazid, acetylhydrazine and diacetylhydrazine in biological fluids by HPLC 338(1985)113

Determination of the antibiotic fludalanine in plasma and urine by HPLC using a packed-bed, post-column reactor with *o*-phthalaldehyde and 2-mercaptoethanol 338(1985)357

Simplified procedure for specific determination of imino acids in human blood plasma by HPLC 339(1985)59

HPLC analysis of free hydroxyproline and proline in blood plasma and of free and peptide-bound hydroxyproline in urine 339(1985)285

Determination of the enantiomers of indoprofen in blood plasma by HPLC after rapid derivatization by means of ethyl chloroformate 339(1985)339

HPLC analysis of melphalan in plasma, brain and peripheral tissue by *o*-phthalaldehyde

derivatization and fluorescence detection 339(1985)434

Drug level monitoring: Sedative hypnotics (Review) 340(1985)139

Antibiotic monitoring in body fluids (Review) 340(1985)361

Determination of amino acids by separation of their ion pairs with dodecyl sulphate 341(1985)11

HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 341(1985)115

Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 341(1985)213

Derivatization in aqueous solution, isolation and separation of tetrahydro-β-carbolines and their precursors by LC 341(1985)287

Determination of diclofensine, an antidepressant agent, and its major metabolites in human plasma by HPLC with fluorometric detection 341(1985)313

3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 341(1985)426

Simultaneous determination of cholestanol and cholesterol in human serum by HPLC with fluorescence detection 341(1985)432

Quantitative method for the detection of glucose in body fluids by HPLC with fluorescence detection 342(1985)179

Determination of dopamine-3- and -4-O-sulphate in human plasma and urine by anion-exchange HPLC with fluorimetric detection 342(1985)261

HPLC analysis of guanidino compounds using ninhydrin reagent. II. Guanidino compounds in blood of patients on haemodialysis therapy 342(1985)269

Determination of plasma tranexamic acid using cation-exchange HPLC with fluorescence detection 343(1985)109

Determination of (15*R*)- and (15*S*)-15-methylprostaglandin E_2 in human plasma with picogram per milliliter sensitivity by column-switching HPLC 343(1985)271

Determination of α-keto acids in serum and urine by HPLC with fluorescence detection 344(1985)33

HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61

HPLC determination of plasma and brain histamine without previous purification of biological samples: cation-exchange chromatography coupled with post-column derivatization fluorometry 344(1985)115

LC determination of mexiletine and tocainide in human plasma with fluorescence detection after reaction with a modified *o*-phthalaldehyde reagent 344(1985)221

Determination of *p*-hydroxybestatin in human serum by HPLC using fluorescence detection 344(1985)267

Automated determination of amoxycillin in biological fluids by column switching in ion-pair reversed-phase LC systems with post-column derivatization 344(1985)285

HPLC with fluorimetric detection in biological tissues of the 4-bromomethyl-7-methoxycoumarin ester derivative of 5-pyrrolidone-2-carboxylic acid 344(1985)325

Determination of total ascorbic acid in human serum by HPLC with fluorescence detection 344(1985)351

Fluorimetric determination of the quaternary compound trospium and its metabolite in biological material after derivatization with benoxaprofen chloride 345(1985)99

Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 345(1985)229

HPLC method for monitoring leupeptin in mouse serum and muscle by pre-column fluorescence derivatization with benzoin 345(1985)259

Capillary GC method for the determination of complex mixture of isocyanates and amines 346(1985)191

3-Bromomethyl-6,7-dimethoxy-1-methyl-2(1*H*)-quinoxalinone as a new fluorescence derivatization reagent for carboxylic acids in HPLC 346(1985)227

Cinnamoylation of the sugar hydroxyls of 3-methylthymidine. Formation of a relatively stable ester derivative 347(1985)99

Selective determination of secondary amino acids using precolumn derivatization with 9-fluorenylmethylchloroformate and reversed-phase HPLC 348(1985)213

Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Rapid analysis of the major classes of retinoids by step gradient reversed-phase HPLC using retinal (O-ethyl) oxime derivatives 348(1985)241

Determination of phenylethanolamine N-methyltransferase by HPLC with fluorescence detection 348(1985)407

Fluorescent labelling of amino acids with 9-anthryldiazomethane and its applications to HPLC 348(1985)425

HPLC determination of naphthols as 4-aminoantipyrine derivatives. Application to carbaryl 349(1985)63

Complete HPLC separation of 4-N,N-dimethylaminoazobenzene-4'-thiohydantoin and 4-dimethylaminoazobenzene-4'-sulphonyl chloride amino acids utilizing the same reversed-phase column at room temperature 349(1985)77

Separation of phenylthiohydantoin-amino acids by HPLC and some applications in dansyl Edman sequence analysis 349(1985)99

Determination of trace amounts of lanthanides in rocks and minerals by HPLC 349(1985)137

HPLC separation and determination of prostaglandins, oxidized by pyridinium dichromate. Optimization and applications 349(1985)431

HPLC method for the separation and estimation of choline, glycinebetaine aldehyde and related compounds 350(1985)245

HPLC of anthranilate synthase using gel filtration and a post-column reactor 350(1985)332

Determination of phenols in water and wastewater by post-column reaction detection HPLC 350(1985)407

Diastereomeric separations of natural glycero derivatives as their 1-(1-naphthyl)ethyl carbamates by HPLC 350(1985)417

Determination of phenylthiocarbamyl amino acids by reversed-phase HPLC 350(1985)453

Derivatization, TLC
Determination of salbutamol in human plasma and urine by HPTLC 345(1985)111

Desacetylcefotaxime
Micromethod for the determination of cefotaxime and desacetylcefotaxime in plasma and urine by HPLC 336(1985)438

Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 341(1985)131

Desacetylcentapicrin
Analytical and preparative reversed-phase LC of secoiridoid glycosides 346(1985)396

Desacetyldiltiazem
Diltiazem and desacetyldiltiazem analysis in human plasma using HPLC: improved sensitivity without derivatization 342(1985)216

Desamido-insulin
Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27

Descarboxyloflazepate
Determination of circulating ethyl loflazepate metabolites in the baboon by radio-HPLC with injection of crude plasma samples: comparison with solvent extraction and TLC 342(1985)159

Desferoxamine
Determination of desferoxamine and a major metabolite by HPLC. Application to the treatment of aluminium-related disorders 341(1985)123

Desipramine
HPLC determination of imipramine and desipramine in human serum 336(1985)410

Desisopropyl indecainide
LC determination of indecainide, a new antiarrhythmic drug, and its major metabolite, desisopropyl indecainide, in biological samples 337(1985)329

N-Desisopropylpropranolol
Rapid and simultaneous extraction of propranolol, its neutral and basic metabolites from plasma and assay by HPLC 343(1985)349

N-Desmethylclobazam
Determination of clobazam, N-desmethylclobazam and their hydroxy metabolites in plasma and urine by HPLC 343(1985)369

Desmetryn
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Detection limit
HPLC theory for the practitioner (Review) 334(1985)197

Detector cell
Performance of some cell designs for laser-induced fluorescence detection in open-tubular LC 346(1985)149

Detector response factors
Use of dual-column fused-silica capillary GC in combination with detector response factors for analytical toxicology 341(1985)81

Detectors, electrophoresis
Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193

Detectors, GC
Hypercoulometry in a pulsed conventional electron-capture detector 330(1985)87

Temperature stabilization of a commercial photoionization detector. Performance evaluation 330(1985)365

Long pathlength UV absorbance detection for GC with concurrent fluorescence measurement 331(1985)77

Theory of anion contributions to non-linear electron-capture detector response 346(1985)1

GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

Effects of contact potentials on the pulsed electron-capture detector 349(1985)155

Selectivity of the redox chemiluminescence detector for complex sample analysis 349(1985)395

Optimization of helium ionization detector 349(1985)405

Integration of the deformed peak from the helium ionization detector 349(1985)415

Detector signal
HPLC theory for the practitioner (Review) 334(1985)197

Detectors, isotachophoresis
Automation of analytical isotachophoresis (Review) 334(1985)83

Multichannel spectrophotometric detector for fused-silica capillary tube isotachophoresis 346(1985)167

Isotachophoretic control of peptide synthesis and purification. A novel approach using UV detection at 206 nm 346(1985)271

Detectors, LC
Selection of mobile phases for reversed-phase LC with on-line electron-capture detection 328(1985)93

Lanthanide ions as luminescent chromophores for LC detection 328(1985)101

Thin-layer polarographic detector for the HPLC detection of thiourea derivatives 328(1985)127

High-performance UV absorption detector for LC. I. Preliminary experiments 329(1985)25

Electrochemiluminescence as a detection technique for reversed-phase HPLC. II. Low-frequency a.c. electrochemiluminescence 329(1985)281

Low dead volume coulometric detector for LC 332(1985)117

The Abbé refractometer as a LC detector using an image sensor for boundary location 332(1985)195

Recent progress in gel packing materials and detectors for modern LC in Japan 332(1985)227

Radioactivity detection system with a CaF_2 (Eu) scintillator for HPLC 333(1985)83

Nitrogen-sensitive thermionic detection in microcolumn LC 333(1985)291

Improvement of ion chromatography with UV photometric detection and comparison with conductivity detection for the determination of serum cations 341(1985)53

Photodiode array UV spectrophotometric profiling of carboxylic acids in physiological fluids 344(1985)23

Determination of cysteine and glutathione in plasma and blood by LC with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 345(1985)43

Evaluation of combined electrochemical and UV detection in the HPLC analysis of plasma for low-molecular-weight constituents 345(1985)125

Continuous-flow nuclear magnetic resonance 346(1985)17

Performance of some cell designs for laser-induced fluorescence detection in open-tubular LC 346(1985)149

Narrow-bore normal phase LC with on-line electron-capture detection 347(1985)75

Spectral properties of trifluoroacetic acid-acetonitrile gradient systems for separation of picomole quantities of peptides by reversed-phase HPLC 347(1985)83

Expert system for pharmaceutical analysis. I. Selection of the detection system in HPLC analysis: UV *versus* amperometric detection 348(1985)97

LC with voltammetric detection for quantitation of phenolic acids 348(1985)397

Coupling a high-performance liquid chromatograph with a liquid scientillation detector: example in the field of pyrethroids 349(1985)105

High-performance liquid chromatograph coupled with two detectors: a UV spectrometer and a polarimeter. Example in the field of pyrethroids: identification of enantiomers 349(1985)109

Amperometric detector for use with small-bore LC columns 350(1985)169

HPLC method utilising single or multi-wavelength detection for the comparison of disperse dyes extracted from polyester fibres 350(1985)205

Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285

Detector time constant
HPLC theory for the practitioner (Review) 334(1985)197

Detector volume
HPLC theory for the practitioner (Review) 334(1985)197

Dexamethasone
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Determination of dexamethasone in human plasma and urine by electron-impact MS 343(1985)231

Dexpanthenol
Determination of optical purity by enantioselective capillary GC: panthenol and related compounds 328(1985)357

Dextrans
GPC with interference refractometry for the rapid assay of polydisperse dextrans in biological fluids 337(1985)29

Confidence limits for a gel-permeation system 341(1985)420

Dextromethorphan
Determination of dextromethorphan and metabolites in human plasma and urine by HPLC with fluorescence detection 338(1985)99

Dextropropoxyphene
HPLC method for the determination of dextropropoxyphene and nordextropropoxyphene in serum 345(1985)413

Diacetoxyscirpenol
Analysis of some trichothecene mycotoxins by LC 347(1985)429

Diacetylhydrazine
Determination of isoniazid, actylisoniazid, acetylhydrazine and diacetylhydrazine in biological fluids by HPLC 338(1985)113

Diacylglycerols
Formation and separation of fluorescent derivatives of diacylglycerols 331(1985)177

Highly sensitive diastereoisomeric analysis of secondary alcohols and short-chain 1,2-diacylglycerols by capillary GC-MS 331(1985)295

Diastereomeric separations of natural glycero derivatives as their 1-(1-naphthyl)ethyl carbamates by HPLC 350(1985)417

Dialkyldithiocarbamates
Reversed-phase ion-interaction chromatography of sodium dialkyldithiocarbamates in the presence of tetraalkylammonium salts 348(1985)389

Dialkylglycerols
Diastereomeric separations of natural glycero derivatives as their 1-(1-naphthyl)ethyl carbamates by HPLC 350(1985)417

Dialkyl sulphides
Ligand-exchange GC of dialkyl sulphides 350(1985)63

Diamines
Frequency-pulsed electron-capture GLC studies of chemical changes in sera of patients with schistosomiasis 339(1985)243

2,3-Diaminonaphthalene
LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

1,8-Diaminonaphthalene
LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

Diaminopimelic acid
Simple method for the determination of diaminopimelic acid in rumen liquor hydrolysates 329(1985)199

***cis*-Diamminedichloroplatinum**
Drug level monitoring: cytostatics (Review) 340(1985)31

Dianthrone
Analysis of anthralin in dermatological products by reversed-phase HPLC 328(1985)317

Diatomic gases
Noble, diatomic and aliphatic gas analysis by aqueous HPLC 346(1985)93

Diazomethane
N-Methylation and N,N-dimethylation of amino acids. An artifact production in the analysis of organic acids using diazomethane as derivatizing agent 338(1985)33

Dibenzo[*b,f*]thiepins
Relationships between the chromatographic behaviour and structure of some substituted dibenzo[*b,f*]thiepins and their analogues 329(1985)113

Dibenz[*b,f*]-1,4-oxazepine
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379

Dibrohexamidine
Determination of preservatives in cosmetic products by ion-pair reversed-phase HPLC. III 348(1985)321

1,2-Dibromoethane
Separation factors for the system 1,2-dibromoethane-1,2-dibromo[$^2H_4$]ethane 328(1985)333

Dibromopropamidine
Determination of preservatives in cosmetic products by ion-pair reversed-phase HPLC. III 348(1985)321

Dicarboxylic acids
4-methyl-7-methoxycoumarin as a fluorescent label for HPLC analysis of dicarboxylic acids 328(1985)111

HPLC of microbial acid metabolites 336(1985)125

Absolute configuration of 3-hydroxyadipic acid in human urine 337(1985)9

Dichlorobenzene
New liquid crystalline stationary phases for GC of positional and geometrical isomers having similar volatilities 331(1985)69

3,3'-Dichloro-4,4'-diaminodiphenylmethane
Capillary GC method for the determination of complex mixture of isocyanates and amines 346(1985)191

2,2-Dichloroethyl esters
GLC analyses. XLIII. Retention increments for 2-chloro-, 2,2-dichloro- and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns 329(1985)43

Dichloromethane
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

Diclobutrazol
Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181

Diclofenac
HPLC determination of diclofenac and its monohydroxylated metabolites in biological fluids 338(1985)151

Diclofensine
Determination of diclofensine, an antidepressant agent, and its major metabolites in human plasma by HPLC with fluorometric detection 341(1985)313

Dicumyl peroxide
Analysis technique for determining the levels of organic additives in an ethylene-propylene diene monomer rubber for use in underwater acoustic applications 350(1985)187

2-(Diethylamino)ethyl diphenylpropylacetate
HPLC separation and determination of 2-(diethylamino)ethyl diphenylpropylacetate and its metabolites from biological samples 342(1985)135

Diethylene glycol
HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Diethylenetriamine
Ion-pair HPLC determination of dansylated aliphatic polyamines with fluorescence and UV detection 333(1985)220

Di(2-ethylhexyl) phthalate
Determination of phthalic acid, mono-(2-ethylhexyl) phthalate and di(2-ethylhexyl) phthalate in human plasma and in blood products 337(1985)279

Determination of di- and mono(2-ethylhexyl) phthalate in plasma by GC 342(1985)399

Determination of di(2-ethylhexyl) phthalate and four of its metabolites in blood plasma by GC-MS 344(1985)167

Diethylstilboestrol
Rapid extraction and purification of diethylstilboestrol in bovine urine hydrolysates using reversed-phase C_{18} columns before determination by radioimmunoassay 328(1985)413

Application of a capillary GC-selected-ion recording MS technique to the analysis of diethylstilbestrol and its phosphorylated precursors in plasma and tissues 339(1985)87

Separation and detection of enantiomers of stilbestrol analogues by combined HPLC-thermospray MS 344(1985)378

Screening of parts-per-billion levels of dietyhlstilbestrol in bovine urine by HPLC with UV detection 347(1985)379

Diflunisal
Assay methodology for quantification of the ester and ether glucuronide conjugates of diflunisal in human urine 337(1985)363

α-Difluoromethylornithine
Determination of amino acids by separation of their ion pairs with dodecyl sulphate 341(1985)11

Digitalis glycosides
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Diglycerides
HPLC technique for the separation of lipid classes 329(1985)273

Dihydrocapsaicin
Formation and metabolism of pungent principle of *Capsicum* fruits. XV. Microdetermination of capsaicin by HPLC with electrochemical detection 329(1985)99

Dihydrocodeine
Improved method for the simultaneous determination of morphine, codeine and dihydrocodeine in blood by HPLC with electrochemical detection 347(1985)398

3,4-Dihydro-6-[4-(3,4-(dimethoxybenzoyl)-1-piperazinyl]-2(1*H*)-quinolinone
HPLC procedure for the determination of a new positive inotropic agent, 3,4-dihydro-6-[4-(3,4-(dimethoxybenzoyl)-1-piperazinyl]-2(1*H*)-quinolinone, in human plasma and urine 338(1985)450

Dihydrofuroquinolinic alkaloids
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

4,5-Dihydro-6-[4-(1H-imidazol-1-yl)phenyl]-3(2H)-pyridazinone
Determination of 4,5-dihydro-6[4-(1H-imidazol-1-yl)phenyl]-3(2H)-pyridazinone hydrochloride, a new cardiotonic, in plasma and urine by reversed-phase HPLC 336(1985)446

Dihydrolevobunolol
Determination of levobunolol and dihydrolevobunolol in blood and urine by HPLC using fluorescence detection 338(1985)444

Dihydropterins
Estimation of tetrahydro, dihydro and fully oxidised pterins by HPLC using sequential electrochemical and fluorometric detection 343(1985)35

Dihydropyridines
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

1,8-Dihydroxyanthraquinones
HPLC separation and quantitative determination of 1,8-dihydroxyanthraquinones in plant cell cultures 329(1985)311

Dihydroxybenzenes
Comparative GC studies of bonded and physically coated PEG 20M and its derivative phases with reference to some high-boiling isomeric solutes 346(1985)359

3,4-Dihydroxycinnamic acid
LC with voltammetric detection for quantitation of phenolic acids 348(1985)397

Dihydroxycoumarin methyl ethers
GC separation of monomethyl ethers of dihydroxycoumarin isomers 333(1985)175

trans-*10,11-Dihydroxy-10,11-dihydrocarbamazepine*
Simultaneous determination of anticonvulsant drugs and metabolites in plasma by HPLC 342(1985)469

Dihydroxydiphenylmethanes
Separation and identification of phenol-formaldehyde condensates by GC-MS. I. Acid-catalysed condensation products 329(1985)290

3,4-Dihydroxymandelic acid
Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

3,4-Dihydroxyphenylacetic acid
Comparison of HPLC with electrochemical detection and GC-mass fragmentography for the assay of salsolinol, dopamine and dopamine metabolites in food and beverage samples 336(1985)199

Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

3,4-Dihydroxyphenylethanol
Comparison of HPLC with electrochemical detection and GC-mass fragmentography for the assay of salsolinol, dopamine and dopamine metabolites in food and beverage samples 336(1985)199

Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

3,4-Dihydroxyphenylethylene glycol
Simultaneous assay of 3,4-dihydroxyphenylethylene glycol and norepinephrine in human plasma by HPLC with electrochemical detection 338(1985)401

Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

25,26-Dihydroxyvitamin D_3
Specific mass fragmentographic assay for 25,26-dihydroxyvitamin D in human plasma using a deuterated internal standard 338(1985)289

Diketogulonic acid
Separation of ascorbic acid, dehydroascorbic acid, diketogulonic acid and glucose by isocratic elution from a column of a hydrophilic gel 332(1985)283

1,3-Diketones
Adsorption chromatographic separations on buffered silica gel (Review) 334(1985)35

Diltiazem
Diltiazem and desacetyldiltiazem analysis in human plasma using HPLC: improved sensitivity without derivatization 342(1985)216

HPLC determination of diltiazem and four of its metabolites in plasma. Application to pharmacokinetics 345(1985)355

Dilution factor
HPLC theory for the practitioner (Review) 334(1985)197

Dimethophrine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

N-(3',4'-Dimethoxycinnamoyl)anthranilic acid
Simultaneous determination of tranilast and metabolites in plasma and urine using HPLC 345(1985)345

Dimethylamine
GC-MS analysis of volatile amines produced by several strains of *Clostridium* 337(1985)213

4-N,N-Dimethylaminoantipyrine
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

***p*-Dimethylaminobenzaldehyde**
Detection of citrazinic acid by PC and TLC 331(1985)202

Dimethylbenzamidines
Retention indices of dimethylbenzamidines and benzylideneamines on a non-polar column 330(1985)79

Dimethylbenzoic acid
Urinary dimethylbenzoic acid excretion as an indicator of occupational exposure to white spirit 337(1985)146

Dimethylcyclohexanes
Resolution of *cis*- and *trans*-dimethylcyclohexanes by partition GC through cyclodextrin complexes 349(1985)3

Dimethyl disulphide
GC and MS identification of sulphur gases in kraft digestion plants 346(1985)405

N^2,N^2-Dimethylguanosine
Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 339(1985)75

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

N-5-Dimethyl-9-[(2-methoxy-4-methylsulphonylamino)phenylamino]-4-acridinecarboxamide
Determination of N-5-dimethyl-9-[(2-methoxy-4-methylsulfonylamino)phenylamino]-4-acridinecarboxamide in plasma by HPLC 342(1985)431

Dimethylnaphthalenes
Correlations between retention on liquid crystalline stationary phases and chemical structure. I. Dimethylnaphthalenes 333(1985)11

2,6-Dimethylpyridine
GC separation of 2,6-dimethylpyridine and its chloro derivatives 347(1985)393

Dimethyl sulphide
GC and MS identification of sulphur gases in kraft digestion plants 346(1985)405

GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

Dimethyl sulphoxide
GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

***p*-(3,3-Dimethyl-1-triazeno)benzoic acid**
HPLC assay for the determination of *p*-(3,3-dimethyl-1-triazeno)benzoic acid in mouse plasma 345(1985)323

N-(3,5-Dinitrobenzoyl)-α-amino acid esters
Resolution of enantiomeric amino acid derivatives by HPLC on chiral stationary phases 348(1985)275

2,4-Dinitrophenol
Plasma levels and kinetic disposition of 2,4-dinitrophenol and its metabolites 2-amino-4-nitrophenol and 4-amino-2-nitrophenol in the mouse 344(1985)177

Dinoracetylmethadol
Quantitation of *l*-α-acetylmethadol and its metabolites in human serum by capillary GLC and nitrogen detection 343(1985)339

Diode-array detectors
Photodiode array UV spectrophotometric profiling of carboxylic acids in physiological fluids 344(1985)23

Dioleoyl phosphatidyl choline
Interaction of some non-ionic tensides with dioleoyl phosphatidyl choline, studied by charge-transfer chromatography 349(1985)295

Dioxane
Effects of test solutes on the characterization of stationary phases in GC using a Snyder triangle 329(1985)5

1,2-Dioxetanes
Determination of the thermal stability of some 1,2-dioxetanes by HPLC 347(1985)183

Dipeptides
Separation of free radicals by HPLC with electron spin resonance detection 332(1985)71

Direct liquid introduction MS of some underivatized dipeptides and polypeptides 350(1985)399

Diphenylamine
Measurement of trace aromatic amines in seawater using HPLC with electrochemical detection 348(1985)265

1,3-Diphenyl-2-aminopropane
Identification of 1,3-diphenyl-2-aminopropane metabolites by GC, GC with Fourier transform IR spectroscopy and GC-MS 337(1985)121

1,2-Diphenylethylenediamine
HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61

Diprobutine
Determination of diprobutine in human plasma using GLC with nitrogen-selective detection 339(1985)186

4-(2-Di-*n*-propylaminoethyl)-7-hydroxy-2-(3H)-indolone
Determination of 4-(2-di-*n*-propylaminoethyl)-7-hydroxy-2-(3H)-indolone and N-[2'-hydroxy-5'-(N,N-di-*n*-propylaminoethylphenyl)]methane sulfonamide in plasma by HPLC with electrochemical detection 345(1985)203

Direct liquid introduction interface
Gaseous acidity and basicity scales as guides to chemical ionization in reversed-phase LC-MS with direct liquid introduction 347(1985)61

Direct liquid introduction MS of some underivatized dipeptides and polypeptides 350(1985)399

Disopyramide
Determination of (*R*)- and (*S*)-disopyramide in human plasma using a chiral α_1-acid glycoprotein column 336(1985)321

Disperse dyes
HPLC method utilising single or multi-wavelength detection for the comparison of disperse dyes extracted from polyester fibres 350(1985)205

Displacement chromatography
Investigation of operating parameters in high-performance displacement chromatography 330(1985)1

Dissociation constants
Determination of the dissociation constant of oligomeric proteins by size-exclusion HPLC: application to human haemoglobin 327(1985)313

Energy effects in the retention of aromatic acids in LC 332(1985)189

Distribution coefficients
Possibility and consequences of using different concentration scales in the study of solution thermodynamics by GLC 350(1985)339

Disulphide bridges
Novel approach to rapid and sensitive localization of protein disulfide bridges by HPLC and electrochemical detection 326(1985)339

Disulphides
Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Dithiocarbamates
Reversed-phase ion-interaction chromatography of sodium dialkyldithiocarbamates in the presence of tetraalkylammonium salts 348(1985)389

Diuretics
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Influence of the chain length of chemically bonded phases on the behaviour of several thiazide, potassium-sparing and loop diuretics in HPLC 349(1985)301

DNA
HPLC of DNA (Review) 336(1985)3

Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 341(1985)182

Preparative isolation of plasmid DNA with sedimentation field flow fractionation 342(1985)67

DNA-cellulose chromatography
Application of DNA-cellulose chromatography in the isolation of immunoglobulin M and complement component C4b-binding protein from human serum 347(1985)129

DNA restriction fragments
HPLC of DNA restriction fragments 326(1985)251

Dobutamine
Rapid and simple method for the routine determination of dobutamine in human plasma by HPLC 345(1985)425

Dodine
GLC method for the determination of dodine in fruit 348(1985)437

Dolichyl pyrophosphate
Quantitation of tissue levels of dolichyl pyrophosphate by HPLC on silica 343(1985)413

DOPA
Determination of catecholamines by HPLC with catalytic photometric detection 329(1985)147

Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

Simultaneous determination of L-dopa and 3-O-methyldopa in human serum by HPLC 336(1985)380

Simultaneous assay of 3,4-dihydroxyphenylalanine, catecholamines and O-methylated metabolites in human plasma using HPLC 337(1985)239

Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

DOPA decarboxylase
1-DOPA decarboxylase and dopamine-β-hydroxylase isolated from sympathetic splenic nerves estimated by sodium dodecyl sulphate polyacrylamide gel electrophoresis with and without mercaptoethanol 329(1985)300

Assay of aromatic L-amino acid decarboxylase by HPLC with radiochemical detection 342(1985)175

Dopamine
Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Comparison of HPLC with electrochemical detection and GC-mass fragmentography for the assay of salsolinol, dopamine and dopamine metabolites in food and beverage samples 336(1985)199

3-Methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, and homovanillic acid in human cerebrospinal fluid. Storage and measurement by reversed-phase HPLC and coulometric detection using 3-methoxy-4-hydroxyphenyllactic acid as an internal standard 336(1985)259

Direct clean-up and analysis of urinary catecholamines 336(1985)374

Estimation of adrenal catecholamines by elevated-temperature LC with amperometric detection 342(1985)363

Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370

HPLC assay of free norepinephrine, epinephrine, dopamine, vanillylmandelic acid and homovanillic acid 343(1985)9

Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61

Rapid and reliable estimation of urinary free catecholamines in patients with pheochromocytoma: comparison with plasma catecholamines and vanillylmandelic acid excretion 344(1985)313

Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 345(1985)140

Amperometric detector for use with small-bore LC columns 350(1985)169

Dopamine β-hydroxylase
Characterization of the subunit structure of dopamine β-hydroxylase by anion-exchange HPLC 327(1985)351

1-DOPA decarboxylase and dopamine-β-hydroxylase isolated from sympathetic splenic nerves estimated by sodium dodecyl sulphate polyacrylamide gel electrophoresis with and without mercaptoethanol 329(1985)300

Simple and sensitive assay of dopamine β-hydroxylase in human cerebrospinal fluid by HPLC with electrochemical detection 341(1985)176

Dopamine sulphates
Determination of dopamine-3- and -4-O-sulphate in human plasma and urine by anion-exchange HPLC with fluorimetric detection 342(1985)261

Doxapram
Sensitive nitrogen-phosphorus capillary GC assay for doxapram in premature infants 344(1985)372

Doxazosin
Doxazosin determination by HPLC using fluorescence detection 344(1985)403

Doxorubicin
see Adriamycin

Droperidol
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Droplet counter-current chromatography
Increasing the speed of droplet counter-current chromatography separations 329(1985)395

Drug monitoring
Analytical strategies for therapeutic monitoring of drugs in biological fluids (Review) 340(1985)3

Drug level monitoring: cytostatics (Review) 340(1985)31

Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Drug level monitoring: Sedative hypnotics (Review) 340(1985)139

Analysis of tricyclic antidepressant drugs in plasma and serum by chromatographic techniques (Review) 340(1985)173

Determination of antiepileptic drugs (Review) 340(1985)199

Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Antibiotic monitoring in body fluids (Review) 340(1985)361

Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

Drugs
TLC screening procedure for undeclared synthetic drugs in Chinese herbal preparations 329(1985)107

Use of dual-column fused-silica capillary GC in combination with detector response factors for analytical toxicology 341(1985)81

Qualitative organic analysis. I. Identification of drugs by principal components analysis of standardized TLC data in four eluent systems 350(1985)151

Dyes
Identification of natural red dyes in old Indian textiles. Evaluation of TLC systems 347(1985)447

HPLC method utilising single or multi-wavelength detection for the comparison of disperse dyes extracted from polyester fibres 350(1985)205

Dyphylline
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

E

Ecdysteroids
Catalysed derivatisation of trimethylsilyl ethers of ecdysterone. A preliminary study 333(1985)186

Simultaneous determination of free and conjugated ecdysteroids by LC 346(1985)301

Ecdysterone
Catalysed derivatisation of trimethylsilyl ethers of ecdysterone. A preliminary study 333(1985)186

Effluent fractionation
High-sensitivity radioassay of chromatographic effluents. Automatic fraction collector/concentrator for quantitative autoradiography 349(1985)267

Elastomers
Chromatographic analysis of thermoplastic, medical-grade, polyether-based polyurethane elastomers 349(1985)347

Electrochemiluminescence
Electrochemiluminescence as a detection technique for reversed-phase HPLC. II. Low-frequency a.c. electrochemiluminescence 329(1985)281

Electroendosmosis
High-performance electrophoresis. Elimination of electroendosmosis and solute adsorption 347(1985)191

Electrokinetic chromatography
Electrokinetic chromatography with 2-O-carboxymethyl-β-cyclodextrin as a moving "stationary" phase 332(1985)211

Electrokinetic chromatography with micellar solutions. Separation of phenylthiohydantoin-amino acids 332(1985)219

Electrokinetic chromatography with micellar solutions. Retention behaviour and separation of chlorinated phenols 348(1985)39

Electron-capture detector
Selection of mobile phases for reversed-phase LC with on-line electron-capture detection 328(1985)93

Hypercoulometry in a pulsed conventional electron-capture detector 330(1985)87

Theory of anion contributions to non-linear electron-capture detector response 346(1985)1

Narrow-bore normal phase LC with on-line electron-capture detection 347(1985)75

Effects of contact potentials on the pulsed electron-capture detector 349(1985)155

Electron spin resonance detection
Separation of free radicals by HPLC with electron spin resonance detection 332(1985)71

Electrophoresis
Adaptation of the equipment for high-performance electrophoresis to isoelectric focusing 346(1985)265

High-performance electrophoresis. Elimination of electroendosmosis and solute adsorption 347(1985)191

Elution
Elution and washing in fixed bed affinity chromatography 330(1985)356

Emodin
HPLC separation and quantitative determination of 1,8-dihydroxyanthraquinones in plant cell cultures 329(1985)311

Enantiomer separation
Determination of optical purity by enantioselective capillary GC: panthenol and related compounds 328(1985)357

Enantiomeric resolution of *dl-threo*-methylphenidate, U.S.P. (Ritalin), by HPLC 328(1985)378

Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181

LC on triacetylcellulose. Preparative separation of enantiomers on an axially compressed column 329(1985)307

Rapid separation of enantiomers by TLC on a chiral stationary phase 330(1985)375

Application of micro HPLC to the separation of chiral amino acids 331(1985)99

Determination of the enantiomeric composition of chiral epoxides using chiral derivatization and LC 331(1985)349

Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59

Direct chiral resolution in an aqueous two-phase system using the counter-current distribution principle 333(1985)211

Resoltuion of *RS*-abscisic acid and the separation of abscisic acid metabolites from plant tissue by HPLC 336(1985)221

Determination of (*R*)- and (*S*)-disopyramide in human plasma using a chiral α_1-acid glycoprotein column 336(1985)321

HPLC assay for mexiletine enantiomers in human plasma 337(1985)172

Determination of the enantiomers of indoprofen in blood plasma by HPLC after rapid derivatization by means of ethyl chloroformate 339(1985)339

Urinary dihydrodiol metabolites of phenytoin: HPLC assay of diastereometic composition 342(1985)323

Separation and detection of enantiomers of stilbestrol analogues by combined HPLC-thermospray MS 344(1985)378

Column LC for the simultaneous determination of the enantiomers of loxoprofen sodium and its metabolites in human urine 345(1985)373

Isolation and stereospecific determination of the enantiomers of oxindazac by direct LC resolution on triacetylcellulose 346(1985)321

Molecular imprinting of amino acid derivatives in macroporous polymers. Demonstration of substrate- and enantio-selectivity by chromatographic resolution of racemic mixtures of amino acid derivatives 347(1985)1

Analytical separation of enantiomeric polygodials by GC of pyrrole derivatives 347(1985)455

Chromatographic separation of the enantiomers of 2-carboalkoxyindolines and N-aryl-α-amino esters on chiral stationary phases derived from N-(3,5-dinitrobenzoyl)-α-amino acids 348(1985)89

Resolution of enantiomeric amino acid derivatives by HPLC on chiral stationary phases 348(1985)275

High-performance liquid chromatograph coupled with two detectors: a UV spectrometer and a polarimeter. Example in the field of pyrethroids: identification of enantiomers 349(1985)109

Chiral recognition mechanisms in the enantioselectivity of chiral hydrogen bonding additives in LSC 349(1985)143

Chromatographic resolution. IX. HPLC resolution of enantiomers on optically active poly(triphenylmethyl methacrylate) 350(1985)127

Application of chromatographic chiral stationary phases to pharmaceutical analysis. Enantiomeric purity of D-penicillamine 350(1985)179

Enantiomer separation of metoprolol and its analogues and metabolites by capillary column GC after derivatization with phosgene 350(1985)328

Enatiomer separation
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 341(1985)213

β-Endorphin
HPLC of amino acids, peptides and proteins. LXV Studies on the optimisation of the reversed-phase gradient elution of polypeptides: evaluation of retention relationships with β-endorphin-related polypeptides 327(1985)115

HPLC analysis of *in vitro* central neuropeptide processing 327(1985)279

π-Energy effect
Energy effects in the retention of aromatic acids in LC 332(1985)189

Enkephalins
Improved detection limits in the analysis of tyrosine-containing polypeptide hormones by using electrochemical detection 327(1985)165

Amperometric detector for use with small-bore LC columns 350(1985)169

5-Enolpyruvylshikimate 3-phosphate
HPLC of shikimate pathway intermediates 329(1985)268

Enoxacin
HPLC and preliminary pharmacokinetics of enoxacin and its 4-oxo metabolite in human plasma, urine and saliva 343(1985)449

Enthalpy of solution
Possibility and consequences of using different concentration scales in the study of solution thermodynamics by GLC 350(1985)339

Enzyme entrapment
Use of silane monomers for molecular imprinting and enzyme entrapment in polysiloxane-coated porous silica 347(1985)11

Enzymes
see also Proteins

HPLC investigations on some enzymes of papaya latex 326(1985)103

Purification of oestrogen synthetase by HPLC. Two membrane-bound enzymes from the human plancenta 326(1985)137

Purification of gelatin-specific neutral protease from human skin by conventional and high-performance LC 326(1985)147

Purification of a uridine-specific acid nuclease from chicken liver by fast protein LC 326(1985)157

High-performance liquid affinity chromatography of β-lactamase 326(1985)163

The bacterial phosphoenolpyruvate-dependent phosphotransferase system. Isolation of active site peptides by reversed-phase HPLC and determination of their primary structure 326(1985)363

Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399

Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Application of a new ion exchanger TSK-GEL DEAE-5PW, to the purification of Cu,Zn-superoxide dismutase of bovine erythrocytes 327(1985)301

Characterization of the subunit structure of dopamine β-hydroxylase by anion-exchange HPLC 327(1985)351

Affinity electrophoresis in polyacrylamide gel. Influence of the gel concentration on the apparent affinity of cholinesterase for an anionic site ligand (in French) 328(1985)135

Purification of an acid protease and a serine carboxypeptidase from *Aspergillus niger* using metal-chelate affinity chromatography 329(1985)165

1-DOPA decarboxylase and dopamine-β-hydroxylase isolated from sympathetic splenic nerves estimated by sodium dodecyl sulphate polyacrylamide gel electrophoresis with and without mercaptoethanol 329(1985)300

Separation and partial characterization of *Trichoderma reesei* cellulase by fast chromatofocusing 329(1985)379

Separation of two forms of glutathione peroxidase from human erythrocytes by hydrophobic chromatography 330(1985)383

Use of Toyopearl as a support for the immobilization of urease 332(1985)237

Preparative high-performance ion-exchange chromatography 333(1985)29

High-performance affinity chromatography of trypsins on Asahipak GS-gel coupled with *p*-aminobenzamidine 333(1985)107

Preparative high-performance hydrophobic interaction chromatography of proteins on TSKgel Phenyl-5PW 333(1985)202

Use of chromatofocusing for separation of β-lactamases. V. Inducible chromosomally mediated β-lactamase of the *Enterobacter cloacae* 53 strain 333(1985)239

Use of chromatofocusing for separation of β-lactamases. VI. Comparative studies on two chromatofocusing polybuffer exchanger matrices with the R46 plasmid coded β-lactamase carried by the *Escherichia coli* K12 J5-3 strain 333(1985)244

Separation of tryptic phosphopeptides of ribosomal origin by reversed-phase HPLC 336(1985)51

Reversed-phase HPLC method for determination of brain glutamate decarboxylase suitable for use in kinetic studies 336(1985)385

Clinical analysis of steroids. XXXI. Assay of oestradiol 17-sulphate 4-hydroxylase activity by HPLC with electrochemical detection 337(1985)114

HPLC assays of phenylpyruvate and phenylpyruvate oxidase 337(1985)397

Assay of human erythrocyte pyrimidine and deoxypyrimidine 5'-nucleotidase by isocratic reversed-phase HPLC 339(1985)293

Chromatography of γ-glutamyl transferases from ascites hepatoma AH-66 cells and human primary hepatoma on *Phaseolus vulgaris* erythroagglutinating lectin agarose 339(1985)394

Rapid purification of human placental angiotensin I converting enzyme by captopril affinity chromatography 339(1985)399

Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 341(1985)23

Simple and sensitive assay of dopamine β-hydroxylase in human cerebrospinal fluid by HPLC with electrochemical detection 341(1985)176

Assay of aromatic L-amino acid decarboxylase by HPLC with radiochemical detection 342(1985)175

High-performance liquid chromatographic detection of XMP as a basis for an improved IMP dehydrogenase assay 343(1985)160

Simultaneous measurement of prolidase and prolinase activities in erythrocytes using an isotachophoretic analyser 343(1985)179

Highly sensitive assay for acetylcholinesterase activity by HPLC with electrochemical detection 344(1985)93

Combined recycling affinity column-filtration technique 346(1985)247

Isoelectric focusing separation of *Gliocladium* enzyme components 347(1985)310

Purification of D-amino acid oxidase apoenzyme by affinity chromatography on Cibacron Blue Sepharose 347(1985)316

Studies on lectins. LVIII. Sugar-binding properties, as determined by affinity electrophoresis, of α-D-galactosidases from *Vicia faba* seeds possessing erythroagglutinating activity 347(1985)343

HPLC of proteins. Rapid chromatofocusing of plant enzymes 348(1985)167

Highly sensitive assay for PZ-peptidase activity by HPLC 348(1985)205

Electrophoretic detection of peroxidase isoenzymes in polyacrylamide gels using hydrogen peroxide generated by rose bengal-glutathione in the presence of light 348(1985)313

Determination of phenylethanolamine N-methyltransferase by HPLC with fluorescence detection 348(1985)407

Simultaneous HPLC assay of the activities of erythrocytic hypoxanthine-guanine phosphoribosyl transferase and purine nucleoside phosphorylase 349(1985)275

HPLC of anthranilate synthase using gel filtration and a post-column reactor 350(1985)332

Ephedrine
Determination of sodium, ephedrine and procaine in pharmaceuticals by capillary isotachophoresis 330(1985)436

Epidermal cell derived thymocyte-activating factor
HPLC separation of distinct epidermal cell-derived cytokines 326(1985)129

Epidermal growth factors
Macroscale HPLC separation and instrumental identification of components of diethylaminoethyl murine epidermal growth factor 336(1985)73

Ion-exchange HPLC of mouse epidermal growth factor and its congeners: mobile phase optimization with ion-pairing additives 345(1985)168

Epinephrine
see Adrenaline

Epipolythiopiperazine-3,6-diones
TLC of mycotoxins (Review) 334(1985)211

Epoxides
Determination of the enantiomeric composition of chiral epoxides using chiral derivatization and LC 331(1985)349

Derivatization of alkyl halides and epoxides with picric acid salts for improved HPLC detection 333(1985)349

10,11-Epoxycarbamazepine
Simultaneous determination of anticonvulsant drugs and metabolites in plasma by HPLC 342(1985)469

Epoxyeicosatrienoic acids
Analysis of epoxyeicosatrienoic acids by GC-MS using chlorohydrin adducts 339(1985)175

Equivalent chain lengths
Use of equivalent chain lengths for the characterization of fatty acid methyl esters separated by linear temperature-programmed GC 346(1985)33

Ergopeptines
Retention indices of some ergopeptines in reversed-phase HPLC 331(1985)245

Ergot alkaloids
Retention indices of some ergopeptines in reversed-phase HPLC 331(1985)245

HPLC system for the separation of ergot alkaloids with applicability to the analysis of illicit lysergide (LSD) 346(1985)423

Erythromycins
Separation of erythromycin and related substances by HPLC on poly(styrene-divinylbenzene) packing materials 330(1985)275

Isolation of erythromycins and related substances from fermentation residues of *Streptomyces erythreus* by HPLC on silica gel 346(1985)309

Escherichia coli
Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 345(1985)11

Eserine
Sub-nanogram analysis of yohimbine and related compounds by HPLC 342(1985)333

Esmolol
HPLC method for the determination of esmolol hydrochloride in solutions and parenteral formulations 330(1985)420

Essential oils
see also Volatile oils

GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

General method for the HPLC prefractionation of essential oils and flavor mixtures for GC-MS analysis. Identification of

new constituents in cold pressed lime oil 330(1985)141

PC of the essential oils occurring in the genus Stachys 333(1985)288

Variation in the GC stationary phase properties of tetra-*n*-butylammonium salts as a function of the anion type 349(1985)235

Esters
GLC analyses. XLII. Retention behaviour of C_1-C_{12} *n*-alkyl esters of benzoic and pentafluorobenzoic acids on SE-30 and OV-351 capillary columns 328(1985)325

GLC analyses. XLIII. Retention increments for 2-chloro-, 2,2-dichloro- and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns 329(1985)43

GLC analyses. XLIV. Benzoyl and pentafluorobenzoyl esters of lower saturated branched-chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 329(1985)359

Ethambutol
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Ethanol
Effects of test solutes on the characterization of stationary phases in GC using a Snyder triangle 329(1985)5

HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Ether-bonded phases
High-performance hydrophobic-interaction chromatography on ether-bonded phases. Chromatographic characteristics and gradient optimization 326(1985)45

Ethionamide
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Ethosuximide
Rapid extraction method for ethosuximide and other antiepileptics in serum for determination by HPLC 338(1985)460

***o*-Ethoxybenzamide**
Analysis of antipyretics by semimicro LC 332(1985)269

7-Ethoxycoumarin
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Ethoxymethacrylate-ethylenedimethacrylate copolymer
Preparation of low-capacity anion-exchange resins for ion chromatography on a methacrylic copolymer matrix 350(1985)296

Ethyl chloroformate
Determination of the enantiomers of indoprofen in blood plasma by HPLC after rapid derivatization by means of ethyl chloroformate 339(1985)339

Ethylene
Determination of ethylene by a GC enrichment technique with a flame ionization detector (in German) 346(1985)43

Ethylenediamine
Ion-pair HPLC determination of dansylated aliphatic polyamines with fluorescence and UV detection 333(1985)220

Ethylenediammonium tartrate
Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147

Ethylene-propylene diene monomer rubber
Analysis technique for determining the levels of organic additives in an ethylene-propylene diene monomer rubber for use in underwater acoustic applications 350(1985)187

Ethylenic compounds
Paper electrophoretic characterization of ethylenic compounds as dimethylsulphoniomethylthio derivatives 328(1985)436

Ethylephrine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Ethyl loflazepate
Determination of circulating ethyl loflazepate metabolites in the baboon by radio-HPLC with injection of crude plasma samples: comparison with solvent extraction and TLC 342(1985)159

Ethyl-5-methoxysalicylate
HPLC determination of certain salicylates and their major metabolites in plasma following topical administration of a liniment to healthy subjects 342(1985)315

Ethylphosphine
Quantitative GC analysis of low-molecular-weight alkylphosphines in the presence of phosphine and hydrocarbons 330(1985)400

Ethyl phosphite
Determination of ethyl phosphite, phosphite and phosphate in plant tissues by anion-exchange HPLC and GC 347(1985)267

Etiocholanolone
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 341(1985)426

Etomidate
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Etoposide
HPLC determination of etoposide in plasma using electrochemical detection 336(1985)434

HPLC of etoposide in plasma and urine 339(1985)419

Drug level monitoring: cytostatics (Review) 340(1985)31

Evoxine
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Exopeptidase
Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399

Expert systems
Expert system for pharmaceutical analysis. I. Selection of the detection system in HPLC analysis: UV *versus* amperometric detection 348(1985)97

Explosives
Reversed-phase gradient HPLC of nitramine munitions and characterization of munitions process samples by GC-MS 329(1985)285

Microcolumn clean-up and recovery techniques for organic explosives compounds and for propellants traces in firearms discharge residues 330(1985)121

Metabolic studies of explosives. II. HPLC-MS of metabolites of 2,4,6-trinitrotoluene 339(1985)127

Extraction solvents
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

F

Factor VIII:C
Isolation of human factor VIII:C by preparative high-performance size-exclusion chromatography 326(1985)217

Famotidine
Analytical method for the quantification of famotidine, an H_2-receptor blocker, in plasma and urine 338(1985)438

Farnesol
Preparative HPLC of (2*E*)- and (2*Z*)-farnesol 347(1985)188

Fast-atom bombardment mass spectrometry
Direct coupling of micro HPLC with fast atom bombardment MS 346(1985)161

Fatty acid methyl esters
Analysis of labelled fatty acid methyl esters by argentation and reversed-phase two-dimensional TLC 330(1985)396

Evaluation of a rapid method for preparation of fatty acid methyl esters for analysis by GLC 331(1985)141

Use of equivalent chain lengths for the characterization of fatty acid methyl esters separated by linear temperature-programmed GC 346(1985)33

Fatty acids
Adsorption and reversed-phase HPLC of *p*-nitrobenzyl esters of monohydroxy fatty acids 329(1985)57

HPLC technique for the separation of lipid classes 329(1985)273

HPLC of fatty acids as their *p*-phenylazophenacyl esters 331(1985)187

HPLC separation of fatty acids as pentafluorobenzyl esters 336(1985)115

Assay of lipids in dog myocardium using capillary GC and derivatization with boron trifluoride and methanol 339(1985)25

HPLC of cardiolipin 339(1985)35

Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary GC-MS 341(1985)139

Unusual fatty acids from amniotic fluid phospholipids 341(1985)146

Fatty acid composition of human erythrocyte membranes by capillary GC-MS 342(1985)1

Differentiation between major species of the *Actinobacillus-Haemophilis-Pasteurella* group by GC of trifluoroacetic acid anhydride derivatives from whole-cell methanolysates 342(1985)13

Fatty acid analysis in phosphatidylethanolamine subclasses of human erythrocyte membranes by HPLC 342(1985)359

Purification of the peracetylated glycosphingolipids of the Gala series (galactosyl- and galabiosylceramides) 342(1985)385

Qualitative and quantitative measurement of hydroxy fatty acids, thromboxanes and prostaglandins using stable isotope dilutions and detection by GC-MS 344(1985)11

One-step separation of free fatty acids and phospholipids in brain tissue extracts by HPLC 344(1985)297

Application of silica gel CC in the assessment of non-esterified fatty acids and phosphoglycerides in myocardial tissue 344(1985)304

Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 345(1985)229

Improved solvent-extraction based procedure for the GC analysis of resin and fatty acids in pulp mill effluents 346(1985)205

3-Bromomethyl-6,7-dimethoxy-1-methyl-2(1*H*)-quinoxalinone as a new fluorescence derivatization reagent for carboxylic acids in HPLC 346(1985)227

Semi-preparative isolation of crepenynic acid, a potential inhibitor of essential fatty acid metabolism 346(1985)431

Gradient separation of fatty acids (C_{14}-C_{30}) by reversed-phase HPLC 346(1985)453

HPLC of triglycerides of Flacourtiaceae seed oils containing cyclopentenyl fatty acids (chaulmoogric oils) 348(1985)441

Preseparation of polar, substituted, polycyclic aromatic hydrocarbons in aerosol extracts by HPLC on acidic- or basic-buffered silica 349(1985)331

FD&C Yellow No. 6
Determination of unsulfonated aromatic amines in FD&C Yellow No. 6 by the diazotization and coupling procedure followed by reversed-phase HPLC 331(1985)109

Determination of the lower sulfonated subsidiary colors in FD&C Yellow No. 6 by reversed-phase HPLC 347(1985)163

Felodipine
Bimodal column switching LC assay of six metabolites of [^{14}C]felodipine in rat urine 343(1985)85

Fenac compounds
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77

Fenamates
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77

Fenbufen
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77

Fenoterol
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Fentanyl
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Fetoprotein
Simple purification procedure of rat α-fetoprotein by a combination of Cibacron Blue gel affinity chromatography and anion-exchange HPLC 338(1985)410

Fractionation of rat α-fetoprotein by HPLC 341(1985)452

Fezolamine
Determination of fezolamine and its desmethyl metabolite in human plasma and urine by HPLC. Intravenous pharmacokinetics in the beagle hound 344(1985)211

Fibronectin
Interaction between fluorescence-labeled fibronectin fragments studied by gel HPLC 327(1985)343

Field flow fractionation
Preparative isolation of plasmid DNA with sedimentation field flow fractionation 342(1985)67

Film thickness
Contribution to the study of micropacked columns in GC 348(1985)327

Flavin adenine dinucleotide
Use of HPLC in the preparation of flavin adenine dinucleotide analyte conjugates 326(1985)301

Flavin adenine dinucleotide theophylline conjugate
Use of HPLC in the preparation of flavin adenine dinucleotide analyte conjugates 326(1985)301

Flavin N^6-(N-6-aminohexyl)adenine dinucleotide
Use of HPLC in the preparation of flavin adenine dinucleotide analyte conjugates 326(1985)301

Flavivirus proteins
Separation of flavivirus membrane and capsid proteins by multistep HPLC optimized by immunological monitoring 326(1985)113

Flavonoids
Separation and determination of the various constituents of troxerutin by HPLC. Application to the pharmaceutical control (in French) 346(1985)382

HPLC of caffeoylquinic acid derivatives of *Cynara scolymus* L. leaves 348(1985)447

Flecainide acetate
HPLC method for the quantitation of a *meta*-O-dealkylated metabolite of flecainide acetate, a new antiarrhythmic 343(1985)119

Flow-pressure gradients
Gradient programming and combined gradient-pressure programming in supercritical fluid chromatography 349(1985)317

Flow programming
Optimized flow programming for temperature-programmed GC 328(1985)49

Flow-rate
HPLC theory for the practitioner (Review) 334(1985)197

Flow velocity
HPLC theory for the practitioner (Review) 334(1985)197

Fludalanine
Determination of the antibiotic fludalanine in plasma and urine by HPLC using a packed-bed, post-column reactor with *o*-phthalaldehyde and 2-mercaptoethanol 338(1985)357

9-Fluorenylmethylchloroformate
Selective determination of secondary amino acids using precolumn derivatization with 9-fluorenylmethylchloroformate and reversed-phase HPLC 348(1985)213

Fluorescein
HPLC analysis for fluorescein and fluorescein monoglucuronide in plasma 338(1985)213

5-Fluorocytosine
Micro determination of 5-fluorocytosine by HPLC and fluorimetric detection (in French) 345(1985)436

5-Fluorouracil
Drug level monitoring: cytostatics (Review) 340(1985)31

Analysis of 5'-deoxy-5-fluorouridine and 5-fluorouracil in human plasma and urine by HPLC 342(1985)303

Highly sensitive determination of 5-fluorouracil in human plasma by capillary GC and negative ion chemical ionization MS 343(1985)59

Sensitive HPLC method for the determination of 5-fluorouracil in plasma 344(1985)385

Flurazepam
Simple TLC method for the investigation of six related compounds in flurazepam hydrochloride and its capsules 350(1985)273

Flurbiprofen
Rapid and selective derivatization method for the nitrogen-selective detection of carboxylic acids in biological fluids prior to GC analysis 336(1985)241

Fluzinamide
Simultaneous determination of fluzinamide and three of its active metabolites in plasma by HPLC 336(1985)329

Fly ash
Semi-preparative HPLC analysis of complex organic mixtures 332(1985)169

Folates
Stabilization of 5-methyltetrahydrofolic acid and subsequent analysis by reversed-phase HPLC 328(1985)417

Formaldehyde
Determination of low concentrations of low-molecularweight aldehydes and ketones in aqueous samples 329(1985)81

Determination of formaldehyde as its lutidine derivative in the atmospheric liquid phase by HPLC 333(1985)262

Characterization of neutralized β-naphthalenesulfonic acid and formaldehyde condensates 347(1985)249

Formic acid
Characterization of a solvent system for separation of water-insoluble poliovirus proteins by reversed-phase HPLC 326(1985)91

N-Formyl-DL-tryptophan
Chromatographic determination of N-acetyl-DL-tryptophan and octanoic acid in human albumin solutions 333(1985)381

N^{in}-Formyltryptophan
Purification of synthetic lipid associating peptides and the monitoring of the deformylation of N^{in}-formyltryptophan by reversed-phase HPLC 347(1985)237

Fraction collection
Monitoring radioactive compounds in HPLC eluates: fraction collection versus on-line detection 342(1985)25

High-sensitivity radioassay of chromatographic effluents. Automatic fraction collector/concentrator for quantitative autoradiography 349(1985)267

Free energy of solution
Possibility and consequences of using different concentration scales in the study of solution thermodynamics by GLC 350(1985)339

Free radicals
Separation of free radicals by HPLC with electron spin resonance detection 332(1985)71

Fructose
Elevation of certain polyols in the cerebrospinal fluid of patients with multiple sclerosis 336(1985)351

Isolation and quantitation of carbohydrates in sheep plasma by HPLC 338(1985)303

HPLC analysis of glucose, fructose, sorbitol and low levels of the aldose reductase inhibitor sorbinil in human lens and plasma 343(1985)402

HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Fumitremorgin
TLC of mycotoxins (Review) 334(1985)211

Functional group contribution concept
Utilization of the functional group contribution concept in LC on chemically bonded reversed phases 348(1985)49

Furans
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

Furocoumarins
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Furoquinolinic alkaloids
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Furosine
Determination of furosine by GLC 346(1985)363

G

Gabapentin
Determination of gabapentin in plasma and urine by HPLC and pre-column labelling for UV detection 341(1985)473

Gabexate mesilate
HPLC assay of anti-inflammatory drugs incorporated in gel ointments. Separation and stability testing 331(1985)383

Galabiosylceramides
Purification of the peracetylated glycosphingolipids of the Gala series (galactosyl- and galabiosylceramides) 342(1985)385

α-D-Galactosidases
Studies on lectins. LVIII. Sugar-binding properties, as determined by affinity electrophoresis, of α-D-galactosidases from *Vicia faba* seeds possessing erythroagglutinating activity 347(1985)343

Galactosylceramides
Purification of the peracetylated glycosphingolipids of the Gala series (galactosyl- and galabiosylceramides) 342(1985)385

Gallamine
Quantitation of gallamine (Flaxedil) in human plasma using HPLC 345(1985)184

Gangliosides
Normal-phase HPLC separation of non-derivatized ganglioside mixtures 348(1985)371

Gas chromatographic peak yield
Application of radioisotope tracer techniques to analytical GC: determination of GC peak yield 341(1985)251

Gases
Noble, diatomic and aliphatic gas analysis by aqueous HPLC 346(1985)93

Determination of permanent gases and light hydrocarbons by simultaneous operation on packed and capillary columns with thermal conductivity detection and flame-ionization detection on a single computer integrator 349(1985)17

Gas sampling
The re-injection method for recovery of components adsorbed during syringe gas sampling: a statistical evaluation 329(1985)119

Gastrin
Rapid, small-scale preparation of gastrointestinal hormones by HPLC on a C_{18} column 326(1985)121

Gelatinase
Purification of gelatin-specific neutral protease from human skin by conventional and high-performance LC 326(1985)147

Gel concentration
Affinity electrophoresis in polyacrylamide gel. Influence of the gel concentration on the apparent affinity of cholinesterase for an anionic site ligand (in French) 328(1985)135

Gel packing materials
Recent progress in gel packing materials and detectors for modern LC in Japan 332(1985)227

Gentamicin
Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Gentiopicrin
Analytical and preparative reversed-phase LC of secoiridoid glycosides 346(1985)396

Gentisic acid
HPLC determination of certain salicylates and their major metabolites in plasma following topical administration of a liniment to healthy subjects 342(1985)315

Geranylgeraniol
Separation of geranylgeraniol isomers by HPLC and identification by ^{13}C nuclear magnetic resonance spectroscopy 347(1985)275

Ghatti gum
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Glass-lined stainless-steel tubing columns
Advanced microcolumn techniques in LC. Use of glass-lined stainless-steel tubing columns 332(1985)3

Gliadins
Detection of quality differences among wheats by HPLC 327(1985)333

Glibenclamide
Determination of glibenclamide, chlorpropamide and tolbutamide in plasma by HPLC with UV detection 339(1985)203

***Gliocladium* enzymes**
Isoelectric focusing separation of *Gliocladium* enzyme components 347(1985)310

Globulins
Purification of human serum gamma globulins by hydrophobic interaction HPLC 326(1985)235

Isotachophoresis as a useful tool for monitoring neurological complications of acute leukaemia in children 342(1985)285

Glucagon
Homogenous [mono-^{125}I-Tyr^{1}]- and [mono-^{125}I-Tyr^{13}] glucagon 331(1985)125

Purification of radioiodinated photoactivable glucagon by isocratic HPLC 331(1985)133

1-(β-D-Glucopyranosyl)phenobarbital
HPLC analysis of 1-(β-D-glucopyranosyl)phenobarbital in urine 343(1985)219

Glucose
Amperometric detection of reducing carbohydrates in HPLC using an amino-bonded column and acetonitrile-water as the eluent 330(1985)333

Separation of ascorbic acid, dehydroascorbic acid, diketogulonic acid and glucose by isocratic elution from a column of a hydrophilic gel 332(1985)283

Isolation and quantitation of carbohydrates in sheep plasma by HPLC 338(1985)303

Plasma amino acid analysis by GC. Removal of glucose interference 339(1985)366

Quantitative method for the detection of glucose in body fluids by HPLC with fluorescence detection 342(1985)179

HPLC analysis of glucose, fructose, sorbitol and low levels of the aldose reductase inhibitor sorbinil in human lens and plasma 343(1985)402

HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Glucose-1-phosphate
HPLC determination of glucose-1-phosphate and glucuronic acid-1-phosphate 330(1985)412

3-O-Glucosyl 11-deoxyjervine
Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

Glucuronic acid-1-phosphate
HPLC determination of glucose-1-phosphate and glucuronic acid-1-phosphate 330(1985)412

Glucuronides
Simultaneous determination of baicalein, wogonin, oroxylin-A and their glucuronides in Scutellariae Radix by ion-pair HPLC 328(1985)289

Glutamate decarboxylase
Reversed-phase HPLC method for determination of brain glutamate decarboxylase suitable for use in kinetic studies 336(1985)385

γ-Glutamylcysteine
Rapid and simple method for quantitative determination of non-protein sulphydryls in mouse liver by reversed-phase HPLC 338(1985)209

γ-Glutamyl transferase
Chromatography of γ-glutamyl transferases from ascites hepatoma AH-66 cells and human primary hepatoma on *Phaseolus vulgaris* erythroagglutinating lectin agarose 339(1985)394

Glutathione
Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193

Rapid and simple method for quantitative determination of non-protein sulphydryls in mouse liver by reversed-phase HPLC 338(1985)209

Determination of cysteine and glutathione in plasma and blood by LC with electrochemical detection using a chemically modified electrode containing cobalt phthalocyanine 345(1985)43

Glutathione peroxidase
Separation of two forms of glutathione peroxidase from human erythrocytes by hydrophobic chromatography 330(1985)383

Glutenins
Detection of quality differences among wheats by HPLC 327(1985)333

Glyceollin
HPLC of isoflavonoid phytoalexins from soybean (*Glycine max*, L. Harosoy 63) 333(1985)225

Glycerol
Determination of glycerol in pharmaceutical preparations by reversed-phase HPLC using a refractive index detector 329(1985)434

HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Glycerolipids
Diastereomeric separations of natural glycero derivatives as their 1-(1-naphthyl)ethyl carbamates by HPLC 350(1985)417

D-Glycero-D-mannoheptose
Differentiation between major species of the *Actinobacillus-Haemophilis-Pasteurella* group by GC of trifluoroacetic acid anhydride derivatives from whole-cell methanolysates 342(1985)13

Glycinebetaine aldehyde
HPLC method for the separation and estimation of choline, glycinebetaine aldehyde and related compounds 350(1985)245

Glycoalkaloids
Determination of the total C_{27}-steroidal alkaloid composition of *Solanum* species by high resolution GC 331(1985)285

Glycolipids
Isolation and quantitative analysis of phosphatidylglycerol and glycolipid molecular species using reversed-phase HPLC with flame ionization detection 346(1985)291

Glycopeptides
Automated tandem HPLC system for separation of extremely complex peptide mixtures 326(1985)407

Glycoproteins
Tandem separation of labelled human blood platelet membrane glycoproteins by anion-exchange and gel fast protein LC 326(1985)179

Isolation of the subunits of the coronavirus envelope glycoprotein E2 by hydroxyapatite HPLC 326(1985)191

Purification of hemopexin and its domain fragments by affinity chromatography and HPLC 326(1985)373

Isolation and characterization of two minor fractions of α_1-antitrypsin by HPLC chromatofocusing 327(1985)173

Interaction between fluorescence-labeled fibronectin fragments studied by gel HPLC 327(1985)343

Interaction of synthetic glycoproteins with immobilized lectins 330(1985)392

Measurement of neutral sugars in glycoproteins as dansyl derivatives by automated HPLC 333(1985)123

Simple purification procedure of rat α-fetoprotein by a combination of Cibacron Blue gel affinity chromatography and anion-exchange HPLC 338(1985)410

Evidence for the degradation of sugars during the resin hydrolysis of glycoproteins 339(1985)182

Two-step purification of human α_1-acid glycoprotein 341(1985)457

Glycosides
HPLC assay for the antitumor glycoside phyllanthoside and its stability in plasma of several species 342(1985)129

Glycosphingolipids
Purification of the peracetylated glycosphingolipids of the Gala series (galactosyl- and galabiosylceramides) 342(1985)385

Glycylglycine
Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Glycylmesidide
Determination of trimecaine metabolites in blood plasma by capillary isotachophoresis 342(1985)167

Gonadotropin
HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25

Gossypol
HPLC analysis of terpene aldehydes in cotton 329(1985)91

HPLC with electrochemical detection of gossypol in human plasma 343(1985)387

Gradient elution
High-performance hydrophobic-interaction chromatography on ether-bonded phases. Chromatographic characteristics and gradient optimization 326(1985)45

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Application to method development and the choice of column configuration 327(1985)93

HPLC of amino acids, peptides and proteins. LXV. Studies on the optimisation of the reversed-phase gradient elution of polypeptides: evaluation of retention relationships with β-endorphin-related polypeptides 327(1985)115

Spectral properties of trifluoroacetic acid-acetonitrile gradient systems for separation of picomole quantities of peptides by reversed-phase HPLC 347(1985)83

Effect of column temperature on HPLC behaviour of inorganic polyphosphates. II. Gradient ion-exchange chromatography 350(1985)119

Grape aroma components
The aroma of grapes. I. Extraction and determination of free and glycosidically bound fractions of some grape aroma components 331(1985)83

Graphitized carbon black
Adsorption of mixtures of acetone and benzene vapors on graphitized carbon black 328(1985)63

Growth factors
Macroscale HPLC separation and instrumental identification of components of diethylaminoethyl murine epidermal growth factor 336(1985)73

Partial purification of a specific inhibitor of the insulin-like growth factors by reversed-phase HPLC 336(1985)87

Ion-exchange HPLC of mouse epidermal growth factor and its congeners: mobile phase optimization with ion-pairing additives 345(1985)168

Growth hormone
Preparative high-performance ion-exchange chromatography 333(1985)29

HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25

Guaiacol glycerol ether
HPLC of antipyretics on chemically modified porous glass 332(1985)265

Analysis of antipyretics by semimicro LC 332(1985)269

Guanidino compounds
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

HPLC analysis of guanidino compounds using ninhydrin reagent. II. Guanidino compounds in blood of patients on haemodialysis therapy 342(1985)269

Quantitative overestimation of free taurocyamine using short liquid cation-exchange chromatography columns? 345(1985)215

Guanidinosuccinic acid
HPLC analysis of guanidino compounds using ninhydrin reagent. II. Guanidino compounds in blood of patients on haemodialysis therapy 342(1985)269

Guanosine
Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 345(1985)457

Guar gum
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Gums
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

H

Haemoglobin
Determination of the dissociation constant of oligomeric proteins by size-exclusion HPLC: application to human haemoglobin 327(1985)313

Electrophoretic and chromatographic techniques for the differential diagnosis of a haemoglobin abnormality: Hb E heterozygosity 330(1985)299

Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Isolation of normal and variant human hemoglobin subunits 343(1985)168

New less temperature-sensitive microchromatographic method for the separation and quantitation of glycosylated hemoglobins using a non-cyanide buffer system 344(1985)81

Analytical isoelectric focusing of native and modified haemoglobins after treatment with 4-hydroxymercuribenzoate 346(1985)401

Haemopexin
Purification of hemopexin and its domain fragments by affinity chromatography and HPLC 326(1985)373

Haemophilus
Differentiation between major species of the *Actinobacillus-Haemophilis-Pasteurella* group by GC of trifluoroacetic acid anhydride derivatives from whole-cell methanolysates 342(1985)13

Halides
Trace analysis of organic compounds in volatile inorganic halides by the dynamic headspace technique and GC-MS 350(1985)324

Hall electrolytic conductivity detector
Temperature selection for chlorinated hydrocarbon reduction for the Hall electrolytic conductivity detector 328(1985)342

Halogenated organic compounds
Temperature-programmed retention indices of 221 halogenated organic compounds with 1-bromoalkanes as references 328(1985)35

Halogenobicyclononene carbonitriles
Separation of isomeric halogenobicyclononene carbonitriles 347(1985)297

Haloperidol
Improved GC-MS assay for haloperidol utilizing ammonia chemical ionization and selected-ion monitoring 339(1985)321

Sensitive analysis of butyrophenone neuroleptics by HPLC with UV detection at 254 nm 341(1985)465

Hapten number
Clinical analysis of steroids. XXXII. New method for the determination of hapten number of the antigen prepared for the radioimmunoassay of steroidal sulphates using ion chromatography 338(1985)404

Heliocides
HPLC analysis of terpene aldehydes in cotton 329(1985)91

Helium ionization detector
Optimization of helium ionization detector 349(1985)405

Integration of the deformed peak from the helium ionization detector 349(1985)415

Hemiformal compounds
Separation and identification of phenol-formaldehyde condensates by GC-MS. III. Hemiformal isomers in resol 333(1985)161

Heptadec-10-enoate
HPLC separation of fatty acids as pentafluorobenzyl esters 336(1985)115

***n*-Heptane**
Reversed-flow GC technique applied to measurement of mass transfer coefficients of *n*-hydrocarbons on Porapak P 348(1985)339

Heterometallic clusters
Catalytic hydrogenation and exchange reactions on GC columns, in the presence of supported $(\eta^5\text{-}C_5H_5)NiOs_3(\mu\text{-}H)_3(CO)_9$ 349(1985)173

***Z*-11-Hexadecenal**
Determination of trimer in *Z*-11-hexadecenal pheromone by HPLC with a rotating disc flame ionization detector 329(1985)142

Hexamidine
Determination of preservatives in cosmetic products by ion-pair reversed-phase HPLC. III 348(1985)321

***n*-Hexane**
Reversed-flow GC technique applied to measurement of mass transfer coefficients of *n*-hydrocarbons on Porapak P 348(1985)339

High-density lipoproteins
HPLC of apolipoproteins in serum high-density lipoproteins 336(1985)151

High-mobility group proteins
Analysis and preparation of chromosomal high-mobility group proteins by ion-exchange HPLC 344(1985)332

High-performance electrophoresis
Micropreparative version of high-performance electrophoresis. The electrophoretic counterpart of narrow-bore HPLC 327(1985)157

Adaptation of the equipment for high-performance electrophoresis to isoelectric focusing 346(1985)265

High-performance electrophoresis. Elimination of electroendosmosis and solute adsorption 347(1985)191

Histamine
HPTLC analysis of dialysable materials in allergen extracts 329(1985)153

Separation of cadaverine from putrescine, histamine and polyamines in rat kidney by phosphocellulose chromatography 337(1985)103

Comparative evaluation of the radioenzymatic method for the determination of urinary histamine with a MS assay 338(1985)41

Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

HPLC determination of plasma and brain histamine without previous purification of biological samples: cation-exchange chromatography coupled with post-column derivatization fluorometry 344(1985)115

Histidine
Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

HMX
Reversed-phase gradient HPLC of nitramine munitions and characterization of munitions process samples by GC-MS 329(1985)285

Microcolumn clean-up and recovery techniques for organic explosives compounds and for propellants traces in firearms discharge residues 330(1985)121

Homocysteine
Determination of homocysteine in urine 338(1985)201

Homovanillic acid
Measurement of urinary vanilmandelic acid and homovanillic acid by HPLC with electrochemical detection following extraction by ion-exchange and ion-moderated partition 336(1985)173

3-Methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, and homovanillic acid in human cerebrospinal fluid. Storage and measurement by reversed-phase HPLC

and coulometric detection using 3-methoxy-4-hydroxyphenyllactic acid as an internal standard 336(1985)259

GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

HPLC assay of free norepinephrine, epinephrine, dopamine, vanillylmandelic acid and homovanillic acid 343(1985)9

Determination of homovanillic acid in human plasma by HPLC with electrochemical detection 343(1985)155

Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

Hop acids
Adsorption chromatographic separations on buffered silica gel (Review) 334(1985)35

Hormone receptors
Isolation of unliganded steroid receptor proteins by high-performance size-exclusion chromatography. An investigation of steroid-dependent structural alterations 327(1985)221

High-performance hydrophobic-interaction chromatography of steroid hormone receptors 327(1985)237

High-performance chromatofocusing of steroid receptor proteins in the presence and absence of steroid. Investigation of steroid-dependent alterations in surface charge heterogeneity 327(1985)247

Rapid, high-resolution procedure for assessment of estrogen receptor heterogeneity in clinical samples 327(1985)369

Hormones
see also Plant hormones and steroids

Rapid, small-scale preparation of gastrointestinal hormones by HPLC on a C_{18} column 326(1985)121

Improved detection limits in the analysis of tyrosine-containing polypeptide hormones by using electrochemical detection 327(1985)165

Quantitative affinity HPLC of neuroendocrine polypeptides using porous and non-porous glass derivatives 327(1985)193

Comparison of reversed-phase and cation-exchange HPLC for separating closely related peptides: separation of Asp^{76}-human parathyroid hormone (1-84) from Asn^{76}-human parathyroid hormone (1-84) 327(1985)327

Homogenous [mono-^{125}I-Tyr^{1}]- and [mono-^{125}I-Tyr^{13}] glucagon 331(1985)125

Purification of radioiodinated photoactivable glucagon by isocratic HPLC 331(1985)133

Preparative high-performance ion-exchange chromatography 333(1985)29

HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25

Measurement and chromatographic characterization of vasoactive intestinal peptide from guinea-pig enteric nerves 336(1985)41

HPLC and studies of neurophysin-neurohypophysial hormone pathways 336(1985)63

Reversed-phase HPLC of C_{21} metabolites of progesterone 336(1985)361

Determination of testosterone propionate in human plasma by GC-MS 337(1985)205

Total oestriol in maternal serum or plasma as measured by LC 337(1985)379

Determination of plasma testosterone by mass fragmentography using [3,4-^{13}C]testosterone as an internal standard 339(1985)233

Efficient chromatographic fractionation of steroids in human serum through regulation of liquid-liquid distribution ratios 339(1985)253

Determination of oestriol in pregnancy urine by normal-phase HPLC with electrochemical detection 341(1985)271

Determination of progesterone by LC-MS using a moving-belt interface and isotope dilution 343(1985)397

Reversed-phase HPLC of insulin and insulin derivatives. A comparative study 348(1985)347

Identification and isolation of human insulin A and B chains by HPLC 349(1985)91

HT-2 toxin
GC assay with pharmacokinetic applications for monitoring T-2 and HT-2 toxins in plasma 343(1985)67

Humic acids
Size-exclusion chromatography-low-angle laser light-scattering photometry of lignite macromolecules 349(1985)337

Hyaluronate
Comparative study of affinity chromatography of components of the hyaluronate-proteoglycan complex to immobilized hyaluronate 350(1985)227

Hyaluronate oligosaccharides
Comparison of relationships between the chemical structures and mobilities of hyaluronate oligosaccharides in TLC and HPLC 328(1985)73

Hydralazine
Determination of hydralazine and metabolites in urine by LC with electrochemical detection 343(1985)99

Quantitation of hydralazine hydrochloride in pharmaceutical dosage forms by HPLC 347(1985)414

Hydrazine derivatives
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Hydrocarbon gases
On the possibility of increasing the effectiveness of chromatographic columns filled with zeolites 331(1985)402

Hydrocarbons
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

Trace analysis of organic pollutants in air by polymer adsorption and dual flame ionization detection-electron-capture detection capillary GC 328(1985)367

Quantitative GC analysis of low-molecular-weight alkylphosphines in the presence of phosphine and hydrocarbons 330(1985)400

Noble, diatomic and aliphatic gas analysis by aqueous HPLC 346(1985)93

Quantitative analysis of hydrocarbons in gasolines by capillary GLC. II. Isothermal and temperature-programmed analyses 346(1985)177

Unified retention index of hydrocarbons separated on squalane 347(1985)366

Reversed-flow GC technique applied to measurement of mass transfer coefficients of *n*-hydrocarbons on Porapak P 348(1985)339

Determination of permanent gases and light hydrocarbons by simultaneous operation on packed and capillary columns with thermal conductivity detection and flame-ionization detection on a single computer integrator 349(1985)17

Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. I. Introduction and characterization of the cuticular hydrocarbon patterns of *Solenopsis invicta* and *S. richteri* 349(1985)31

Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. II. Comparison of the cuticular hydrocarbon patterns between different colonies of *Solenopsis richteri* 349(1985)39

Hydrogen-bonding energy
Energy effects in the retention of aromatic acids in LC 332(1985)189

Hydrogen sulphide
GC and MS identification of sulphur gases in kraft digestion plants 346(1985)405

On-line analysis of sulphur compounds in natural gas with flame photometric detection 349(1985)465

Hydroperoxides
HPLC as a tool for identification of linolenic acid hydroperoxide prepared with soybean lipoxygenase 350(1985)435

HPLC of linoleic acid hydroperoxides and their corresponding alcohol derivatives 350(1985)445

13-Hydroperoxylinolenic acid
HPLC as a tool for identification of linolenic acid hydroperoxide prepared with soybean lipoxygenase 350(1985)435

Hydrophobicity
see also Lipophilicity

LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. II. Quantitative determination of hydrophobicity 350(1985)105

Hydroquinones
HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. II. Retention behavior, quantitative structure-retention relationships and octanol-water partition coefficients 330(1985)181

Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 345(1985)11

Hydroxy acids
Counter-ion effects in partition chromatography 332(1985)203

3-Hydroxyadipic acid
Absolute configuration of 3-hydroxyadipic acid in human urine 337(1985)9

Hydroxyamphetamine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Hydroxyanthraquinones
TLC of mycotoxins (Review) 334(1985)211

4-Hydroxybenzoic acid
Analysis of antipyretics by semimicro LC 332(1985)269

GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

Studies on the analysis of food additives by HPLC. V. Simultaneous determination of preservatives and saccharin in foods by ion-pair chromatography 346(1985)333

p-Hydroxybestatin
Determination of p-hydroxybestatin in human serum by HPLC using fluorescence detection 344(1985)267

6β-Hydroxybetamethasone
Quantitative determination of betamethasone and its major metabolite in equine urine by micro-LC-MS 341(1985)261

3-Hydroxy-p-butyrophenetidine
HPLC of antipyretics on chemically modified porous glass 332(1985)265

Analysis of antipyretics by semimicro LC 332(1985)269

Hydroxychloroquine
Estimation of plasma hydroxychloroquine by HPLC with UV detection 338(1985)422

HPLC assay for hydroxychloroquine and metabolites in blood and plasma, using a stationary phase of poly(styrene divinylbenzene) and a mobile phase at pH 11, with fluorimetric detection 344(1985)241

25-Hydroxycholesterol
Analysis of 25-hydroxycholesterol in plasma and cholesterol-containing foods by HPLC 330(1985)339

7-Hydroxycoumarin
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

15β-Hydroxycyproterone acetate
15-Hydroxycyproterone acetate and cyproterone acetate levels in plasma and urine 338(1985)179

3-Hydroxydescarboxyloflazepate
Determination of circulating ethyl loflazepate metabolites in the baboon by radio-HPLC with injection of crude plasma samples: comparison with solvent extraction and TLC 342(1985)159

Hydroxydiclofenac
HPLC determination of diclofenac and its monohydroxylated metabolites in biological fluids 338(1985)151

10-Hydroxy-10,11-dihydrocarbamazepine
Simultaneous determination of anticonvulsant drugs and metabolites in plasma by HPLC 342(1985)469

N-[2'-Hydroxy5'-(N,N-di-n-propylaminoethylphenyl)]methane-sulphonamide
Determination of 4-(2-di-n-propylaminoethyl)-7-hydroxy-2-(3H)-indolone and N-[2'-hydroxy-5'-(N,N-di-n-propylaminoethylphenyl)]methane sulfonamide in plasma by HPLC with electrochemical detection 345(1985)203

Hydroxy fatty acids
Adsorption and reversed-phase HPLC of p-nitrobenzyl esters of monohydroxy fatty acids 329(1985)57

Qualitative and quantitative measurement of hydroxy fatty acids, thromboxanes and prostaglandins using stable isotope dilutions and detection by GC-MS 344(1985)11

5-Hydroxyflavones
Reversed-phase HPLC of 5-hydroxyflavones bearing tri- or tetrasubstituted A rings 347(1985)443

3-Hydroxyhexanoic acid
3-Hydroxyhexanoic acid: an abnormal metabolite in urine and serum of diabetic ketoacidotic patients 337(1985)1

4-Hydroxyhippuric acid
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

5-Hydroxyindole-3-acetic acid
3-Methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, and homovanillic acid in human cerebrospinal fluid. Storage and measurement by reversed-phase HPLC and coulometric detection using 3-methoxy-4-hydroxyphenyllactic acid as an internal standard 336(1985)259

Determination of urinary 5-hydroxyindole-3-acetic acid using solid-phase extraction and reversed-phase HPLC with electrochemical detection 341(1985)167

Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 345(1985)140

Hydroxyl aromatic compounds
Solubility parameter and window diagram optimization methods for the reversed-phase chromatographic optimization of complex hydroxyl aromatic mixtures 328(1985)81

Class separation of polycyclic aromatic hydrocarbons, nitrogen heterocycles and hydroxyl aromatics by LC 331(1985)55

2-[β-(4-Hydroxylphenylethyl)aminomethyl]tetralone
Rapid and simple purification of the ^{125}I-labeled α-1 adrenergic radioligand 2-[β-(4-hydroxylphenylethyl)aminomethyl] tetralone (BE 2254) using reversed-phase HPLC 338(1985)236

Hydroxyl-terminated stationary phases
New approach to capillary columns for GC? Condensation of hydroxyl-terminated stationary phases 347(1985)351

4-Hydroxymandelic acid
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

4-Hydroxymercuribenzoate
Analytical isoelectric focusing of native and modified haemoglobins after treatment with 4-hydroxymercuribenzoate 346(1985)401

7-Hydroxymethotrexate
Determination of serum methotrexate and 7-hydroxymethotrexate concentrations. Method evaluation showing advantages of HPLC 337(1985)81

4'-Hydroxy-3'-methoxynomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 341(1985)333

3'-Hydroxy-4'-methoxynomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 341(1985)333

4-Hydroxy-3-methoxyphenylpropionic acid
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

4-(4'-Hydroxy-1'-methylbutylamino)-7-chloroquinoline
Chloroquine metabolism in man: urinary excretion of 7-chloro-4-hydroxyquinoline and 7-chloro-4-aminoquinoline metabolites 345(1985)209

3-Hydroxymethyldibenzo[*b,f*]thiepin 5,5-dioxide
HPLC determination of 3-hydroxymethyl-dibenzo[*b,f*]thiepin 5,5-dioxide and its acid metabolite in human plasma and urine 338(1985)253

***p*-Hydroxymethylphenidate**
GC-MS analysis of methylphenidate and *p*-hydroxymethylphenidate using deuterated internal standards 343(1985)329

4-[(2-Hydroxynaphthalene-1-yl)azo]benzenesulphonic acid
Determination of the lower sulfonated subsidiary colors in FD&C Yellow No. 6 by reversed-phase HPLC 347(1985)163

6-Hydroxynicotinic acid
Availability of 6-hydroxynicotinic acid for rapid identification of *Pseudomonas aeruginosa* and *Serratia marcescens* 338(1985)51

4'-Hydroxynomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 341(1985)333

7-Hydroxyphenprocoumon
Identification of phenprocoumon metabolites in human urine by HPLC and GC-MS 338(1985)325

4-Hydroxyphenylacetic acid
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

4-Hydroxyphenylglycine
Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

4-Hydroxyphenyllactic acid
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

4-Hydroxyphenylpyruvic acid
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

N-(4-Hydroxyphenyl)retinamide
Simple HPLC method for the separation of retinoids including N-(4-hydroxyphenyl)-all-*trans*-retinamide 341(1985)187

Hydroxyphenytoin
GC determination of *m*- and *p*-hydroxyphenytoin in the urine of epileptic patients 339(1985)331

5-Hydroxy-N-(2-piperidylmethyl)-2-(2,2,2-trifluoroethoxy)benzamide
HPLC method for the quantitation of a *meta*-O-dealkylated metabolite of flecainide acetate, a new antiarrhythmic 343(1985)119

Hydroxyproline
Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1

Simplified procedure for specific determination of imino acids in human blood plasma by HPLC 339(1985)59

HPLC analysis of free hydroxyproline and proline in blood plasma and of free and peptide-bound hydroxyproline in urine 339(1985)285

4-Hydroxypropranolol
Rapid and simultaneous extraction of propranolol, its neutral and basic metabolites from plasma and assay by HPLC 343(1985)349

4'-Hydroxypropranolol sulphate
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 341(1985)213

3'-Hydroxysepiapterin
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

6β-Hydroxy-7α-thiomethylspirolactone
Determination of the serum concentration of spironolactone and its metabolites by HPLC 341(1985)279

5-Hydroxytryptamine
see Serotonin

5-Hydroxytryptophan
Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

25-Hydroxyvitamin D
Solid phase extraction system for vitamin D and its major metabolites in human plasma 343(1985)43

Hyodeoxycholic acid
Determination of bile acids in rat bile by HPLC 338(1985)195

Hyoscyamine
Comparison between TLC-densitometry and HPLC for the determination of hyoscyamine and hyoscyne in leaves, fruits and seeds of *Datura* (*Datura* spp.) 329(1985)415

Hyoscyne
Comparison between TLC-densitometry and HPLC for the determination of hyoscyamine and hyoscyne in leaves, fruits and seeds of *Datura* (*Datura* spp.) 329(1985)415

Hypercoulometry
Hypercoulometry in a pulsed conventional electron-capture detector 330(1985)87

Hypotaurine
Determination of cysteinesulfinate, hypotaurine and taurine in physiological samples by reversed-phase HPLC 343(1985)303

Hypoxanthine
Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Hypoxanthine-guanine phosphoribosyl transferase
Simultaneous HPLC assay of the activities of erythrocytic hypoxanthine-guanine phosphoribosyl transferase and purine nucleoside phosphorylase 349(1985)275

I

Ibuprofen
Improved HPLC assay of ibuprofen in plasma 344(1985)408

Imidazoleacetic acid
Measurement of imidazoleacetic acid in urine by GC-MS 343(1985)249

N-[4-(1H-Imidazol-1-yl)butyl]-2-(1-methylethyl)-11-oxo-11H-pyrido[2,1-*b*]quinazoline-8-carboxamide
Determination of the antiallergenic agent, N-[4-(1H-imidazol-1-yl)butyl]-2-(1-methylethyl)-11-oxo-11H-pyrido[2,1-*b*]quinazoline-8-carboxamide, in plasma by reversed-phase HPLC analysis using fluorometric detection 336(1985)301

Imidazotetrazines
Antitumour imidazotetrazines. VII. Quantitative analysis of mitozolomide in biological fluids by HPLC 337(1985)178

Imino acids
Simplified procedure for specific determination of imino acids in human blood plasma by HPLC 339(1985)59

HPLC analysis of free hydroxyproline and proline in blood plasma and of free and peptide-bound hydroxyproline in urine 339(1985)285

Imipramine
HPLC determination of imipramine and desipramine in human serum 336(1985)410

Immobilized pH gradients
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Immunoglobulin A
Chromatographic study of the interrelationships of immunoglobulin A and α_1-microglobulin in myelomatosis 327(1985)179

Immunoglobulin E
Isolation and quantitation of serum IgE levels by high-performance immunoaffinity chromatography 327(1985)205

Immunoglobulin G
Protein A-coated glass beads. Universal support medium for high-performance immunoaffinity chromatography 327(1985)213

Methods for chromatofocusing of cerebrospinal fluid and serum immunoglobulin G 327(1985)293

Immunoglobulin M
Application of DNA-cellulose chromatography in the isolation of immunoglobulin M and

complement component C4b-binding protein from human serum 347(1985)129

Immunoglobulins
Preparation of sheep and cattle immunoglobulins with antibody activity by HPLC 326(1985)225

Purification of human serum gamma globulins by hydrophobic interaction HPLC 326(1985)235

Immunosuppressives
Drug level monitoring: chromatography of some minor groups of drugs (Review) 340(1985)401

IMP dehydrogenase
High-performance liquid chromatographic detection of XMP as a basis for an improved IMP dehydrogenase assay 343(1985)160

Indecainide
LC determination of indecainide, a new antiarrhythmic drug, and its major metabolite, desisopropyl indecainide, in biological samples 337(1985)329

Indenestrol
Separation and detection of enantiomers of stilbestrol analogues by combined HPLC-thermospray MS 344(1985)378

Indole ethylamines
Derivatization in aqueous solution, isolation and separation of tetrahydro-β-carbolines and their precursors by LC 341(1985)287

Indoprofen
Determination of the enantiomers of indoprofen in blood plasma by HPLC after rapid derivatization by means of ethyl chloroformate 339(1985)339

Injection peak
Occurrence and origin of system peaks in non-suppressed ion chromatography of inorganic anions with indirect UV absorption detection 346(1985)125

Injection techniques
Rejection of the sample plug after on-column injection into capillaries at high temperatures 333(1985)156

On-column injection of large sample volumes using the retention gap technique in capillary GC (Review) 334(1985)129

Improved direct injection method and extractive methylation method for determination of valproic acid in serum by GC 342(1985)89

Injection volume
HPLC theory for the practitioner (Review) 334(1985)197

Trace analysis of anions by use of a back-flush method and large injection volumes in ion chromatography 350(1985)317

Injectors
GC retention indices of chemical warfare agents and simulants 331(1985)47

Inlet pressure
A problem of optimization in capillary GC 348(1985)416

Inorganic anions
Recent developments in isotachophoresis (Review) 334(1985)157

Occurrence and origin of system peaks in non-suppressed ion chromatography of inorganic anions with indirect UV absorption detection 346(1985)125

Pentafluorobenzyl *p*-toluenesulphonate as a new derivatizing reagent for GC determination of anions 346(1985)215

Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Inosine
Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 339(1985)75

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 345(1985)457

Inosine 5'-monophosphate
High-performance liquid chromatographic detection of XMP as a basis for an improved IMP dehydrogenase assay 343(1985)160

Inositols
Identification of inositols and their mono-O-methyl ethers by GLC 333(1985)167

Insulin
Reversed-phase HPLC of insulin and insulin derivatives. A comparative study 348(1985)347

Identification and isolation of human insulin A and B chains by HPLC 349(1985)91

Insulin-like growth factor inhibitor
Partial purification of a specific inhibitor of the insulin-like growth factors by reversed-phase HPLC 336(1985)87

Interfaces
Versatile HPLC computer system for solvent delivery, gradient control and data acquisition using the Commodore 64 personal computer 333(1985)301

Interfaces, LC-MS
Direct liquid introduction MS of some underivatized dipeptides and polypeptides 350(1985)399

Interfacial adsorption
Contributions to hydrocarbon retention in gas-liquid-solid chromatography with polyethylene glycol 6000 as stationary phase 350(1985)15

Interferometry
Application of interferometry in studies of transport processes near ionites and adsorbents 333(1985)198

Interferon
Automated high-performance immunosorbent assay for recombinant leukocyte A interferon 327(1985)189

Analysis of different forms of recombinant human leukocyte interferons and synthetic fragments by HPLC 327(1985)359

Interleukin-1
HPLC separation of distinct epidermal cell-derived cytokines 326(1985)129

Interleukin-2
Characterization of recombinant human interleukin-2 with micromethods 326(1985)357

Interleukin-3
HPLC separation of distinct epidermal cell-derived cytokines 326(1985)129

Internal standards
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

Interstrand distance
Effect of interstrand distance upon chiral recognition by a chiral stationary phase 328(1985)1

Inulin
Application of GPC systems to the determination of the molecular weight of inulin 348(1985)187

Ion exchangers
Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Separation of peptides by strong cation-exchange HPLC 327(1985)147

Application of a new ion exchanger TSK-GEL DEAE-5PW, to the purification of Cu,Zn-superoxide dismutase of bovine erythrocytes 327(1985)301

Preparative high-performance ion-exchange chromatography 333(1985)29

Performance characteristics of some commercially available low-capacity anion-exchange columns suitable for non-suppressed ion chromatography 346(1985)139

Preparation of low-capacity anion-exchange resins for ion chromatography on a methacrylic copolymer matrix 350(1985)296

Ionites
Application of interferometry in studies of transport processes near ionites and adsorbents 333(1985)198

Ionization constants
Ionization constants and lipophilicity of 1-arylpiperazines investigated by reversed-phase HPLC 333(1985)191

Iothalamate
Quantitation of iothalamate in serum and urine by HPLC 337(1985)434

Ipratropium bromide
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Iron
Simultaneous determination of iron(II) and -(III) by ion chromatography with post-column reaction 329(1985)247

Irritants
GC retention indices of chemical warfare agents and simulants 331(1985)47

Isoalkyl acrylates
Chromatography of monomers. IV. GLC studies of C_1-C_6 *n*-alkyl and C_3-C_6 isoalkyl acrylates and their hydrogen halide and halogen addition derivatives 348(1985)141

Isobergapten
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Isobutylamine
GC-MS analysis of volatile amines produced by several strains of *Clostridium* 337(1985)213

2-(4-Isobutylphenyl)propionic acid
Determination of 2-(4-isobutylphenyl)propionic acid in bulk drug and compressed tablets by reversed-phase HPLC 331(1985)169

Isocyanate-based copolyamide resins
GC analysis of isocyanate-based copolyamide resins 329(1985)132

Isocyanates
Trace analysis of amines and isocyanates using glass capillary GC and selective detection. V. Direct determination of isocyanates using nitrogen-selective and electron-capture detection 331(1985)331

Capillary GC method for the determination of complex mixture of isocyanates and amines 346(1985)191

Isoelectric focusing
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Adaptation of the equipment for high-performance electrophoresis to isoelectric focusing 346(1985)265

Isoetharine
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Isoflavones
Analysis of isoflavones in *Puerariae radix* by HPLC with amperometric detection 347(1985)438

3-O-α-Isomaltosyl-D-glucose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Isomaltotriose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Isonamid-7030
GC analysis of isocyanate-based copolyamide resins 329(1985)132

Isoniazid
Determination of isoniazid, actylisoniazid, acetylhydrazine and diacetylhydrazine in biological fluids by HPLC 338(1985)113

Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Ion-pair HPLC determination of isoniazid and acetylisoniazid in plasma and urine. Application for acetylator phenotyping 341(1985)193

Isopanose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Isophephrine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Isopimpinellin
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Isoplast
GC analysis of isocyanate-based copolyamide resins 329(1985)132

***d*-Isoprenaline**
Reversed-phase HPLC and amperometric detection of 3-O-methyl isoprenaline sulphate: application to studies on the presystemic metabolism of *d*-isoprenaline in man 337(1985)311

Isoproterenol
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Isoptelefolonium
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Isoputreanine
Determination of polyamines and related compounds by reversed-phase HPLC: improved separation systems 339(1985)45

Isosorbide-5-mononitrate
GLC determination of isosorbide-5-mononitrate derivatized with perfluorinated anhydrides for pharmacokinetic and bioavailability investigations 345(1985)390

Isotachophoresis
Automation of analytical isotachophoresis (Review) 334(1985)83

Recent developments in isotachophoresis (Review) 334(1985)157

Isotachophoresis as a useful tool for monitoring neurological complications of acute leukaemia in children 342(1985)285

Isotope effects
Separation factors for the system 1,2-dibromoethane-1,2-dibromo[$^2H_4$]ethane 328(1985)333

Isoxicam
LC assay of isoxicam in human plasma and urine 337(1985)156

J

Janthitrems
TLC of mycotoxins (Review) 334(1985)211

Jasmonic acid
Simultaneous determination of abscisic acid and jasmonic acid in plant extracts using HPLC 330(1985)347

Jervine
Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

Josamycin
Determination of josamycin in plasma, erythrocytes and leucocytes via HPLC by direct injection of plasma or homogenized blood cells, using alternating sample clean-up columns 344(1985)416

K

Kahweol
Use of silver nitrate impregnated silica cartridges in the separation of kahweol and cafestol esters by preparative LC 328(1985)422

Kanamycin
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Karaya gum
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Kerosene
GC determination of total aromatics in kerosene 328(1985)207

1-Kestose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Ketamine
Quantitative deconvolution of heavily fused chromatographic peaks of biological components using a multi-wavelength UV detector 332(1985)139

Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Ketanserin
Simultaneous determination of ketanserin and ketanserinol in biological fluids using ion-pair LC and fluorometric detection 341(1985)208

Ketanserinol
Simultaneous determination of ketanserin and ketanserinol in biological fluids using ion-pair LC and fluorometric detection 341(1985)208

Keto acids
Quantification of branched-chain α-keto acids as quinoxalinols: importance of excluding oxygen during derivatization 337(1985)98

GC determination of serum branched-chain α-keto acids derivatized by extractive alkylation 338(1985)265

Determination of α-keto acids in serum and urine by HPLC with fluorescence detection 344(1985)33

Ketones
Determination of low concentrations of low-molecularweight aldehydes and ketones in aqueous samples 329(1985)81

Determination of aldehydes and ketones by derivatization and LC-MS 333(1985)337

Ketoprofen
Highly sensitive determination of ketoprofen in human serum and urine and its application to pharmacokinetic study 345(1985)419

6-Keto-prostaglandin $F_{1\alpha}$
Simultaneous determination of prostanoids in plasma by GC-negative-ion chemical-ionization MS 338(1985)273

Assay methods for 6-keto-prostaglandin $F_{1\alpha}$ in human urine. Comparison of chromatographic techniques with radioimmunoassay and GC-negative-ion chemical-ionization MS 339(1985)1

Ketotifen
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

17-Ketotrilostane
HPLC assay for trilostane and its major metabolite, 17-ketotrilostane, in human plasma 339(1985)440

Kokusaginine
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Kraft digestion
GC and MS identification of sulphur gases in kraft digestion plants 346(1985)405

Ky-5
Synthesis and GC-MS determination of polychlorinated dibenzo-*p*-dioxins and related compounds in the technical chlorophenol formulation Ky-5 331(1985)410

L

Labetalol
Convenient GC method for the determination of the proportions of the two racemic modifications present in labetalol hydrochloride 333(1985)178

Measurement of labetalol by HPLC with electrochemical detection 344(1985)412

β-Lactamases
High-performance liquid affinity chromatography of β-lactamase 326(1985)163

Use of chromatofocusing for separation of β-lactamases. V. Inducible chromosomally mediated β-lactamase of the *Enterobacter cloacae* 53 strain 333(1985)239

Use of chromatofocusing for separation of β-lactamases. VI. Comparative studies on two chromatofocusing polybuffer exchanger matrices with the R46 plasmid coded β-lactamase carried by the *Escherichia coli* K12 J5-3 strain 333(1985)244

Lactate dehydrogenase
Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Lactogen
HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25

Lacto-N-hexaose
Characterization of lacto-N-hexaose and two fucosylated derivatives from human milk by HPLC and proton NMR spectroscopy 328(1985)259

4-O-β-Laminaribiosyl-D-glucose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Laminaritriose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Lanthanide β-diketonates
Selectivity of complexing GC stationary phases containing lanthanide β-diketonates 349(1985)457

Lanthanide ions
Lanthanide ions as luminescent chromophores for LC detection 328(1985)101

Lanthanides
Determination of trace amounts of lanthanides in rocks and minerals by HPLC 349(1985)137

Laser-induced fluorescence detection
Performance of some cell designs for laser-induced fluorescence detection in open-tubular LC 346(1985)149

Laudanosine
Determination of atracurium, laudanosine and related compounds in plasma by HPLC 343(1985)431

Lecithin
Estimation of amniotic fluid phospholipids by HPLC 336(1985)139

Lectins
Anomalous behavior of lectins in size-exclusion HPLC and gel electrophoresis 336(1985)397

Studies on lectins. LVIII. Sugar-binding properties, as determined by affinity electrophoresis, of α-D-galactosidases from *Vicia faba* seeds possessing erythroagglutinating activity 347(1985)343

Lenperone
Determination of lenperone in dog plasma by HPLC 345(1985)430

Leucine
GC of amino acids in urine and haemofiltrate 345(1985)19

Leucine enkephalin
Improved detection limits in the analysis of tyrosine-containing polypeptide hormones by using electrochemical detection 327(1985)165

Leukotrienes
Quantitation of sulfidopeptide leukotrienes by reversed-phase HPLC 343(1985)213

Leupeptin
HPLC method for monitoring leupeptin in mouse serum and muscle by pre-column fluorescence derivatization with benzoin 345(1985)259

Levamisole
Rapid Extrelut column method for determination of levamisole in milk using HPLC 337(1985)151

Levobunolol
Determination of levobunolol and dihydrolevobunolol in blood and urine by HPLC using fluorescence detection 338(1985)444

Levorphanol
Quantitation of levorphanol in plasma using HPLC with electrochemical detection 341(1985)239

Lidocaine
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Lignin
High-performance size-exclusion chromatography of lignin and its derivatives 328(1985)299

Lignosulphonates
High-performance size-exclusion chromatography of lignosulphonates 331(1985)419

Lime oil
General method for the HPLC prefractionation of essential oils and flavor mixtures for GC-MS analysis. Identification of new constituents in cold pressed lime oil 330(1985)141

Linoleic acid hydroperoxides
HPLC of linoleic acid hydroperoxides and their corresponding alcohol derivatives 350(1985)445

Linolenic acid hydroperoxide
HPLC as a tool for identification of linolenic acid hydroperoxide prepared with soybean lipoxygenase 350(1985)435

Lipid associating peptides
Purification of synthetic lipid associating peptides and the monitoring of the deformylation of N^{in}-formyltryptophan by reversed-phase HPLC 347(1985)237

Lipids
HPLC technique for the separation of lipid classes 329(1985)273

Assay of lipids in dog myocardium using capillary GC and derivatization with boron trifluoride and methanol 339(1985)25

Effect of copper(II) sulphate impregnation of Chromarods on the sensitivity of the Iatroscan detection system to lipids 346(1985)468

HPLC determination of lipids in vesicles 348(1985)221

Lipophilicity
see also Hydrophobicity

Ionization constants and lipophilicity of 1-arylpiperazines investigated by reversed-phase HPLC 333(1985)191

TLC and HPLC in the evaluation of the lipophilicity of aryloxoalkanoic and arylhydroxyalkanoic acids 347(1985)335

Interaction of some non-ionic tensides with dioleoyl phosphatidyl choline, studied by charge-transfer chromatography 349(1985)295

Lipopolysaccharides
Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 345(1985)119

Lipoproteins
HPLC of apolipoproteins in serum high-density lipoproteins 336(1985)151

Rapid separation of plasma lipoproteins by GPC on agarose gel Superose 6B 341(1985)154

Evaluation of immunoaffinity chromatography for isolating human lipoproteins containing apolipoprotein B 342(1985)47

Use of reversed-phase HPLC to analyse the cholesterol ester profiles present in lipoproteins of patients exhibiting hyperalphalipoproteinaemia or hypoalphalipoproteinaemia (Tangier disease) 343(1985)149

Lipoxidase
Preparative high-performance ion-exchange chromatography 333(1985)29

Liquefied gases
Liquefied gases as eluents for TLC 347(1985)362

Liquid chromatography-mass spectrometry
Direct coupling of micro HPLC with fast atom bombardment MS 346(1985)161

Gaseous acidity and basicity scales as guides to chemical ionization in reversed-phase LC-MS with direct liquid introduction 347(1985)61

Analysis of candidate anticancer drugs by thermospray HPLC-MS and by high-resolution MS 347(1985)137

Direct liquid introduction MS of some underivatized dipeptides and polypeptides 350(1985)399

Liquid crystals
New liquid crystalline stationary phases for GC of positional and geometrical isomers having similar volatilities 331(1985)69

Correlations between retention on liquid crystalline stationary phases and chemical structure. I. Dimethylnaphthalenes 333(1985)11

Liquid scintillation detector
Coupling a high-performance liquid chromatograph with a liquid scientillation detector: example in the field of pyrethroids 349(1985)105

Lithocholic acid
Determination of bile acids in rat bile by HPLC 338(1985)195

Rapid and accurate reversed-phase HPLC determination of conjugated bile acids in human bile for routine clinical applications. Therapeutic control during gallstone dissolution therapy 339(1985)263

Loading techniques
Bracketing, a simple loading technique that increases sample recovery, improves resolution, and increases sensitivity in HPLC 348(1985)123

Ion-exchange columns in HPLC of proteins; a simple loading technique that improves resolution 350(1985)299

Loflazepate
Determination of circulating ethyl loflazepate metabolites in the baboon by radio-HPLC with injection of crude plasma samples: comparison with solvent extraction and TLC 342(1985)159

Low-angle laser light-scattering photometry
Size-exclusion chromatography-low-angle laser light-scattering photometry of lignite macromolecules 349(1985)337

Loxoprofen
Column LC for the simultaneous determination of the enantiomers of loxoprofen sodium and its metabolites in human urine 345(1985)373

Luteinizing hormone-releasing hormone
Improved detection limits in the analysis of tyrosine-containing polypeptide hormones by using electrochemical detection 327(1985)165

Lutidine
GC separation of 2,6-dimethylpyridine and its chloro derivatives 347(1985)393

Lysergic acid diethylamide
HPLC system for the separation of ergot alkaloids with applicability to the analysis of illicit lysergide (LSD) 346(1985)423

Lysergide
HPLC system for the separation of ergot alkaloids with applicability to the analysis of illicit lysergide (LSD) 346(1985)423

Lysophospholipids
Reversed-phase HPLC separation of molecular species of alkyl ether, vinyl ether, and monoacyl lyosphospholipids 338(1985)61

Lysozyme
Comparison of reversed-phase columns for the separation of tryptic peptides by gradient elution. Correlation of experimental results and model prediction 326(1985)419

Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269

M

α_2-Macroglobulin
Evidence for the degradation of sugars during the resin hydrolysis of glycoproteins 339(1985)182

Macrolides
Antibiotic monitoring in body fluids (Review) 340(1985)361

Macromolecules
A new, stabilized, hydrophilic silica packing for the high-performance gel chromatography of macromolecules 326(1985)63

Affinity chromatographic examination of a retention model for macromolecules 331(1985)1

Magnesium
Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. I. Cation binding in the mobile phase 329(1985)321

Improvement of ion chromatography with UV photometric detection and comparison with conductivity detection for the determination of serum cations 341(1985)53

Malotilate
Determination of malotilate and its metabolites in plasma and urine 342(1985)349

Maltooligosaccharides
Ion-exchange chromatography of fluorogenic derivatives of maltooligosaccharides for preparation of α-amylase substrates 336(1985)368

Maltotriose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Mandelic acid
Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59

Determination of mandelic and phenylglyoxylic acids in rat urine by HPLC and by isotachophoresis 343(1985)175

Mass flow control system
Constant mass flow system for GC 330(1985)172

Mass spectrometry-mass spectrometry
Some results from an investigation into the analytical potential of MS-MS techniques, with particular reference to industrial applications 328(1985)167

Mass transfer coefficients
Reversed-flow GC technique applied to measurement of mass transfer coefficients of *n*-hydrocarbons on Porapak P 348(1985)339

MATRIXCOMP
Microcomputer programming in BASIC for the evaluation of capillary GC in the analysis of pesticide residues. I. MATRIXCOMP: a program that facilitates the recognition of interfering peaks from the biological matrix 349(1985)49

Matrix effects
Influence of the biological matrix on retention behaviour in TLC: evidence of systematic differences between pure and extracted drugs 342(1985)241

Mebeverine
Reversed-phase HPLC of mebeverine hydrochloride and related compounds 350(1985)135

Medroxyprogesterone acetate
Simple HPLC method for the determination of medroxyprogesterone acetate in human plasma 341(1985)437

Specific and sensitive determination of medroxyprogesterone acetate in human serum by GC-MS 345(1985)1

Mefenamic acid
Analysis of antipyretics by semimicro LC 332(1985)269

Mefloquine
GC determination of mefloquine in human and dog plasma using electron-capture detection 337(1985)47

Melamine
HPLC determination of melamine extracted from cups made of malemine resin 346(1985)450

Melezitose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Melphalan
Modified extraction and chromatography for the measurement of plasma melphalan by ion-pair HPLC 336(1985)429

HPLC analysis of melphalan in plasma, brain and peripheral tissue by *o*-phthalaldehyde derivatization and fluorescence detection 339(1985)434

Drug level monitoring: cytostatics (Review) 340(1985)31

Membrane filtration
Combined recycling affinity column-filtration technique 346(1985)247

Membrane glycoproteins
Tandem separation of labelled human blood platelet membrane glycoproteins by anion-exchange and gel fast protein LC 326(1985)179

Membrane proteins
Separation of flavivirus membrane and capsid proteins by multistep HPLC optimized by immunological monitoring 326(1985)113

Isolation of detergent-solubilized monomers of bacteriorhodopsin by size-exclusion HPLC 326(1985)243

Rapid delipidation of the membrane protein bacteriorhodopsin by HPLC 350(1985)427

Membranes
Affinity partitioning of biopolymers and membranes in Ficoll-dextran aqueous two-phase systems 331(1985)11

Membrane-stabilizing agents
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Meperidine
Analysis of meperidine and normeperidine in serum and urine by HPLC 338(1985)141

Mephenytoin
Determination of antiepileptic drugs (Review) 340(1985)199

Meprobamate
Determination of meprobamate as an *n*-butylboronate ester derivative in serum by GLC 341(1985)462

Meptazinol
Measurement of meptazinol in plasma by HPLC with electrochemical detection 341(1985)223

Mercaptalbumin
HPLC studies on non-mercapt mercapt conversion of human serum albumin II 332(1985)19

Ion-exchange HPLC studies on sulphydryl-catalysed structural alterations of bovine mercaptalbumin 332(1985)29

Mercaptoethanol
1-DOPA decarboxylase and dopamine-β-hydroxylase isolated from sympathetic splenic nerves estimated by sodium dodecyl sulphate polyacrylamide gel electrophoresis with and without mercaptoethanol 329(1985)300

GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

2-Mercaptopropionic acid
Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193

6-Mercaptopurine
Sensitive HPLC determination of 6-mercaptopurine, 6-thioguanine, 6-mercaptopurine riboside and 6-thioguanosine in biological fluids 336(1985)422

Drug level monitoring: cytostatics (Review) 340(1985)31

Assay of 6-mercaptopurine in human plasma 345(1985)441

6-Mercaptopurine riboside
Sensitive HPLC determination of 6-mercaptopurine, 6-thioguanine, 6-mercaptopurine riboside and 6-thioguanosine in biological fluids 336(1985)422

Mesh-size distribution
Effect of the mesh-size distribution of porous polymer beads on efficiency and selectivity of stationary phases 349(1985)189

Metal chelates
HPLC of copper, cobalt, nickel, chromium and vanadium chelates with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol 333(1985)133

Applications of fused-silica wall-coated open-tubular columns. I. Schiff base chelates 350(1985)353

Metal complexes
Phenol effect in PC of metal complexes 331(1985)273

Metal ions
Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147

Metalloendopeptidase
Purification of gelatin-specific neutral protease from human skin by conventional and high-performance LC 326(1985)147

Metapramine
Simultaneous determination of metapramine and its dimethylated metabolites in plasma by GC-MS 345(1985)59

Metaproterenol
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Metaraminol
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Methadone
HPLC analysis of methadone hydrochloride in pharmaceuticals 329(1985)438

Methanesulphonatobicyclononene carbonitriles
Separation of isomeric halogenobicyclononene carbonitriles 347(1985)297

Methanethiol
GC and MS identification of sulphur gases in kraft digestion plants 346(1985)405

Methanol
HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Methionine
GC of amino acids in urine and haemofiltrate 345(1985)19

Methionine sulphone
Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1

Methotrexate
Determination of serum methotrexate and 7-hydroxymethotrexate concentrations. Method evaluation showing advantages of HPLC 337(1985)81

Drug level monitoring: cytostatics (Review) 340(1985)31

HPLC assay for methotrexate utilizing a cold acetonitrile purification and separation of plasma or cerebrospinal fluid 342(1985)424

Determination of the antineoplastic agent methotrexate in body fluids by HPLC with electrochemical detection 344(1985)249

2-[3'-(Methoxycarbonylamino)-phenyl]-3-phenyl-6-methoxycarbonylamino-4-(3H)-quinazolone
HPLC determination of 2-[3'-(methoxycarbonylamino)-phenyl]-3-phenyl-6-methoxycarbonylamino-4-(3H)-quinazolone (NSC-251635) in human serum 339(1985)445

3-Methoxy-4-hydroxyphenylethylene glycol
3-Methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, and homovanillic acid in human cerebrospinal fluid. Storage and measurement by reversed-phase HPLC and coulometric detection using 3-methoxy-4-hydroxyphenyllactic acid as an internal standard 336(1985)259

Assay of 3-methoxy-4-hydroxyphenylglycol in human plasma using HPLC with amperometric detection 338(1985)396

Routine determination of unconjugated 3-methoxy-4-hydroxyphenylglycol in plasma using HPLC with electrochemical detection 341(1985)160

Determination of low plasma concentrations of 3-methoxy-4-hydroxyphenylethylene glycol using GC-negative-ion chemical ionization MS 343(1985)239

Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

2-[2-Methoxy-4-(methylsulphinyl)phenyl]-1H-imidazo-[4,5-*c*]pyridine hydrochloride
HPLC assay of 2-[2-methoxy-4-(methylsulfinyl)phenyl]-1H-imidazo-[4,5-*c*]pyridine hydrochloride, a potent cardiotonic agent, and two metabolites in plasma 342(1985)228

(+)-3-Methoxymorphinan
Determination of dextromethorphan and metabolites in human plasma and urine by HPLC with fluorescence detection 338(1985)99

Methoxynaphthalene
New liquid crystalline stationary phases for GC of positional and geometrical isomers having similar volatilities 331(1985)69

8-Methoxypsoralen
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

5-Methoxysalicylic acid
HPLC determination of certain salicylates and their major metabolites in plasma following topical administration of a liniment to healthy subjects 342(1985)315

3-Methoxy-4-sulphooxyphenylethylene glycol
Quantitative analysis of (3-methoxy-4-sulphooxyphenyl)ethylene glycol (MHPG sulphate) in human urine 337(1985)441

Quantitative analysis of (3-methoxy-4-sulphooxyphenyl)ethylene glycol (MHPG sulphate) in human urine. Reply to Murray and Davies 337(1985)443

3-Methoxytyramine
Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

6α-Methyl-17α-acetoxyprogesterone
Reversed-phase HPLC with isocratic elution of 6α-methyl-17α-acetoxyprogesterone (MAP) and its impurities 330(1985)415

Methylalkanes
Kováts retention indices of high-molecular-weight monomethyl-, cyclopentyl-, cyclohexyl- and phenylalkanes 331(1985)219

Methylamino acids
N-Methylation and N,N-dimethylation of amino acids. An artifact production in the analysis of organic acids using diazomethane as derivatizing agent 338(1985)33

5-(Methylaminosulphonyl)-N-[(1-allyl-2-pyrrolidinyl)methyl]-2-methoxy-4-aminobenzamide
HPLC determination of a new benzamide in biological fluids for use in a human pharmacokinetic study 343(1985)443

Methylaniline
Measurement of trace aromatic amines in seawater using HPLC with electrochemical detection 348(1985)265

Methylated cyclodextrins
Methylated cyclodextrins as mobile phase additives in LC 330(1985)253

3-Methylbutylamine
GC-MS analysis of volatile amines produced by several strains of *Clostridium* 337(1985)213

3-Methylclonazepam
Plasma determination of 3-methylclonazepam by capillary GC 338(1985)347

S-Methylcysteine
Reversed-phase HPLC separation and quantitation of phenylthiohydantoin derivatives of 25 amino acids, including those of cysteic acid, 4-hydroxyproline, methionine sulfone, S-carboxymethylcysteine and S-methylcysteine 327(1985)1

A method for the determination of S-methylcysteine by HPLC: application to the study of carcinogenic methylating agents 329(1985)202

Methyldopa
Simultaneous determination of L-dopa and 3-O-methyldopa in human serum by HPLC 336(1985)380

Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

N-Methyldopamine
HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61

Methylene blue
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Methylene chloride
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233

***cis*-9,10-Methylene-hexadecanoate**
HPLC separation of fatty acids as pentafluorobenzyl esters 336(1985)115

Methyl enol ethers
Methyl enol ethers as artefacts in capillary GC profiles of aldehyde dimethyl acetals 345(1985)219

3-Methyl-7-ethyl-9α-ol-(4-chlorobenzoyloxy)3,7-diazabicyclo(3.3.1)-nonane monohydrochloride
HPLC method for quantitative determination of Yutac, a new antiarrhythmic agent, in dog plasma 338(1985)428

Methylfurylbutyrolactone
Investigation of operating parameters in high-performance displacement chromatography 330(1985)1

Methylguanidine
HPLC analysis of guanidino compounds using ninhydrin reagent. II. Guanidino compounds in blood of patients on haemodialysis therapy 342(1985)269

Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Methylguanosine
Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 339(1985)75

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Methylhippuric acid
Determination of methylhippuric acid in human urine by HPLC and by isotachophoresis 343(1985)419

1-Methylhistamine
Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Methylhistidine
GC quantitative determination of 1- and 3-methylhistidine in urine and muscles: comparison with glass capillary determination 336(1985)356

6-Methyl-2-(2-hydroxyethyl)pyridine
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

1-Methylinosine
Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 339(1985)75

3-O-Methyl isoprenaline sulphate
Reversed-phase HPLC and amperometric detection of 3-O-methyl isoprenaline sulphate: application to studies on the presystemic metabolism of *d*-isoprenaline in man 337(1985)311

Methyl methacrylate
Assay of methyl methacrylate in blood samples by headspace capillary GC 345(1985)386

(+)-17-Methylmorphinan-3-ol
Determination of dextromethorphan and metabolites in human plasma and urine by HPLC with fluorescence detection 338(1985)99

Methylnaphthalenes
α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131

N-Methylnicotinium
HPLC with electrochemical detection for the determination of nicotine and N-methylnicotinium ion 347(1985)405

Methyl paraoxon
Determination of methyl paraoxon in dog plasma by reversed-phase HPLC 338(1985)389

Methylphenidate
Enantiomeric resolution of *dl-threo*-methylphenidate, U.S.P. (Ritalin), by HPLC 328(1985)378

GC-MS analysis of methylphenidate and *p*-hydroxymethylphenidate using deuterated internal standards 343(1985)329

Methylphosphine
Quantitative GC analysis of low-molecular-weight alkylphosphines in the presence of phosphine and hydrocarbons 330(1985)400

10-(4-Methylpiperazino)-10,11-dihydrodibenzo[*b,f*]thiepin
Relationships between the chromatographic behaviour and structure of some substituted dibenzo[*b,f*]thiepins and their analogues 329(1985)113

Methylprednisolone
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Rapid method for the measurement of methylprednisolone and its hemisuccinate in plasma and urine following "pulse therapy" by HPLC 342(1985)251

HPLC assay for methylprednisolone and its soluble prodrug esters in dog plasma 344(1985)41

(15*R*)-15-Methylprostaglandin E_2
Determination of (15*R*)- and (15*S*)-15-methylprostaglandin E_2 in human plasma with picogram per milliliter sensitivity by column-switching HPLC 343(1985)271

Methyltestosterone
Quantitative determination of methyltestosterone and methyltestosterone-d_3 in serum by GC-MS 338(1985)281

5-Methyltetrahydrofolic acid
Stabilization of 5-methyltetrahydrofolic acid and subsequent analysis by reversed-phase HPLC 328(1985)417

6-Methylthioguanine
HPLC assay of 6-methylthioguanine, a major metabolite of 6-thioguanine, in urine 337(1985)321

3-Methylthymidine
Cinnamoylation of the sugar hydroxyls of 3-methylthymidine. Formation of a relatively stable ester derivative 347(1985)99

O-Methyltransferases
HPLC of proteins. Rapid chromatofocusing of plant enzymes 348(1985)167

Metipranolol
Determination of ophthalmic therapeutic metipranolol and its degradation product by reversed-phase HPLC 348(1985)304

Metoprolol
Simultaneous determination of metoprolol and deuterium-labelled metoprolol in human plasma by GC-negative-ion MS 339(1985)404

Simple and rapid analysis of atenolol and metoprolol in plasma using solid-phase extraction and HPLC 339(1985)429

Quantitative analysis of metoprolol and three of its metabolites in urine and liver microsomes by HPLC 342(1985)199

Enantiomer separation of metoprolol and its analogues and metabolites by capillary column

GC after derivatization with phosgene 350(1985)328

Mexiletine
HPLC assay for mexiletine enantiomers in human plasma 337(1985)172

LC determination of mexiletine and tocainide in human plasma with fluorescence detection after reaction with a modified *o*-phthalaldehyde reagent 344(1985)221

Micellar chromatography
Determination of drugs in untreated body fluids by micellar chromatography with fluorescence detection 342(1985)293

Microbore columns
see also Capillary columns and columns

Advanced microcolumn techniques in LC. Use of glass-lined stainless-steel tubing columns 332(1985)3

Polyallylamine-coated silica gel microbore column for LC 332(1985)15

On-column preparation of chemically bonded stationary phase with maximum surface coverage and high reproducibility, and its application to packed microcapillary columns 332(1985)57

α_1-Microglobulin
Chromatographic study of the interrelationships of immunoglobulin A and α_1-microglobulin in myelomatosis 327(1985)179

Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269

β_2-Microglobulin
Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269

Efficient purification of human plasma β_2-microglobulin from the haemodialysate of a patient with chronic renal failure by use of HPLC 339(1985)67

Micropacked columns
Contribution to the study of micropacked columns in GC 348(1985)327

Midazolam
Determination of midazolam (Versed) and its metabolites in plasma by HPLC 344(1985)199

Mitindomide
Analysis of candidate anticancer drugs by thermospray HPLC-MS and by high-resolution MS 347(1985)137

Mitomycin
Drug level monitoring: cytostatics (Review) 340(1985)31

HPLC assay of mitomycin in biological fluids 345(1985)197

Mitoxantrone
Ion-paired HPLC determination of mitoxantrone in physiological fluids 337(1985)73

Detection and separation of mitoxantrone and its metabolites in plasma and urine by HPLC 342(1985)119

Mitozolomide
Antitumour imidazotetrazines. VII. Quantitative analysis of mitozolomide in biological fluids by HPLC 337(1985)178

Mobile phase composition
Solubility parameter and window diagram optimization methods for the reversed-phase chromatographic optimization of complex hydroxyl aromatic mixtures 328(1985)81

Selection of mobile phases for reversed-phase LC with on-line electron-capture detection 328(1985)93

Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

Methylated cyclodextrins as mobile phase additives in LC 330(1985)253

Application of micro HPLC to the separation of chiral amino acids 331(1985)99

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. II. Influence of organic co-solvents on retention at 25°C 333(1985)41

Some applications of reversed-phase HPLC to oligosaccharide separations 336(1985)161

Computer-aided mobile phase optimization and chromatogram simulation in HPLC (Review) 338(1985)1

Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 339(1985)75

Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370

Ion-exchange HPLC of mouse epidermal growth factor and its congeners: mobile phase optimization with ion-pairing additives 345(1985)168

Influence of the mobile phase composition on the reversed-phase TLC behaviour of a series of prostaglandins. Comparison of the extrapolated R_M values 347(1985)209

Liquefied gases as eluents for TLC 347(1985)362

Reversed-phase HPLC of insulin and insulin derivatives. A comparative study 348(1985)347

α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131

HPLC determination of contaminants in salicylic acids after computer-assisted optimization of the separation parameters (in German) 350(1985)471

Modifier gradients
Gradient programming and combined gradient-pressure programming in supercritical fluid chromatography 349(1985)317

Molar refractivities
Relationships between GLC behaviour and structure of arylaliphatic acids 333(1985)21

Molecular imprinting
Molecular imprinting of amino acid derivatives in macroporous polymers. Demonstration of substrate- and enantio-selectivity by chromatographic resolution of racemic mixtures of amino acid derivatives 347(1985)1

Use of silane monomers for molecular imprinting and enzyme entrapment in polysiloxane-coated porous silica 347(1985)11

Molecular sieve 13X
Sorption properties of some modified molecular sieves 13X towards thiophene and benzene 329(1985)128

Molecular weight determination
Determination of molecular weights of coal derived liquids by GPC-HPLC 329(1985)65

Anomalous behavior of lectins in size-exclusion HPLC and gel electrophoresis 336(1985)397

GPC with interference refractometry for the rapid assay of polydisperse dextrans in biological fluids 337(1985)29

Confidence limits for a gel-permeation system 341(1985)420

Application of GPC systems to the determination of the molecular weight of inulin 348(1985)187

Size-exclusion chromatography-low-angle laser light-scattering photometry of lignite macromolecules 349(1985)337

Determination of molecular weight data in size-exclusion chromatography by use of braod-molecular-weight-distributed standards 349(1985)357

Molecular weight standards
Determination of molecular weight data in size-exclusion chromatography by use of braod-molecular-weight-distributed standards 349(1985)357

Monoacetyldapsone
Rapid and sensitive LC method for the determination of dapsone and monoacetyldapsone in plasma and urine 345(1985)447

Monoacetylputrescine
Determination of polyamines and related compounds by reversed-phase HPLC: improved separation systems 339(1985)45

Monoacetylspermidines
Determination of polyamines and related compounds by reversed-phase HPLC: improved separation systems 339(1985)45

Monoacyl lysophospholipids
Reversed-phase HPLC separation of molecular species of alkyl ether, vinyl ether, and monoacyl lysophospholipids 338(1985)61

Monoalkyl-*sn*-glycerols
Diastereomeric separations of natural glycero derivatives as their 1-(1-naphthyl)ethyl carbamates by HPLC 350(1985)417

Monoclonal light chains
Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269

Monoethylglycylmesidide
Determination of trimecaine metabolites in blood plasma by capillary isotachophoresis 342(1985)167

Mono(2-ethylhexyl) phthalate
Determination of phthalic acid, mono-(2-ethylhexyl) phthalate and di(2-ethylhexyl) phthalate in human plasma and in blood products 337(1985)279

Determination of di- and mono(2-ethylhexyl) phthalate in plasma by GC 342(1985)399

Monoglycerides
HPLC technique for the separation of lipid classes 329(1985)273

Monomers
Chromatography of monomers. IV. GLC studies of C_1-C_6 *n*-alkyl and C_3-C_6 isoalkyl acrylates and their hydrogen halide and halogen addition derivatives 348(1985)141

(+)-Morphinan-3-ol
Determination of dextromethorphan and metabolites in human plasma and urine by HPLC with fluorescence detection 338(1985)99

Morphine
LC with pre-column dansyl derivatization and fluorimetric detection applied to the assay of morphine in biological samples 330(1985)323

Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Determination of drugs in untreated body fluids by micellar chromatography with fluorescence detection 342(1985)293

Improved method for the simultaneous determination of morphine, codeine and dihydrocodeine in blood by HPLC with electrochemical detection 347(1985)398

Motilin
Rapid, small-scale preparation of gastrointestinal hormones by HPLC on a C_{18} column 326(1985)121

Muldamine
Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

Multichannel detectors
Multichannel spectrophotometric detector for fused-silica capillary tube isotachophoresis 346(1985)167

Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285

Munitions
Reversed-phase gradient HPLC of nitramine munitions and characterization of munitions process samples by GC-MS 329(1985)285

Muramic acid
Modifications in the alditol acetate method for analysis of muramic acid and other neutral and amino sugars by capillary GC-MS with selected ion monitoring 347(1985)109

Muricholic acid
Determination of bile acids in rat bile by HPLC 338(1985)195

Mycophenolic acid
TLC of mycotoxins (Review) 334(1985)211

Mycotoxins
Rapid and sensitive determination of zearalenone in cereals by HPLC with fluorescence detection 328(1985)271

TLC of mycotoxins (Review) 334(1985)211

GC assay with pharmacokinetic applications for monitoring T-2 and HT-2 toxins in plasma 343(1985)67

Analysis of some trichothecene mycotoxins by LC 347(1985)429

Myelin basic protein
Separation and isolation of human myelin basic protein peptides by reversed-phase HPLC in volatile buffers 329(1985)295

Myoglobin
Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

N

Nabumetone
HPLC determination of a new anti-inflammatory agent, nabumetone, and its major metabolite in plasma using fluorimetric detection 336(1985)234

NADH
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Nadolol
Assay of nadolol in serum by reversed-phase HPLC with fluorometric detection 342(1985)206

NADPH-cytochrome P-450 reductase
Purification of oestrogen synthetase by HPLC. Two membrane-bound enzymes from the human plancenta 326(1985)137

Nafidimide
Sensitive reversed-phase HPLC assay for the antitumor agent benzisoquinolinedione (Nafidimide) employing direct on-column injection of plasma and urine, and its use for pharmacokinetic studies in a dog 345(1985)402

HPLC separation and quantitation of benzisoquinolinedione (Nafidimide) from biological fluids 345(1985)408

Naftidrofuryl
Determination of naftidrofuryl in human plasma by HPLC with fluorescence detection 338(1985)433

Naphtha
Naphtha analysis. The advantages of a specific olefin trap 349(1985)377

Naphthalenes
Evaluation of tetraalkylammonium tetrafluoroborate salts as high-temperature stationary phases for packed and open-tubular column GC 349(1985)249

β-Naphthalenesulphonic acid
Characterization of neutralized β-naphthalenesulfonic acid and formaldehyde condensates 347(1985)249

2-Naphthalenesulphonyl chloride
Derivatization of secondary amines with 2-naphthalenesulfonyl chloride for HPLC analysis of spectinomycin 333(1985)365

Naphthenes
Quantitative analysis of hydrocarbons in gasolines by capillary GLC. II. Isothermal and temperature-programmed analyses 346(1985)177

Naphthols
Effects of various strengths of bases on the electrophoretic behavior of weak acids in a non-aqueous solvent 330(1985)19

Comparative GC studies of bonded and physically coated PEG 20M and its derivative phases with reference to some high-boiling isomeric solutes 346(1985)359

Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

Naphtho-γ-pyrones
TLC of mycotoxins (Review) 334(1985)211

Naphthylamines
Comparative GC studies of bonded and physically coated PEG 20M and its derivative phases with reference to some high-boiling isomeric solutes 346(1985)359

Measurement of trace aromatic amines in seawater using HPLC with electrochemical detection 348(1985)265

Naphtols
HPLC determination of naphthols as 4-aminoantipyrine derivatives. Application to carbaryl 349(1985)63

Neomycin
Analysis of aminoglycoside antibiotics as benzoyl derivatives by HPLC and its application to the quantitation of neomycin in the perilymph 337(1985)187

Neuroendocrine polypeptides
Quantitative affinity HPLC of neuroendocrine polypeptides using porous and non-porous glass derivatives 327(1985)193

Neurophysins
Quantitative affinity HPLC of neuroendocrine polypeptides using porous and non-porous glass derivatives 327(1985)193

HPLC and studies of neurophysin-neurohypophysial hormone pathways 336(1985)63

Neurotensin
Weak anion-exchange HPLC of peptides (Review) 334(1985)49

Neurotransmitters
Determination of neurotransmitter amino acids by HPLC with fluorescence detection 337(1985)391

Nickel chelates
HPLC of copper, cobalt, nickel, chromium and vanadium chelates with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol 333(1985)133

Nicotine
Isolation and analysis of N-oxide metabolites of tertiary amines: quantitation of nicotine-1'-N-oxide formation in mice 341(1985)349

Rapid GC method for the determination of nicotine and cotinine in urine (in French) 343(1985)424

Simple HPLC method for rapid determination of nicotine and cotinine in urine 344(1985)391

HPLC with electrochemical detection for the determination of nicotine and N-methylnicotinium ion 347(1985)405

Nicotine-1'-N-oxide
Isolation and analysis of N-oxide metabolites of tertiary amines: quantitation of nicotine-1'-N-oxide formation in mice 341(1985)349

Nifedipine
Determination of nifedipine in human plasma by HPLC with electrochemical detection 341(1985)341

Study of nifedipine photodecomposition in plasma and whole blood using capillary GLC 342(1985)193

Nilvadipine
Determination of nilvadipine in human plasma by capillary column GC-negative-ion chemical-ionization MS 345(1985)51

Ninhydrin
HPLC analysis of guanidino compounds using ninhydrin reagent. II. Guanidino compounds in blood of patients on haemodialysis therapy 342(1985)269

Nitramine munitions
Reversed-phase gradient HPLC of nitramine munitions and characterization of munitions process samples by GC-MS 329(1985)285

Nitrate esters
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Nitrazepam
Simplified GC assay of underivatized nitrazepam in plasma 337(1985)131

Determination of nitrazepam and its main metabolites in urine by TLC and direct densitometry 339(1985)163

Nitrendipine
Quantification of nitrendipine in plasma by a capillary column GC-MS methods 337(1985)402

Nitrilotriacetic acid
Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147

Nitroanilines
α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131

***p*-Nitroanisole**
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Nitrobenzoic acid
Methylated cyclodextrins as mobile phase additives in LC 330(1985)253

Nitrocelluloses
Determination of molecular weight data in size-exclusion chromatography by use of braod-molecular-weight-distributed standards 349(1985)357

Nitrocinnamic acid
Methylated cyclodextrins as mobile phase additives in LC 330(1985)253

Nitrogen heterocycles
Class separation of polycyclic aromatic hydrocarbons, nitrogen heterocycles and hydroxyl aromatics by LC 331(1985)55

Nitroglycerin
Quantitative determination of nitroglycerin in human plasma by capillary GC with electron-capture detection 339(1985)117

Rapid, isothermal GLC determination of nitroglycerine in plasma using an electron-capture detector. II 349(1985)421

Nitromethane
Effects of test solutes on the characterization of stationary phases in GC using a Snyder triangle 329(1985)5

Nitronaphthalenes
α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131

Nitrophenols
Effects of various strengths of bases on the electrophoretic behavior of weak acids in a non-aqueous solvent 330(1985)19

Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

α- and β- cyclodextrins as selective agents for the separation of isomers by reversed-phase HPTLC and column LC 349(1985)131

Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

Nitrosamines
Analysis of volatile nitrosamines by microbore HPLC and thermal energy analyser detection 328(1985)362

Nitroso compounds
Improved staining procedure for nucleic acids in polyacrylamide gels after complexing with nitroso compounds 347(1985)384

Nitrosoureas
Drug level monitoring: cytostatics (Review) 340(1985)31

Nivalenol
Rapid and sensitive determination of zearalenone in cereals by HPLC with fluorescence detection 328(1985)271

Analysis of some trichothecene mycotoxins by LC 347(1985)429

Noble gases
Noble, diatomic and aliphatic gas analysis by aqueous HPLC 346(1985)93

Nomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 341(1985)333

GC-MS determination of nomifensine using a stable isotope-labeled analogue as an internal standard 344(1985)362

Non-amphoteric buffers
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Non-catastrophe theory
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Non-steroidal antirheumatic drugs
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77

Noracetylmethadol
Quantitation of *l*-α-acetylmethadol and its metabolites in human serum by capillary GLC and nitrogen detection 343(1985)339

Noradrenaline
Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Comparison of HPLC with electrochemical detection and GC-mass fragmentography for the assay of salsolinol, dopamine and dopamine metabolites in food and beverage samples 336(1985)199

3-Methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, and homovanillic acid in human cerebrospinal fluid. Storage and measurement by reversed-phase HPLC and coulometric detection using 3-methoxy-4-hydroxyphenyllactic acid as an internal standard 336(1985)259

Direct clean-up and analysis of urinary catecholamines 336(1985)374

Simple and fast analysis of adrenaline and noradrenaline in plasma on microbore HPLC columns using fluorimetric detection 337(1985)384

Determination of adult sheep plasma catecholamines using [^{3}H]norepinephrine as the internal standard 338(1985)192

Simultaneous assay of 3,4-dihydroxyphenylethylene glycol and norepinephrine in human plasma by HPLC with electrochemical detection 338(1985)401

Estimation of adrenal catecholamines by elevated-temperature LC with amperometric detection 342(1985)363

Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370

HPLC assay of free norepinephrine, epinephrine, dopamine, vanillylmandelic acid and homovanillic acid 343(1985)9

Determination of intraoperative plasma catecholamine concentrations using LC with electrochemical detection 343(1985)285

Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Measurement of urinary catecholamines and their catechol metabolites and precursor by LC with column-switching and on-line fluorimetric and electrochemical detection 344(1985)51

HPLC of plasma catecholamines using 1,2-diphenylethylenediamine as precolumn fluorescence derivatization reagent 344(1985)61

Plasma levels of norepinephrine and epinephrine during malignant hyperthermia in susceptible pigs 344(1985)71

Rapid and reliable estimation of urinary free catecholamines in patients with pheochromocytoma: comparison with plasma catecholamines and vanillylmandelic acid excretion 344(1985)313

Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 345(1985)140

19-Norandrostan-3-ol-17-ones
GC-MS of epimeric 19-norandrostan-3-ol-17-ones as the trimethylsilyl ether, methyloxime-trimethylsilyl ether and trimethylsilyl-enol trimethylsilyl ether derivatives 339(1985)11

Norbuprenorphine
^{63}Ni electron-capture GC assay for buprenorphine and metabolites in human urine and feces 337(1985)291

Analysis of buprenorphine and its N-dealkylated metabolite in plasma and urine by selected-ion monitoring 338(1985)89

Nordextropropoxyphene
HPLC method for the determination of dextropropoxyphene and nordextropropoxyphene in serum 345(1985)413

Norepinephrine
see Noradrenaline

Norfloxacin
Improved HPLC determination of pefloxacin and its metabolite norfloxacin in human plasma and tissue 339(1985)214

Nor-ketamine
Quantitative deconvolution of heavily fused chromatographic peaks of biological components using a multi-wavelength UV detector 332(1985)139

Normeperidine
Analysis of meperidine and normeperidine in serum and urine by HPLC 338(1985)141

Normetanephrine
Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

11-Nor-Δ^9-tetrahydrocannabinol-9-carboxylic acid
Antibody-mediated extraction of the main tetrahydrocannabinol metabolite, 11-nor-Δ^9-tetrahydrocannabinol-9-carboxylic acid, from human urine and its identification by GC-MS in the sub-nanogram range 342(1985)393

Norverapamil
Simultaneous quantitation of verapamil, norverapamil, and N-dealkylated metabolites in human plasma following oral administration 337(1985)160

HPLC measurement of verapamil in plasma using a diol column 337(1985)412

Noscapine
New sensitive method to determine noscapine in serum by reversed-phase LC 337(1985)429

Novolacs
Separation and identification of phenol-formaldehyde condensates by GC-MS. I. Acid-catalysed condensation products 329(1985)290

Nuclear magnetic resonance spectroscopy
Continuous-flow nuclear magnetic resonance 346(1985)17

Nuclease
Purification of a uridine-specific acid nuclease from chicken liver by fast protein LC 326(1985)157

Nucleic acids
Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33

HPLC of tRNAs on novel stationary phases 326(1985)79

Affinity partitioning of biopolymers and membranes in Ficoll-dextran aqueous two-phase systems 331(1985)11

Recent progress in gel packing materials and detectors for modern LC in Japan 332(1985)227

HPLC of DNA (Review) 336(1985)3

Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 341(1985)182

Preparative isolation of plasmid DNA with sedimentation field flow fractionation 342(1985)67

Improved staining procedure for nucleic acids in polyacrylamide gels after complexing with nitroso compounds 347(1985)384

Nucleobases
see also Purines and pyrimidines

HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33

Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33

Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255

Recent developments in isotachophoresis (Review) 334(1985)157

HPLC method for the analysis of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in biological fluids 339(1985)380

Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 341(1985)182

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Use of a fundamental elution protocol for the development of reversed-phase HPLC enabling rapid simultaneous determination of purines, pyrimidines and allied compounds commonly found in human biological fluids 344(1985)101

Evaluation of combined electrochemical and UV detection in the HPLC analysis of plasma for low-molecular-weight constituents 345(1985)125

Nucleoproteins
Analysis of nucleoproteins by direct injection of dissolved nuclei or chromosomes into a HPLC system 326(1985)199

Nucleosides
Reversed-phase HPLC of nucleotides and oligonucleotides 326(1985)281

HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33

Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33

Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255

Recent developments in isotachophoresis (Review) 334(1985)157

Reversed-phase HPLC determination of tricyclic nucleoside and tricyclic nucleoside 5'-phosphate in biological specimens 337(1985)55

Mobile phase and temperature studies in the reversed-phase LC of inosine and guanosine compounds 339(1985)75

HPLC method for the analysis of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in biological fluids 339(1985)380

Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 341(1985)182

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Use of a fundamental elution protocol for the development of reversed-phase HPLC enabling rapid simultaneous determination of purines, pyrimidines and allied compounds commonly found in human biological fluids 344(1985)101

Evaluation of combined electrochemical and UV detection in the HPLC analysis of plasma for low-molecular-weight constituents 345(1985)125

Nucleotides
Reversed-phase HPLC separation of diastereomeric phosphorothioate analogues of oligodeoxyribonucleotides and other backbone-modified congeners of DNA 326(1985)263

Reversed-phase HPLC of nucleotides and oligonucleotides 326(1985)281

Use of a C_4 column for reversed-phase HPLC purification of synthetic oligonucleotides 326(1985)293

HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33

Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. I. Cation binding in the mobile phase 329(1985)321

Use of fast protein LC for the purification of synthetic oligonucleotides 329(1985)406

Reversed-phase LC of biotin-labelled nucleotides. A new class of markers in molecular biology 330(1985)429

Behaviour of single-stranded oligodeoxyribonucleotides on a DEAE-5PW anion-exchange column 332(1985)243

Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255

Recent developments in isotachophoresis (Review) 334(1985)157

Purification of synthetic oligodeoxyribonucleotides by ion-exchange HPLC 336(1985)189

Simultaneous determination of cytosine arabinoside, its nucleotides and metabolites by ion-pair HPLC 337(1985)63

Assay of human erythrocyte pyrimidine and deoxypyrimidine 5'-nucleotidase by isocratic reversed-phase HPLC 339(1985)293

High-performance liquid chromatographic detection of XMP as a basis for an improved IMP dehydrogenase assay 343(1985)160

Analyser of adenine nucleotides 344(1985)345

Evaluation of combined electrochemical and UV detection in the HPLC analysis of plasma for low-molecular-weight constituents 345(1985)125

Rapid determination of adenine nucleotides in brain tissue by ion-paired reversed-phase column LC under isocratic conditions 345(1985)267

Multichannel spectrophotometric detector for fused-silica capillary tube isotachophoresis 346(1985)167

Reversed-phase LC of protected oligonucleotide diesters 348(1985)286

Simple separation of adenosine-5'-phosphosulphate, adenosine-5'-phosphoramidate and cyclic adenosine-5'-monophosphate nucleotides involved in adenosine-5'-phosphosulphate degradation 348(1985)296

Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. II. Quantitative determination of hydrophobicity 350(1985)105

Reversed-phase HPLC of protected oligodeoxynucleotides 350(1985)313

O

Ochratoxins
TLC of mycotoxins (Review) 334(1985)211

Octadecyldimethyl-N,N-diethylaminosilane
On-column preparation of chemically bonded stationary phase with maximum surface coverage and high reproducibility, and its application to packed microcapillary columns 332(1985)57

Octadecylsilane
Properties of six octadecylsilane stationary phases for the separation of dansylamides 330(1985)243

***n*-Octane**
Reversed-flow GC technique applied to measurement of mass transfer coefficients of *n*-hydrocarbons on Porapak P 348(1985)339

Octanoic acid
Chromatographic determination of N-acetyl-DL-tryptophan and octanoic acid in human albumin solutions 333(1985)381

Octopamine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Octyl-Sepharose CL-4B
Three independent methods for quantitative determination of octyl covalently coupled to Sepharose CL-4B 346(1985)255

Odorous compounds
Stripping analysis and chromatographic sniffing for the source identification of odorous compounds in drinking water 328(1985)219

Oestradiol 17-sulphate 4-hydroxylase
Clinical analysis of steroids. XXXI. Assay of oestradiol 17-sulphate 4-hydroxylase activity

by HPLC with electrochemical detection 337(1985)114

Oestriol
Total oestriol in maternal serum or plasma as measured by LC 337(1985)379

Determination of oestriol in pregnancy urine by normal-phase HPLC with electrochemical detection 341(1985)271

Oestrogen receptors
Isolation of unliganded steroid receptor proteins by high-performance size-exclusion chromatography. An investigation of steroid-dependent structural alterations 327(1985)221

High-performance hydrophobic-interaction chromatography of steroid hormone receptors 327(1985)237

High-performance chromatofocusing of steroid receptor proteins in the presence and absence of steroid. Investigation of steroid-dependent alterations in surface charge heterogeneity 327(1985)247

Rapid, high-resolution procedure for assessment of estrogen receptor heterogeneity in clinical samples 327(1985)369

Oestrogens
Rapid extraction and purification of diethylstilboestrol in bovine urine hydrolysates using reversed-phase C_{18} columns before determination by radioimmunoassay 328(1985)413

Clinical analysis of steroids. XXXI. Assay of oestradiol 17-sulphate 4-hydroxylase activity by HPLC with electrochemical detection 337(1985)114

Efficient chromatographic fractionation of steroids in human serum through regulation of liquid-liquid distribution ratios 339(1985)253

Oestrogen synthetase
Purification of oestrogen synthetase by HPLC. Two membrane-bound enzymes from the human plancenta 326(1985)137

Oil residues
Group separation of oil residues by supercritical fluid chromatography 349(1985)439

Oils
Adsorption chromatographic separations on buffered silica gel (Review) 334(1985)35

Determination of trace chlorinated benzenes in fuel oil by on-line multidimensional chromatography using packed-capillary LC and capillary GC 349(1985)55

Olefins
Naphtha analysis. The advantages of a specific olefin trap 349(1985)377

Olefin trap
Naphtha analysis. The advantages of a specific olefin trap 349(1985)377

Oligodeoxynucleotides
Reversed-phase HPLC of protected oligodeoxynucleotides 350(1985)313

Oligodeoxyribonucleotides
Reversed-phase HPLC separation of diastereomeric phosphorothioate analogues of oligodeoxyribonucleotides and other backbone-modified congeners of DNA 326(1985)263

Use of a C_4 column for reversed-phase HPLC purification of synthetic oligonucleotides 326(1985)293

Behaviour of single-stranded oligodeoxyribonucleotides on a DEAE-5PW anion-exchange column 332(1985)243

Purification of synthetic oligodeoxyribonucleotides by ion-exchange HPLC 336(1985)189

Oligomers
Supercritical fluid chromatography-IR spectroscopy of oligomers: use of buffer-memory technique 332(1985)47

Separation of oligomers with UV-absorbing side groups by supercritical fluid chromatography using eluent gradients 346(1985)69

Oligonucleotides
Reversed-phase HPLC of nucleotides and oligonucleotides 326(1985)281

Use of a C_4 column for reversed-phase HPLC purification of synthetic oligonucleotides 326(1985)293

Use of fast protein LC for the purification of synthetic oligonucleotides 329(1985)406

Reversed-phase LC of protected oligonucleotide diesters 348(1985)286

Oligo-oxaalkanoyl tetraamides
Oligo-oxaalkanoyl tetraamides derived from L-phenylalanine as stationary phases in capillary GC resolution of D,L-amino acids 349(1985)9

Oligosaccharides
Comparison of relationships between the chemical structures and mobilities of hyaluronate oligosaccharides in TLC and HPLC 328(1985)73

Characterization of lacto-N-hexaose and two fucosylated derivatives from human milk by HPLC and proton NMR spectroscopy 328(1985)259

Some applications of reversed-phase HPLC to oligosaccharide separations 336(1985)161

HPLC analysis of changes of asparagine-linked oligosaccharides in regenerating rat liver 345(1985)134

On-line detection
Monitoring radioactive compounds in HPLC eluates: fraction collection versus on-line detection 342(1985)25

Open stripping system
Optimization of the open stripping system for the analysis of trace organics in water 348(1985)67

Optimization
Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Application to method development and the choice of column configuration 327(1985)93

Optimized flow programming for temperature-programmed GC 328(1985)49

Solubility parameter and window diagram optimization methods for the reversed-phase chromatographic optimization of complex hydroxyl aromatic mixtures 328(1985)81

Computer-aided mobile phase optimization and chromatogram simulation in HPLC (Review) 338(1985)1

Ion-exchange HPLC of mouse epidermal growth factor and its congeners: mobile phase optimization with ion-pairing additives 345(1985)168

Optimization of the open stripping system for the analysis of trace organics in water 348(1985)67

A problem of optimization in capillary GC 348(1985)416

Optimization of helium ionization detector 349(1985)405

Effect of column temperature on HPLC behaviour of inorganic polyphosphates. II. Gradient ion-exchange chromatography 350(1985)119

HPLC determination of contaminants in salicylic acids after computer-assisted optimization of the separation parameters (in German) 350(1985)471

Organic acids
see also Carboxylic acids

Recent developments in isotachophoresis (Review) 334(1985)157

HPLC of microbial acid metabolites 336(1985)125

3-Hydroxyhexanoic acid: an abnormal metabolite in urine and serum of diabetic ketoacidotic patients 337(1985)1

Absolute configuration of 3-hydroxyadipic acid in human urine 337(1985)9

N-Methylation and N,N-dimethylation of amino acids. An artifact production in the analysis of organic acids using diazomethane as derivatizing agent 338(1985)33

Identification of metabolites diagnostic for organic acidurias by simultaneous dual-column capillary GC 341(1985)1

Multichannel spectrophotometric detector for fused-silica capillary tube isotachophoresis 346(1985)167

Organic additives
Analysis technique for determining the levels of organic additives in an ethylene-propylene diene monomer rubber for use in underwater acoustic applications 350(1985)187

Organic bases
Recent developments in isotachophoresis (Review) 334(1985)157

Organic compounds
Retention indices as a tool for the prediction of chromatographic conditions for achieving the separation of mixtures of organic compounds 330(1985)71

Semi-preparative HPLC analysis of complex organic mixtures 332(1985)169

New methods of interpretation of chromatographic-MS data 349(1985)385

Trace analysis of organic compounds in volatile inorganic halides by the dynamic headspace technique and GC-MS 350(1985)324

Organochlorine compounds
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Organohalogen compounds
Trace analysis of organic compounds in volatile inorganic halides by the dynamic headspace technique and GC-MS 350(1985)324

Organometallic compounds
LC retention behavior of organometallic compounds and ligands with amine-, octadecylsilica- and β-cyclodextrin-bonded phase columns 347(1985)51

Organophosphorus compounds
GC retention indices of chemical warfare agents and simulants 331(1985)47

Determination of methyl paraoxon in dog plasma by reversed-phase HPLC 338(1985)389

Capillary column GC-MS of tabun 347(1985)257

Organosulphur compounds
Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Evaluation of the suitability of selected porous polymers for preconcentration of organosulphur compounds from water 346(1985)61

Orosomucoid
Two-step purification of human α_1-acid glycoprotein 341(1985)457

Oroxylin-A
Simultaneous determination of baicalein, wogonin, oroxylin-A and their glucuronides in Scutellariae Radix by ion-pair HPLC 328(1985)289

Ovalbumin
Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Oxcarbazepine
GC assay for oxcarbazepine and its main metabolites in plasma 345(1985)67

Oxido-compounds
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

Oxindazac
Isolation and stereospecific determination of the enantiomers of oxindazac by direct LC resolution on triacetylcellulose 346(1985)321

Oxiranes
Determination of the enantiomeric composition of chiral epoxides using chiral derivatization and LC 331(1985)349

4-Oxo-enoxacin
HPLC and preliminary pharmacokinetics of enoxacin and its 4-oxo metabolite in human plasma, urine and saliva 343(1985)449

Oxosteroids
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 341(1985)426

Oxpentifylline
Determination of oxpentifylline and four metabolites in plasma and urine by automated capillary GC using nitrogen-selective detection 344(1985)187

Determination of 1-(3'-carboxypropyl)-3,7-dimethylxanthine and 1-(4'-carboxybutyl)-3,7-dimethylxanthine, two major metabolites of oxpentifylline, in urine by HPLC 344(1985)397

Oxprenolol
Determination of oxprenolol in human plasma by HPLC, in comparison with GC and GC-MS 345(1985)365

Oxygen
Quantification of branched-chain α-keto acids as quinoxalinols: importance of excluding oxygen during derivatization 337(1985)98

Oxyhaemoglobin
Analytical isoelectric focusing of native and modified haemoglobins after treatment with 4-hydroxymercuribenzoate 346(1985)401

Oxytetracycline
Determination of oxytetracycline, sulphamethazine and sulphamethoxypyridazine in feed premixes 330(1985)425

Reversed-phase HPLC method for the assay of oxytetracycline 347(1985)173

Oxytocin
HPLC and studies of neurophysin-neurohypophysial hormone pathways 336(1985)63

Ozone chemiluminescence
GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

P

Paclobutrazol
Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181

Panose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Panthenol
Determination of optical purity by enantioselective capillary GC: panthenol and related compounds 328(1985)357

Pantothenic acid
Determination of optical purity by enantioselective capillary GC: panthenol and related compounds 328(1985)357

Papaverine
Determination of papaverine by capillary isotachophoresis 350(1985)336

Parabens
Studies on the analysis of food additives by HPLC. V. Simultaneous determination of preservatives and saccharin in foods by ion-pair chromatography 346(1985)333

Paracetamol
Determination of paracetamol, salicylic acid and acetyl salicylic acid in serum by HPLC 329(1985)196

Analysis of antipyretics by semimicro LC 332(1985)269

Micro HPLC assay of acetaminophen and its major metabolites in plasma and urine 339(1985)198

Simultaneous HPLC stability-indicating analysis of acetaminophen and codeine phosphate in tablets and capsules 348(1985)253

Paraffins
Quantitative analysis of hydrocarbons in gasolines by capillary GLC. II. Isothermal and temperature-programmed analyses 346(1985)177

Parathyroid hormone
Comparison of reversed-phase and cation-exchange HPLC for separating closely related peptides: separation of Asp^{76}-human parathyroid hormone (1-84) from Asn^{76}-human parathyroid hormone (1-84) 327(1985)327

Paraxanthine
Improved micro-method for the HPLC determination of caffeine and paraxanthine in biological fluids 336(1985)293

HPLC determination of dimethylxanthine metabolites of caffeine in human plasma 338(1985)369

Particle diameter
Contribution to the study of micropacked columns in GC 348(1985)327

Particle size
Theoretical investigation of the optimum particle size for the resolution of proteins by size-exclusion chromatography 326(1985)3

Partition coefficients
HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. II. Retention behavior, quantitative structure-retention relationships and octanol-water partition coefficients 330(1985)181

Ionization constants and lipophilicity of 1-arylpiperazines investigated by reversed-phase HPLC 333(1985)191

Influence of the mobile phase composition on the reversed-phase TLC behaviour of a series of prostaglandins. Comparison of the extrapolated R_M values 347(1985)209

TLC and HPLC in the evaluation of the lipophilicity of aryloxoalkanoic and arylhydroxyalkanoic acids 347(1985)335

Paspalitrem
TLC of mycotoxins (Review) 334(1985)211

Pasteurella
Differentiation between major species of the *Actinobacillus-Haemophilis-Pasteurella* group by GC of trifluoroacetic acid anhydride derivatives from whole-cell methanolysates 342(1985)13

Pathotoxins
Rapid purification of host-specific pathotoxins from *Alternaria alternata* f. sp. *lycopersici* by solid-phase adsorption on octadecylsilane 350(1985)255

Pattern recognition
Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. I. Introduction and characterization of the cuticular hydrocarbon patterns of *Solenopsis invicta* and *S. richteri* 349(1985)31

Use of computerized pattern recognition in the study of the cuticular hydrocarbons of imported fire ants. II. Comparison of the cuticular hydrocarbon patterns between different colonies of *Solenopsis richteri* 349(1985)39

Patulin
TLC of mycotoxins (Review) 334(1985)211

PCBs
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Vapor pressure estimates of individual polychlorinated biphenyls and commercial fluids using GC retention data 330(1985)203

Recent advances in the analysis of polychlorinated biphenyls in environmental and biological media (Review) 334(1985)277

Novel approaches to the determination of polychlorinated biphenyls 349(1985)425

Peak capacity
HPLC theory for the practitioner (Review) 334(1985)197

Peak integration
Integration of the deformed peak from the helium ionization detector 349(1985)415

Peak overlap
Deconvolution of overlapping chromatograms. II. Analytical solution to the moments of the GEX function 333(1985)152

Peak purity
Application of derivative spectroscopy to the determination of chromatographic peak purity 347(1985)219

Peak shape
HPLC of amino acids, peptides and proteins. LXVI Investigations on the effects of chromatographic dwell in the reversed-phase HPLC separation of proteins 327(1985)47

Integration of the deformed peak from the helium ionization detector 349(1985)415

Peak splitting
Computer simulation of abnormal high-performance liquid chromatograms caused by solvents 329(1985)13

Proposed model for peak splitting in reversed-phase ion-pair HPLC with computer prediction of eluted peak profiles 336(1985)15

Peak volume
HPLC theory for the practitioner (Review) 334(1985)197

Pefloxacin
Improved HPLC determination of pefloxacin and its metabolite norfloxacin in human plasma and tissue 339(1985)214

Pellethane
Chromatographic analysis of thermoplastic, medical-grade, polyether-based polyurethane elastomers 349(1985)347

Penicillamine
Analysis of D-penicillamine by GC utilizing nitrogen-phosphorus detection 337(1985)37

Application of chromatographic chiral stationary phases to pharmaceutical analysis. Enantiomeric purity of D-penicillamine 350(1985)179

Penicillic acid
TLC of mycotoxins (Review) 334(1985)211

Penicillins
Antibiotic monitoring in body fluids (Review) 340(1985)361

HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 341(1985)115

Studies on residual antibacterials in foods. IV. Simultaneous determination of penicillin G, penicillin V and ampicillin in milk by HPLC 348(1985)379

Penitrem
TLC of mycotoxins (Review) 334(1985)211

1,4,5,6,8-Pentaazaacenaphthylene-3-amino-1,5-dihydro-5-methyl-1-β-D-ribofuranosyl 5'-monophosphate
Reversed-phase HPLC determination of tricyclic nucleoside and tricyclic nucleoside 5'-phosphate in biological specimens 337(1985)55

Pentacaine
GC determination of pentacaine in rat serum 339(1985)410

Pentachlorophenol
Rapid and convenient separation of pentachlorophenol from human fat using silica Sep-Pak cartridges 346(1985)435

Pentachloropyridine
Analysis of some chlorinated pyridine isomers in mixtures by reversed-phase HPLC 348(1985)309

Pentaerythritol tetranitrate
Examination of the in vitro degradation of [^{14}C]pentaerythritol tetranitrate in rat and human blood with an improved thin-layer radiochromatographic procedure 343(1985)129

Pentafluorobenzoyl esters
GLC analyses. XLIV. Benzoyl and pentafluorobenzoyl esters of lower saturated branched-chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 329(1985)359

Pentafluorobenzyl *p*-toluenesulphonate
Pentafluorobenzyl *p*-toluenesulphonate as a new derivatizing reagent for GC determination of anions 346(1985)215

O-Pentafluorophenylsulphonylphenols
Relative precision in the determination of some structurally-similar O-pentafluorophenylsulfonylphenols by ultratrace GC 331(1985)396

Pentamidine
HPLC method for quantitation of pentamidine in blood serum 345(1985)91

Pentobarbital
Determination of thiopental and pentobarbital in plasma using HPLC 343(1985)437

Pentopril
Determination of pentopril, an angiotensin converting enzyme inhibitor, and its active metabolite in urine 345(1985)396

Peptidase
Highly sensitive assay for PZ-peptidase activity by HPLC 348(1985)205

Peptides
Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33

Use of reversed-phase and ion-exchange batch extraction in the purification of bovine pituitary peptides 326(1985)329

Novel approach to rapid and sensitive localization of protein disulfide bridges by HPLC and electrochemical detection 326(1985)339

The bacterial phosphoenolpyruvate-dependent phosphotransferase system. Isolation of active site peptides by reversed-phase HPLC and determination of their primary structure 326(1985)363

Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399

Automated tandem HPLC system for separation of extremely complex peptide mixtures 326(1985)407

Comparison of reversed-phase columns for the separation of tryptic peptides by gradient elution. Correlation of experimental results and model prediction 326(1985)419

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27

Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Application to method development and the choice of column configuration 327(1985)93

Separation of peptides by strong cation-exchange HPLC 327(1985)147

Improved detection limits in the analysis of tyrosine-containing polypeptide hormones by using electrochemical detection 327(1985)165

HPLC analysis of *in vitro* central neuropeptide processing 327(1985)279

Comparison of reversed-phase and cation-exchange HPLC for separating closely related peptides: separation of Asp^{76}-human parathyroid hormone (1-84) from Asn^{76}-human parathyroid hormone (1-84) 327(1985)327

Adsorption of SH-containing amino acids, peptides and proteins on the 2-nitrophenylhydrazonopropanedinitrile derivative of Spheron Ara 328(1985)402

Separation and isolation of human myelin basic protein peptides by reversed-phase HPLC in volatile buffers 329(1985)295

Effects of solutions used for storage of size-exclusion columns on subsequent chromatography of peptides and proteins 331(1985)253

Sequence determination of cyanogen bromide-peptides by combined use of HPLC and fast atom bombardment MS 331(1985)425

HPLC analysis of peptides on an Asahipak GS-320 column packed with hydrophilic polymer gel 332(1985)287

Weak anion-exchange HPLC of peptides (Review) 334(1985)49

Recent developments in isotachophoresis (Review) 334(1985)157

Measurement and chromatographic characterization of vasoactive intestinal peptide from guinea-pig enteric nerves 336(1985)41

Separation of tryptic phosphopeptides of ribosomal origin by reversed-phase HPLC 336(1985)51

Separation of biologically important angiotensin peptides by HPLC on a weak cationic exchange bonded phase 336(1985)392

Development of a system to simplify identification of peptides and their catabolites in a HPLC effluent 337(1985)110

New application of HPLC for analysis of urinary C-peptide 338(1985)71

Separation of somatomedins and somatomedin inhibitors by size exclusion HPLC 339(1985)388

HPLC method for monitoring leupeptin in mouse serum and muscle by pre-column fluorescence derivatization with benzoin 345(1985)259

Isotachophoretic control of peptide synthesis and purification. A novel approach using UV detection at 206 nm 346(1985)271

Reversed-phase HPLC of natural and synthetic sauvagines 346(1985)390

Spectral properties of trifluoroacetic acid-acetonitrile gradient systems for separation of picomole quantities of peptides by reversed-phase HPLC 347(1985)83

Purification of synthetic lipid associating peptides and the monitoring of the deformylation of N^{in}-formyltryptophan by reversed-phase HPLC 347(1985)237

Purification and characterization of CAMP-factor from *Streptococcus agalactiae* by hydrophobic interaction chromatography and chromatofocusing 348(1985)363

Reversed-phase HPLC characterization of a novel proline-rich tryptophyllin 349(1985)117

Separation of some peptides and related isopeptides by HPLC: structure-retention time relationships 350(1985)279

Permanent gases
Determination of permanent gases and light hydrocarbons by simultaneous operation on packed and capillary columns with thermal conductivity detection and flame-ionization detection on a single computer integrator 349(1985)17

Integration of the deformed peak from the helium ionization detector 349(1985)415

Permeability
Contribution to the study of micropacked columns in GC 348(1985)327

Peroxidase
Electrophoretic detection of peroxidase isoenzymes in polyacrylamide gels using hydrogen peroxide generated by rose bengal-glutathione in the presence of light 348(1985)313

Perphenazine
Quantitative determination of perphenazine and its dealkylated metabolite using HPLC 341(1985)244

Pesticides
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181

Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Simplified method for the clean-up and reversed-phase HPLC determination of benomyl in mangoes 330(1985)403

A simple TLC method for studies of radiolabelled pesticides on soil surfaces using soil-silica layers 331(1985)200

Determination of methyl paraoxon in dog plasma by reversed-phase HPLC 338(1985)389

Determination of phenylurea herbicides by HPLC with electrochemical detection 346(1985)369

GLC method for the determination of dodine in fruit 348(1985)437

Microcomputer programming in BASIC for the evaluation of capillary GC in the analysis of pesticide residues. I. MATRIXCOMP: a program that facilitates the recognition of interfering peaks from the biological matrix 349(1985)49

HPLC determination of naphthols as 4-aminoantipyrine derivatives. Application to carbaryl 349(1985)63

Coupling a high-performance liquid chromatograph with a liquid scientillation detector: example in the field of pyrethroids 349(1985)105

High-performance liquid chromatograph coupled with two detectors: a UV spectrometer and a polarimeter. Example in the field of pyrethroids: identification of enantiomers 349(1985)109

Petroporphyrins
Computerised capillary GC-MS studies of the petroporphyrins of the Gilsonite bitumen (Eocene, U.S.A.) 350(1985)37

pH effects
Separation of peptides by strong cation-exchange HPLC 327(1985)147

pH gradients
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Phenacetin
HPLC of antipyretics on chemically modified porous glass 332(1985)265

Analysis of antipyretics by semimicro LC 332(1985)269

Phenazopyridine
HPLC determination of phenazopyridine hydrochloride, tetracycline hydrochloride and sulphamethizole in combination 333(1985)249

Phenobarbital
Concurrent determination of valproic acid with other antiepileptic drugs by HPLC 338(1985)131

Simultaneous determination of anticonvulsant drugs and metabolites in plasma by HPLC 342(1985)469

Phenol
Application of thin-layer stick chromatographic identification test methods to drugs contained in preparations in the Pharmacopoeia of Japan 333(1985)115

Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

Phenolalkylamines
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Phenol effect
Phenol effect in PC of metal complexes 331(1985)273

Phenol-formaldehyde condensates
Separation and identification of phenol-formaldehyde condensates by GC-MS. I. Acid-catalysed condensation products 329(1985)290

Separation and identification of phenol-formaldehyde condensates by GC-MS. II. Base-catalysed condensation products 331(1985)91

Phenolic acids
HPLC of microbial acid metabolites 336(1985)125

GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

Rapid method for the separation and quantification of simple phenolic acids in plant material using HPLC 346(1985)440

HPTLC of phenylethylamines and phenolic acids on silanized silica and on ammonium tungstophosphate 347(1985)303

LC with voltammetric detection for quantitation of phenolic acids 348(1985)397

Phenolic compounds
High-performance size-exclusion chromatography of lignin and its derivatives 328(1985)299

Derivatization of butylated hydroxytoluene for GC with electron-capture detection and HPLC with UV detection 333(1985)171

HPLC of caffeoylquinic acid derivatives of *Cynara scolymus* L. leaves 348(1985)447

Phenols
GC and MS studies of the constituents of the rhizome of calamus. I. The volatile constituents of the essential oil 328(1985)179

Pentafluorobenzyl *p*-toluenesulphonate as a new derivatizing reagent for GC determination of anions 346(1985)215

Determination of phenols in water and wastewater by post-column reaction detection HPLC 350(1985)407

Phenprocoumon
Identification of phenprocoumon metabolites in human urine by HPLC and GC-MS 338(1985)325

Phentermine
GLC determination of phentermine in human plasma following oral administration to healthy subjects 337(1985)136

Phenylacetic acids
Correlation of the chromatographic retention of some phenylacetic and phenylpropionic acid derivatives with molecular structure 333(1985)309

Assay of urinary phenylacetic acid by HPLC 341(1985)172

Phenylalanine
Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

Phenylalanine ammonia-lyase
HPLC of proteins. Rapid chromatofocusing of plant enzymes 348(1985)167

Phenylalanine ethyl ester
Molecular imprinting of amino acid derivatives in macroporous polymers. Demonstration of substrate- and enantioselectivity by chromatographic resolution of racemic mixtures of amino acid derivatives 347(1985)1

Phenylalkanes
Kováts retention indices of high-molecular-weight monomethyl-, cyclopentyl-, cyclohexyl- and phenylalkanes 331(1985)219

5-(Phenylazo)-6-hydroxynaphthalene-2-sulphonic acid
Determination of the lower sulfonated subsidiary colors in FD&C Yellow No. 6 by reversed-phase HPLC 347(1985)163

***o*-Phenylenediamine**
LC with crown ether-containing mobile phases. V. Effect of hydrophobicity and cavity size of the crown ether on retention of amino compounds in reversed-phase HPLC 330(1985)43

Phenylephrine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Phenylethanolamine N-methyltransferase
Determination of phenylethanolamine N-methyltransferase by HPLC with fluorescence detection 348(1985)407

2-Phenylethyl alcohols
The aroma of grapes. I. Extraction and determination of free and glycosidically bound fractions of some grape aroma components 331(1985)83

Phenylethylamines
HPTLC of phenylethylamines and phenolic acids on silanized silica and on ammonium tungstophosphate 347(1985)303

Phenylglycine
Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

Monitoring of the production of D-phenylglycine from DL-phenylhydantoin by HPLC 329(1985)428

Phenylglyoxylic acid
Determination of mandelic and phenylglyoxylic acids in rat urine by HPLC and by isotachophoresis 343(1985)175

Phenylhydantoin
Monitoring of the production of D-phenylglycine from DL-phenylhydantoin by HPLC 329(1985)428

Phenylpropionic acids
Correlation of the chromatographic retention of some phenylacetic and phenylpropionic acid derivatives with molecular structure 333(1985)309

3-Phenylpropylsalicylate
HPLC determination of certain salicylates and their major metabolites in plasma following topical administration of a liniment to healthy subjects 342(1985)315

Phenylpyruvate
HPLC assays of phenylpyruvate and phenylpyruvate oxidase 337(1985)397

Phenylpyruvate oxidase
HPLC assays of phenylpyruvate and phenylpyruvate oxidase 337(1985)397

2-Phenylselenopropionic acid
Highly sensitive diastereoisomeric analysis of secondary alcohols and short-chain 1,2-diacylglycerols by capillary GC-MS 331(1985)295

Phenylureas
Determination of phenylurea herbicides by HPLC with electrochemical detection 346(1985)369

Phenytoin
Concurrent determination of valproic acid with other antiepileptic drugs by HPLC 338(1985)131

Analysis of the toxicologically relevant metabolites of phenytoin in biological samples by HPLC 338(1985)242

GC determination of *m*- and *p*-hydroxyphenytoin in the urine of epileptic patients 339(1985)331

Determination of antiepileptic drugs (Review) 340(1985)199

Urinary dihydrodiol metabolites of phenytoin: HPLC assay of diastereomeric composition 342(1985)323

Simultaneous determination of anticonvulsant drugs and metabolites in plasma by HPLC 342(1985)469

Pheromones
Determination of trimer in *Z*-11-hexadecenal pheromone by HPLC with a rotating disc flame ionization detector 329(1985)142

Retention indices as a tool for the prediction of chromatographic conditions for achieving the separation of mixtures of organic compounds 330(1985)71

New liquid crystalline stationary phases for GC of positional and geometrical isomers having similar volatilities 331(1985)69

Pholedrine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

Phosgene
Direct derivatization of alprenolol and its 4-hydroxy metabolite in urine with phosgene and methanol prior to analysis by capillary column GC 349(1985)447

Enantiomer separation of metoprolol and its analogues and metabolites by capillary column GC after derivatization with phosgene 350(1985)328

Phosphate
Determination of O-phosphothreonine, O-phosphoserine, O-phosphotyrosine and phosphate by HPLC 338(1985)315

Determination of ethyl phosphite, phosphite and phosphate in plant tissues by anion-exchange HPLC and GC 347(1985)267

Phosphate esters
Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

Phosphatidic acids
Formation and separation of fluorescent derivatives of diacylglycerols 331(1985)177

Phosphatidylcholine
Reversed-phase HPLC separation and mass detection of individual phospholipid classes 331(1985)313

Phosphatidylethanolamine
Reversed-phase HPLC separation and mass detection of individual phospholipid classes 331(1985)313

Estimation of amniotic fluid phospholipids by HPLC 336(1985)139

Fatty acid analysis in phosphatidylethanolamine subclasses of human erythrocyte membranes by HPLC 342(1985)359

Phosphatidylglycerol
Estimation of amniotic fluid phospholipids by HPLC 336(1985)139

Isolation and quantitative analysis of phosphatidylgylcerol and glycolipid molecular species using reversed-phase HPLC with flame ionization detection 346(1985)291

Phosphatidylinositol
Estimation of amniotic fluid phospholipids by HPLC 336(1985)139

Phosphatidylserine
Estimation of amniotic fluid phospholipids by HPLC 336(1985)139

Phosphine
Quantitative GC analysis of low-molecular-weight alkylphosphines in the presence of phosphine and hydrocarbons 330(1985)400

Phosphinolipids
Silicic acid CC of phosphinolipids. I. Phosphinate analogues of lecithin and cephalin, phosphonic acid and the phosphonate analogue of bis(diacylglycero)phosphate: their separation from other phospholipids and related phosphonolipids 329(1985)160

Phosphite
Determination of ethyl phosphite, phosphite and phosphate in plant tissues by anion-exchange HPLC and GC 347(1985)267

Phosphoamino acids
Determination of O-phosphothreonine, O-phosphoserine, O-phosphotyrosine and phosphate by HPLC 338(1985)315

Phosphoenolpyruvate-dependent phosphotransferase
The bacterial phosphoenolpyruvate-dependent phosphotransferase system. Isolation of active site peptides by reversed-phase HPLC and determination of their primary structure 326(1985)363

Phosphoglycerides
Application of silica gel CC in the assessment of non-esterified fatty acids and phosphoglycerides in myocardial tissue 344(1985)304

Phosphohistidines
The bacterial phosphoenolpyruvate-dependent phosphotransferase system. Isolation of active site peptides by reversed-phase HPLC and determination of their primary structure 326(1985)363

Phospholipids
Improved reproducibility and quantitative analysis of phospholipids by flame ionization detection 331(1985)303

Reversed-phase HPLC separation and mass detection of individual phospholipid classes 331(1985)313

Estimation of amniotic fluid phospholipids by HPLC 336(1985)139

Reversed-phase HPLC separation of molecular species of alkyl ether, vinyl ether, and monoacyl lysophospholipids 338(1985)61

Unusual fatty acids from amniotic fluid phospholipids 341(1985)146

Fatty acid analysis in phosphatidylethanolamine subclasses of human erythrocyte membranes by HPLC 342(1985)359

Rapid isocratic procedure for the separation of platelet-activating factor from phospholipids in human saliva by HPLC 342(1985)380

GC-MS analysis of biologically active phospholipids having and *sn*-2-acetyl group 343(1985)138

One-step separation of free fatty acids and phospholipids in brain tissue extracts by HPLC 344(1985)297

Improved separation of phospholipids by HPLC 344(1985)309

Phosphopeptides
Separation of tryptic phosphopeptides of ribosomal origin by reversed-phase HPLC 336(1985)51

Phosphorothioate analogues of oligodeoxyribonucleotides
Reversed-phase HPLC separation of diastereomeric phosphorothioate analogues of oligodeoxyribonucleotides and other backbone-modified congeners of DNA 326(1985)263

Phosphorus
Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251

O-Phospho-L-serine
Determination of O-phosphothreonine, O-phosphoserine, O-phosphotyrosine and phosphate by HPLC 338(1985)315

O-Phospho-L-threonine
Determination of O-phosphothreonine, O-phosphoserine, O-phosphotyrosine and phosphate by HPLC 338(1985)315

O-Phospho-L-tyrosine
Determination of O-phosphothreonine, O-phosphoserine, O-phosphotyrosine and phosphate by HPLC 338(1985)315

Photoionization detector
Temperature stabilization of a commercial photoionization detector. Performance evaluation 330(1985)365

***o*-Phthalaldehyde**
High-speed LC analysis of amino acids by post-column sodium hypochlorite-*o*-phthalaldehyde reaction 327(1985)17

Phthalates
Application of capillary GC in the stability control of phthalate-ester-type poly(vinyl chloride) plasticisers treated with light and plasma 329(1985)389

Determination of di- and mono(2-ethylhexyl) phthalate in plasma by GC 342(1985)399

Determination of di(2-ethylhexyl) phthalate and four of its metabolites in blood plasma by GC-MS 344(1985)167

Preseparation of polar, substituted, polycyclic aromatic hydrocarbons in aerosol extracts by HPLC on acidic- or basic-buffered silica 349(1985)331

Phthalic acid
Determination of phthalic acid, mono-(2-ethylhexyl) phthalate and di(2-ethylhexyl) phthalate in human plasma and in blood products 337(1985)279

Phyllanthoside
HPLC assay for the antitumor glycoside phyllanthoside and its stability in plasma of several species 342(1985)129

Phylloquinone
Quantitative analysis of phylloquinone (Vitamin K_1) in soy bean oils by HPLC 329(1985)257

Physcion
HPLC separation and quantitative determination of 1,8-dihydroxyanthraquinones in plant cell cultures 329(1985)311

Physostigmine
Sensitive LC method for physostigmine in biological fluids using dual-electrode electrochemical detection 341(1985)361

Phytanic acid
Simple and rapid method for the quantitation of plasma phytanic acid by GC 339(1985)375

Phytoalexins
HPLC of isoflavonoid phytoalexins from soybean (*Glycine max*, L. Harosoy 63) 333(1985)225

Pibenzimol
HPLC assay and preclinical pharmacological studies of pibenzimol (bisbenzimidazole) 341(1985)89

Picrates
Derivatization of alkyl halides and epoxides with picric acid salts for improved HPLC detection 333(1985)349

Pilocarpine
HPLC assay for pilocarpine in aqueous humor 336(1985)337

Pimpinellin
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Pindolol
Determination of pindolol in plasma and urine by TLC 342(1985)458

Pipecuronium bromide
Chromatography of bis-quaternary amino steroids. I. Separation on silica by TLC and HPLC 328(1985)279

Piperine
Determination of piperine in biological tissues by TLC and absorption densitometry 338(1985)259

Pirprofen
Simultaneous determination of pirprofen and its metabolite, the pyrrole derivative, in plasma by HPLC 339(1985)157

Plant hormones
Simultaneous determination of abscisic acid and jasmonic acid in plant extracts using HPLC 330(1985)347

Resoltuion of *RS*-abscisic acid and the separation of abscisic acid metabolites from plant tissue by HPLC 336(1985)221

Plasmids
Preparative isolation of plasmid DNA with sedimentation field flow fractionation 342(1985)67

Plasmin
Separation of human Glu-plasminogen, Lys-plasminogen and plasmin by high-performance affinity chromatography on Asahipak GS gel coupled with *p*-aminobenzamidine 348(1985)199

Plasminogen
Protein precipitation induced by a textile dye. Precipitation of human plasminogen in the presence of Procion Red HE3B 346(1985)111

Separation of human Glu-plasminogen, Lys-plasminogen and plasmin by high-performance affinity chromatography on Asahipak GS gel coupled with *p*-aminobenzamidine 348(1985)199

Plasticizers
Application of capillary GC in the stability control of phthalate-ester-type poly(vinyl chloride) plasticisers treated with light and plasma 329(1985)389

Plastoquinone-9
HPLC of coenzyme Q-related compounds and its application to biological materials 342(1985)35

Plate height equations
LC plate height equations 330(1985)159

Platelet-activating factor
Rapid isocratic procedure for the separation of platelet-activating factor from phospholipids in human saliva by HPLC 342(1985)380

Platelet glycoproteins
Tandem separation of labelled human blood platelet membrane glycoproteins by anion-exchange and gel fast protein LC 326(1985)179

Plate number
HPLC theory for the practitioner (Review) 334(1985)197

Platinum complexes
Phenol effect in PC of metal complexes 331(1985)273

Platydesminium
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Polar compounds
Large-scale preparative countercurrent chromatography for separation of polar compounds 348(1985)131

Polarity
Increments of some polarity parameters for polyoxyethylene glycol dialkyl ethers and for some of their sulphur analogues 330(1985)61

Polarity parameters
Polarity of polyoxyethylene glycol dialkyl ethers and some sulphur analogues measured by GC 333(1985)319

Polarographic detector
Thin-layer polarographic detector for the HPLC detection of thiourea derivatives 328(1985)127

Polar substances
Separation of natural polar substances by reversed-phase HPLC, centrifugal TLC and droplet counter-current chromatography 349(1985)69

Poliovirus proteins
Characterization of a solvent system for separation of water-insoluble poliovirus proteins by reversed-phase HPLC 326(1985)91

Poly(ADP-ribose)
HPLC of *in vitro* synthesized poly(ADP-ribose) on ion-exchange columns, separation of oligomers of varying chain length and estimation of apparent branching 330(1985)287

Polyallylamine
Polyallylamine-coated silica gel microbore column for LC 332(1985)15

Polyamines
Ion-pair HPLC determination of dansylated aliphatic polyamines with fluorescence and UV detection 333(1985)220

Separation of cadaverine from putrescine, histamine and polyamines in rat kidney by phosphocellulose chromatography 337(1985)103

Determination of polyamines and related compounds by reversed-phase HPLC: improved separation systems 339(1985)45

Mass fragmentographic identification of polyamine metabolites in the urine of normal persons and cancer patients, and its relevance to the use of polyamines as tumour markers 339(1985)223

Polybuffer exchanger matrices
Use of chromatofocusing for separation of β-lactamases. VI. Comparative studies on two chromatofocusing polybuffer exchanger matrices with the R46 plasmid coded β-lactamase carried by the *Escherichia coli* K12 J5-3 strain 333(1985)244

Polycarbonates
Determination of molecular weight data in size-exclusion chromatography by use of braod-molecular-weight-distributed standards 349(1985)357

Polychlorinated biphenyls
see PCBs

Polychlorinated dibenzo-*p*-dioxins
Synthesis and GC-MS determination of polychlorinated dibenzo-*p*-dioxins and related compounds in the technical chlorophenol formulation Ky-5 331(1985)410

Semi-preparative HPLC analysis of complex organic mixtures 332(1985)169

Polychlorinated dibenzofurans
Synthesis and GC-MS determination of polychlorinated dibenzo-*p*-dioxins and related compounds in the technical chlorophenol formulation Ky-5 331(1985)410

Semi-preparative HPLC analysis of complex organic mixtures 332(1985)169

Polychlorinated diphenyl ethers
Synthesis and GC-MS determination of polychlorinated dibenzo-*p*-dioxins and related compounds in the technical chlorophenol formulation Ky-5 331(1985)410

Polycyclic aromatic hydrocarbons
see Polynuclear aromatic hydrocarbons

Polycytidylic acid conjugates
Neonucleoproteins. Preparation and HPLC characterization of succinyl-lysozyme-diaminooctyl-polycytidylic acid and related polycytidylic acid conjugates 326(1985)311

Polyethylene glycol
Investigation of operating parameters in high-performance displacement chromatography 330(1985)1

Comparative GC studies of bonded and physically coated PEG 20M and its derivative phases with reference to some high-boiling isomeric solutes 346(1985)359

Contributions to hydrocarbon retention in gas-liquid-solid chromatography with polyethylene glycol 6000 as stationary phase 350(1985)15

Polyethylene oxide
Polyethylene oxide: a ligand for mild hydrophobic interaction chromatography? 347(1985)291

Polygodials
Analytical separation of enantiomeric polygodials by GC of pyrrole derivatives 347(1985)455

Polymers
New gel permeation column for the separation of water-soluble polymers 332(1985)39

Poly-γ-methyl-L-glutamate
Novel GPC packing composed of cross-linked and porous spherical particles of poly-γ-methyl-L-glutamate 347(1985)357

Polynuclear aromatic hydrocarbons
Class separation of polycyclic aromatic hydrocarbons, nitrogen heterocycles and hydroxyl aromatics by LC 331(1985)55

LC retention behavior of large, fused polycyclic aromatics. Normal bonded phases 346(1985)81

Determination of polynuclear aromatic hydrocarbons in petroleum oils by combination of TLC and GC-MS 348(1985)159

Preseparation of polar, substituted, polycyclic aromatic hydrocarbons in aerosol extracts by HPLC on acidic- or basic-buffered silica 349(1985)331

Polyols
Identification of 6-deoxyallitol and 6-deoxygulitol in human urine. Electron-impact mass spectra of eight isomers of 6-deoxyhexitol 336(1985)345

Elevation of certain polyols in the cerebrospinal fluid of patients with multiple sclerosis 336(1985)351

Polyoxyethylene glycol dialkyl ethers
Increments of some polarity parameters for polyoxyethylene glycol dialkyl ethers and for some of their sulphur analogues 330(1985)61

Polarity of polyoxyethylene glycol dialkyl ethers and some sulphur analogues measured by GC 333(1985)319

Polypeptides
HPLC of amino acids, peptides and proteins. LXVI Investigations on the effects of chromatographic dwell in the reversed-phase HPLC separation of proteins 327(1985)47

HPLC of amino acids, peptides and proteins. LXV Studies on the optimisation of the reversed-phase gradient elution of polypeptides: evaluation of retention relationships with β-endorphin-related polypeptides 327(1985)115

Quantitative affinity HPLC of neuroendocrine polypeptides using porous and non-porous glass derivatives 327(1985)193

Homogenous [mono-^{125}I-Tyr^{1}]- and [mono-^{125}I-Tyr^{13}] glucagon 331(1985)125

Purification of radioiodinated photoactivable glucagon by isocratic HPLC 331(1985)133

HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25

HPLC and studies of neurophysin-neurohypophysial hormone pathways 336(1985)63

Macroscale HPLC separation and instrumental identification of components of diethylaminoethyl murine epidermal growth factor 336(1985)73

Purification and characterization of CAMP-factor from *Streptococcus agalactiae* by hydrophobic interaction chromatography and chromatofocusing 348(1985)363

Direct liquid introduction MS of some underivatized dipeptides and polypeptides 350(1985)399

Polyphosphates
Effect of column temperature on HPLC behaviour of inorganic polyphosphates. I. Isocratic ion-exchange chromatography 348(1985)27

Effect of column temperature on HPLC behaviour of inorganic polyphosphates. II. Gradient ion-exchange chromatography 350(1985)119

Computer-assisted prediction of retention times for inorganic polyphosphates in gradient ion-exchange chromatography 350(1985)461

Polysaccharides
Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33

Separation of acidic polysaccharides from *Ulmus glabra* Huds. on Mono P™ 329(1985)315

Some applications of reversed-phase HPLC to oligosaccharide separations 336(1985)161

Comparative study of affinity chromatography of components of the hyaluronate-proteoglycan complex to immobilized hyaluronate 350(1985)227

GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Polysiloxanes
Soviet polysiloxane stationary phases for GC (Review) 334(1985)1

Polystyrene oligomers
Role of polar groups in fractionating isomeric polystyrene oligomers 347(1985)39

Polystyrenes
Swelling propensity (SP factor) of semi-rigid chromatographic packing materials 350(1985)145

Poly(triphenylmethyl methacrylate)
Chromatographic resolution. IX. HPLC resolution of enantiomers on optically active poly(triphenylmethyl methacrylate) 350(1985)127

Polyurethane elastomers
Chromatographic analysis of thermoplastic, medical-grade, polyether-based polyurethane elastomers 349(1985)347

Polyvalent cations
Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147

Poly(vinyl alcohol)
Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255

Porapak Q
Pre-concentration of organic pollutants. Potential interference from the use of styrene copolymer adsorbents 349(1985)181

Porous polymers
Correlation between chemical and physical characteristics and adsorption properties of some styrene copolymers 330(1985)55

Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Evaluation of the suitability of selected porous polymers for preconcentration of organosulphur compounds from water 346(1985)61

Reversed-flow GC technique applied to measurement of mass transfer coefficients of *n*-hydrocarbons on Porapak P 348(1985)339

Effect of the mesh-size distribution of porous polymer beads on efficiency and selectivity of stationary phases 349(1985)189

Porphyrin esters
Rapid isocratic HPLC of porphyrin esters on aminopropyl-bonded silica 333(1985)141

Porphyrins
Ion-pair HPLC separation of porphyrin isomers 349(1985)283

Computerised capillary GC-MS studies of the petroporphyrins of the Gilsonite bitumen (Eocene, U.S.A.) 350(1985)37

Post-column reactors
High-speed LC analysis of amino acids by post-column sodium hypochlorite-*o*-phthalaldehyde reaction 327(1985)17

Determination of anthraquinones in pulping materials by HPLC using online post-column derivatization 328(1985)309

Determination of catecholamines by HPLC with catalytic photometric detection 329(1985)147

Simultaneous determination of iron(II) and -(III) by ion chromatography with post-column reaction 329(1985)247

Off-line LC-MS studies of fluorescent β-aminothiol-*o*-phthalaldehyde derivatives 330(1985)307

Estimation of catecholamines by ion-exchange chromatography on Asahipak ES-502C, using glycylglycine as the post-derivatizing agent 332(1985)9

Determination of thiamine and its phosphate esters in human and rat blood by HPLC with post-column derivatization 332(1985)181

Determination of dopamine-3- and -4-O-sulphate in human plasma and urine by anion-exchange HPLC with fluorimetric detection 342(1985)261

HPLC determination of plasma and brain histamine without previous purification of biological samples: cation-exchange chromatography coupled with post-column derivatization fluorometry 344(1985)115

Automated determination of amoxycillin in biological fluids by column switching in ion-pair reversed-phase LC systems with post-column derivatization 344(1985)285

Determination of trace amounts of lanthanides in rocks and minerals by HPLC 349(1985)137

HPLC of anthranilate synthase using gel filtration and a post-column reactor 350(1985)332

Determination of phenols in water and wastewater by post-column reaction detection HPLC 350(1985)407

Potassium
Improvement of ion chromatography with UV photometric detection and comparison with conductivity detection for the determination of serum cations 341(1985)53

Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251

Prealbumin
Isotachophoresis as a useful tool for monitoring neurological complications of acute leukaemia in children 342(1985)285

Pre-column
Automated pre-column system for trace enrichment and clean-up of plasma samples in narrow-bore LC 330(1985)113

Preconcentration methods
Precolumn sample enrichment device for analysis of ambient volatile organics by GC-MS 329(1985)399

Studies on sample preconcentration in ion chromatography. II. An automated, single pump preconcentration system for non-suppressed ion chromatography with conductivity detection 330(1985)95

Prednisolone
HPLC assay of anti-inflammatory drugs incorporated in gel ointments. Separation and stability testing 331(1985)383

Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Preparative chromatography
Preparative high-performance ion-exchange chromatography 333(1985)29

Preparative high-performance hydrophobic interaction chromatography of proteins on TSKgel Phenyl-5PW 333(1985)202

Preparative HPLC of (2*E*)- and (2*Z*)-farnesol 347(1985)188

Large-scale preparative countercurrent chromatography for separation of polar compounds 348(1985)131

Pressure drop
HPLC theory for the practitioner (Review) 334(1985)197

Contribution to the study of micropacked columns in GC 348(1985)327

Pressure-modifier gradients
Gradient programming and combined gradient-pressure programming in supercritical fluid chromatography 349(1985)317

Primidone
Concurrent determination of valproic acid with other antiepileptic drugs by HPLC 338(1985)131

Determination of antiepileptic drugs (Review) 340(1985)199

Principal components analysis
Qualitative organic analysis. I. Identification of drugs by principal components analysis of standardized TLC data in four eluent systems 350(1985)151

Proadifen
HPLC separation and determination of 2-(diethylamino)ethyl diphenylpropylacetate and its metabolites from biological samples 342(1985)135

Procaine
Determination of sodium, ephedrine and procaine in pharmaceuticals by capillary isotachophoresis 330(1985)436

Procion Red HE3B
Protein precipitation induced by a textile dye. Precipitation of human plasminogen in the presence of Procion Red HE3B 346(1985)111

Proctolin
Exopeptidase-HPLC peptide mapping of small peptides 326(1985)399

Profens
Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77

Progabide
Determination of antiepileptic drugs (Review) 340(1985)199

Determination of progabide and its acid metabolite in biological fluids by HPLC on silica column with UV detection 344(1985)231

Progesterone
Reversed-phase HPLC of C_{21} metabolites of progesterone 336(1985)361

Determination of progesterone by LC-MS using a moving-belt interface and isotope dilution 343(1985)397

Prolidase
Simultaneous measurement of prolidase and prolinase activities in erythrocytes using an isotachophoretic analyser 343(1985)179

Prolinase
Simultaneous measurement of prolidase and prolinase activities in erythrocytes using an isotachophoretic analyser 343(1985)179

Proline
Simplified procedure for specific determination of imino acids in human blood plasma by HPLC 339(1985)59

HPLC analysis of free hydroxyproline and proline in blood plasma and of free and peptide-bound hydroxyproline in urine 339(1985)285

Promethazine
Determination of promethazine in human plasma by automated HPLC with electrochemical detection and by GC-MS 339(1985)105

Prometryn
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Pro-opiomelanocortin
Use of reversed-phase and ion-exchange batch extraction in the purification of bovine pituitary peptides 326(1985)329

Propafenone
Determination of plasma protein binding of propafenone in rats, dogs and humans by highly sensitive GC-MS 341(1985)305

Propazine
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Propellants
Microcolumn clean-up and recovery techniques for organic explosives compounds and for propellants traces in firearms discharge residues 330(1985)121

Propisomide
Determination of propisomide, a new antiarrhythmic agent, in biological samples by GC with a thermionic detector 341(1985)202

Propoxyphylline
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Propranolol
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 341(1985)213

Determination of drugs in untreated body fluids by micellar chromatography with fluorescence detection 342(1985)293

Rapid and simultaneous extraction of propranolol, its neutral and basic metabolites from plasma and assay by HPLC 343(1985)349

Proquazone
Determination of proquazone and its *m*-hydroxy metabolite by HPLC. Clinical application: pharmacokinetics of proquazone in children with juvenile rheumatoid arthritis 341(1985)105

Prostaglandins
HPLC assay of anti-inflammatory drugs incorporated in gel ointments. Separation and stability testing 331(1985)383

Selective trimethylsilylation for the determination of 16(*S*)-amino-$PGF_{2\alpha}$ methyl ester by GLC 333(1985)404

Simultaneous determination of prostanoids in plasma by GC-negative-ion chemical-ionization MS 338(1985)273

Assay methods for 6-keto-prostaglandin $F_{1\alpha}$ in human urine. Comparison of chromatographic techniques with radioimmunoassay and GC-

negative-ion chemical-ionization MS 339(1985)1

Contribution to the measurement of prostaglandins by GC-MS 342(1985)245

Determination of (15*R*)- and (15*S*)-15-methylprostaglandin E_2 in human plasma with picogram per milliliter sensitivity by column-switching HPLC 343(1985)271

Qualitative and quantitative measurement of hydroxy fatty acids, thromboxanes and prostaglandins using stable isotope dilutions and detection by GC-MS 344(1985)11

Assay of prostaglandins in the epithelial cells and fibroblasts of the rat mammary gland 345(1985)380

Influence of the mobile phase composition on the reversed-phase TLC behaviour of a series of prostaglandins. Comparison of the extrapolated R_M values 347(1985)209

Silica capillary GC of prostaglandins with electron-capture detection and its application to the forensic investigation of sexual offences 348(1985)111

HPLC separation and determination of prostaglandins, oxidized by pyridinium dichromate. Optimization and applications 349(1985)431

Prostanoids
Simultaneous determination of prostanoids in plasma by GC-negative-ion chemical-ionization MS 338(1985)273

Protease
Purification of gelatin-specific neutral protease from human skin by conventional and high-performance LC 326(1985)147

Protein kinase
Combined recycling affinity column-filtration technique 346(1985)247

Proteins
see also Enzymes

Theoretical investigation of the optimum particle size for the resolution of proteins by size-exclusion chromatography 326(1985)3

Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33

A new, stabilized, hydrophilic silica packing for the high-performance gel chromatography of macromolecules 326(1985)63

Characterization of a solvent system for separation of water-insoluble poliovirus proteins by reversed-phase HPLC 326(1985)91

Separation of flavivirus membrane and capsid proteins by multistep HPLC optimized by immunological monitoring 326(1985)113

HPLC separation of distinct epidermal cell-derived cytokines 326(1985)129

Combined size-exclusion and reversed-phase HPLC of a detergent extract of Sendai virus 326(1985)173

Analysis of nucleoproteins by direct injection of dissolved nuclei or chromosomes into a HPLC system 326(1985)199

Isolation of human factor VIII:C by preparative high-performance size-exclusion chromatography 326(1985)217

Preparation of sheep and cattle immunoglobulins with antibody activity by HPLC 326(1985)225

Purification of human serum gamma globulins by hydrophobic interaction HPLC 326(1985)235

Isolation of detergent-solubilized monomers of bacteriorhodopsin by size-exclusion HPLC 326(1985)243

General method for the separation of cyanogen bromide digests of proteins by HPLC. Rabbit skeletal troponin I 326(1985)349

Characterization of recombinant human interleukin-2 with micromethods 326(1985)357

Purification and peptide mapping of calmodulin and its chemically modified derivatives by reversed-phase HPLC 326(1985)387

Automated tandem HPLC system for separation of extremely complex peptide mixtures 326(1985)407

Comparison of reversed-phase columns for the separation of tryptic peptides by gradient elution. Correlation of experimental results and model prediction 326(1985)419

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27

HPLC of amino acids, peptides and proteins. LXVI Investigations on the effects of chromatographic dwell in the reversed-phase HPLC separation of proteins 327(1985)47

Vizualization of protein retention and migration in reversed-phase LC 327(1985)67

Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Application to method development and the choice of column configuration 327(1985)93

Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Chromatographic study of the interrelationships of immunoglobulin A and α_1-microglobulin in myelomatosis 327(1985)179

Automated high-performance immunosorbent assay for recombinant leukocyte A interferon 327(1985)189

Isolation and quantitation of serum IgE levels by high-performance immunoaffinity chromatography 327(1985)205

Protein A-coated glass beads. Universal support medium for high-performance immunoaffinity chromatography 327(1985)213

Isolation of unliganded steroid receptor proteins by high-performance size-exclusion chromatography. An investigation of steroid-dependent structural alterations 327(1985)221

High-performance hydrophobic-interaction chromatography of steroid hormone receptors 327(1985)237

High-performance chromatofocusing of steroid receptor proteins in the presence and absence of steroid. Investigation of steroid-dependent alterations in surface charge heterogeneity 327(1985)247

Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269

Methods for chromatofocusing of cerebrospinal fluid and serum immunoglobulin G 327(1985)293

Determination of the dissociation constant of oligomeric proteins by size-exclusion HPLC: application to human haemoglobin 327(1985)313

Detection of quality differences among wheats by HPLC 327(1985)333

Analysis of different forms of recombinant human leukocyte interferons and synthetic fragments by HPLC 327(1985)359

Rapid, high-resolution procedure for assessment of estrogen receptor heterogeneity in clinical samples 327(1985)369

Detection of Sendai virus protein by reversed-phase HPLC combined with immuno-chromatography 327(1985)377

Adsorption of SH-containing amino acids, peptides and proteins on the 2-nitrophenylhydrazonopropanedinitrile derivative of Spheron Ara 328(1985)402

Separation and isolation of human myelin basic protein peptides by reversed-phase HPLC in volatile buffers 329(1985)295

Electrophoretic and chromatographic techniques for the differential diagnosis of a haemoglobin abnormality: Hb E heterozygosity 330(1985)299

Affinity partitioning of biopolymers and membranes in Ficoll-dextran aqueous two-phase systems 331(1985)11

Effects of solutions used for storage of size-exclusion columns on subsequent chromatography of peptides and proteins 331(1985)253

HPLC studies on non-mercapt mercapt conversion of human serum albumin II 332(1985)19

Ion-exchange HPLC studies on sulphydryl-catalysed structural alterations of bovine mercaptalbumin 332(1985)29

Recent progress in gel packing materials and detectors for modern LC in Japan 332(1985)227

Preparative high-performance ion-exchange chromatography 333(1985)29

Resin-based support for reversed-phase chromatography of proteins 333(1985)93

Preparative high-performance hydrophobic interaction chromatography of proteins on TSKgel Phenyl-5PW 333(1985)202

Chromatographic determination of N-acetyl-DL-tryptophan and octanoic acid in human albumin solutions 333(1985)381

Recent developments in isotachophoresis (Review) 334(1985)157

HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25

Anomalous behavior of lectins in size-exclusion HPLC and gel electrophoresis 336(1985)397

Efficient purification of human plasma β_2-microglobulin from the haemodialysate of a patient with chronic renal failure by use of HPLC 339(1985)67

Studies of fish zona pellucida by high-performance ion-exchange chromatography on agarose columns and free zone electrophoresis 341(1985)295

Isotachophoresis as a useful tool for monitoring neurological complications of acute leukaemia in children 342(1985)285

Detection of the changes in cellular proteins in regenerating rat liver by high-resolution two-dimensional electrophoresis 343(1985)51

Isolation of normal and variant human hemoglobin subunits 343(1985)168

Ion-exchange HPLC purification of bovine cardiac and rabbit skeletal muscle troponin subunits 343(1985)315

New less temperature-sensitive microchromatographic method for the separation and quantitation of glycosylated

hemoglobins using a non-cyanide buffer system 344(1985)81

Analysis and preparation of chromosomal high-mobility group proteins by ion-exchange HPLC 344(1985)332

Protein precipitation induced by a textile dye. Precipitation of human plasminogen in the presence of Procion Red HE3B 346(1985)111

Analytical isoelectric focusing of native and modified haemoglobins after treatment with 4-hydroxymercuribenzoate 346(1985)401

Application of DNA-cellulose chromatography in the isolation of immunoglobulin M and complement component C4b-binding protein from human serum 347(1985)129

HPLC of proteins. Rapid chromatofocusing of plant enzymes 348(1985)167

Rapid analysis of bovine milk proteins by fast protein LC 348(1985)177

Separation of human Glu-plasminogen, Lys-plasminogen and plasmin by high-performance affinity chromatography on Asahipak GS gel coupled with *p*-aminobenzamidine 348(1985)199

Reversed-phase HPLC of insulin and insulin derivatives. A comparative study 348(1985)347

Identification and isolation of human insulin A and B chains by HPLC 349(1985)91

High-performance affintiy chromatography of anhydrochymotrypsin on a hydrophilic vinyl-polymer gel coupled with tryptophan 350(1985)265

Ion-exchange columns in HPLC of proteins; a simple loading technique that improves resolution 350(1985)299

Rapid delipidation of the membrane protein bacteriorhodopsin by HPLC 350(1985)427

Protein sequencer
Automated on-line identification of phenylthiohydantoin-amino acids from a vapor phase protein sequencer 350(1985)217

Proteoglycans
Comparative study of affinity chromatography of components of the hyaluronate-proteoglycan complex to immobilized hyaluronate 350(1985)227

Proton affinities
Gaseous acidity and basicity scales as guides to chemical ionization in reversed-phase LC-MS with direct liquid introduction 347(1985)61

Protophomin
TLC of mycotoxins (Review) 334(1985)211

Proxiphomin
TLC of mycotoxins (Review) 334(1985)211

PR toxin
TLC of mycotoxins (Review) 334(1985)211

Pseudomonas aeruginosa
Availability of 6-hydroxynicotinic acid for rapid identification of *Pseudomonas aeruginosa* and *Serratia marcescens* 338(1985)51

Pseudouridine
Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Psoralen
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Pteridines
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

Pterins
Estimation of tetrahydro, dihydro and fully oxidised pterins by HPLC using sequential electrochemical and fluorometric detection 343(1985)35

PTH derivatives
Microscale isocratic separation of phenylthiohydantoin amino acid derivatives 326(1985)321

Puerarin
Analysis of isoflavones in *Puerariae radix* by HPLC with amperometric detection 347(1985)438

Purine nucleoside phosphorylase
Simultaneous HPLC assay of the activities of erythrocytic hypoxanthine-guanine phosphoribosyl transferase and purine nucleoside phosphorylase 349(1985)275

Purines
see also Nucleobases

Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33

Recent developments in isotachophoresis (Review) 334(1985)157

HPLC method for the analysis of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in biological fluids 339(1985)380

Use of a fundamental elution protocol for the development of reversed-phase HPLC enabling rapid simultaneous determination of purines, pyrimidines and allied compounds commonly found in human biological fluids 344(1985)101

Evaluation of combined electrochemical and UV detection in the HPLC analysis of plasma for low-molecular-weight constituents 345(1985)125

Putreanine
Determination of polyamines and related compounds by reversed-phase HPLC: improved separation systems 339(1985)45

Determination of amino acids by separation of their ion pairs with dodecyl sulphate 341(1985)11

Putrescine
Separation of cadaverine from putrescine, histamine and polyamines in rat kidney by phosphocellulose chromatography 337(1985)103

Frequency-pulsed electron-capture GLC studies of chemical changes in sera of patients with schistosomiasis 339(1985)243

Pyrazinamide
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Rapid determination of pyrazinamide in biological fluids by HPLC 345(1985)453

Pyrazoloimidazole
GC determination of pyrazoloimidazole 343(1985)190

Pyrethroids
Coupling a high-performance liquid chromatograph with a liquid scientillation detector: example in the field of pyrethroids 349(1985)105

High-performance liquid chromatograph coupled with two detectors: a UV spectrometer and a polarimeter. Example in the field of pyrethroids: identification of enantiomers 349(1985)109

Pyridinium dichromate
HPLC separation and determination of prostaglandins, oxidized by pyridinium dichromate. Optimization and applications 349(1985)431

Pyridoxal
Quantities of B_6 vitamers in human milk by HPLC. Influence of maternal vitamin B_6 status 337(1985)249

Stability of pyridoxal-5-phosphate semicarbazone: applications in plasma vitamin B_6 analysis and population surveys of Vitamin B_6 nutritional status 342(1985)277

Pyridoxal-5-phosphate
Stability of pyridoxal-5-phosphate semicarbazone: applications in plasma vitamin B_6 analysis and population surveys of Vitamin B_6 nutritional status 342(1985)277

Pyridoxamine
Quantities of B_6 vitamers in human milk by HPLC. Influence of maternal vitamin B_6 status 337(1985)249

Pyridoxine
Quantities of B_6 vitamers in human milk by HPLC. Influence of maternal vitamin B_6 status 337(1985)249

3-(3-Pyridyl)-1-propanol
Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

Pyrimidine 5'-nucleotidase
Assay of human erythrocyte pyrimidine and deoxypyrimidine 5'-nucleotidase by isocratic reversed-phase HPLC 339(1985)293

Pyrimidine photohydrates
HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33

Pyrimidines
see also Nucleobases

HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33

Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33

Recent developments in isotachophoresis (Review) 334(1985)157

HPLC method for the analysis of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in biological fluids 339(1985)380

Use of a fundamental elution protocol for the development of reversed-phase HPLC enabling rapid simultaneous determination of purines, pyrimidines and allied compounds commonly found in human biological fluids 344(1985)101

Pyrimidonesilicones
Improved method for the preparation of immobilised pyrimidonesilicones as polar stationary phases used for the capillary column GC separation of underivatised barbiturates 329(1985)213

Pyrrolidinone
Determination of pyrrolidinone in plasma by HPLC 344(1985)125

5-Pyrrolidone-2-carboxylic acid
HPLC with fluorimetric detection in biological tissues of the 4-bromomethyl-7-methoxycoumarin ester derivative of 5-pyrrolidone-2-carboxylic acid 344(1985)325

Pyrrolizidine alkaloids
Determination of pyrrolizidine alkaloids in *Senecio inaequidens* D.C. by capillary GC 349(1985)23

PZ-peptidase
Highly sensitive assay for PZ-peptidase activity by HPLC 348(1985)205

Q

Quinic acid
HPLC of shikimate pathway intermediates 329(1985)268

Quinidine
Determination of drugs in untreated body fluids by micellar chromatography with fluorescence detection 342(1985)293

Quinine
Determination of drugs in untreated body fluids by micellar chromatography with fluorescence detection 342(1985)293

Cinchona alkaloids for preparing new, easily accessible chiral stationary phases. II. Resolution of binaphthol derivatives on silica-supported quinine 348(1985)79

Quinoxalinols
Quantification of branched-chain α-keto acids as quinoxalinols: importance of excluding oxygen during derivatization 337(1985)98

R

Radioactive compounds
Monitoring radioactive compounds in HPLC eluates: fraction collection versus on-line detection 342(1985)25

Radioassay
High-sensitivity radioassay of chromatographic effluents. Automatic fraction collector/concentrator for quantitative autoradiography 349(1985)267

Radiolabeled anions
Chromatography of radiolabelled anions using reversed-phase LC columns 333(1985)269

Raffinose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Rare-earth elements
Determination of trace amounts of lanthanides in rocks and minerals by HPLC 349(1985)137

RDX
Reversed-phase gradient HPLC of nitramine munitions and characterization of munitions process samples by GC-MS 329(1985)285

Microcolumn clean-up and recovery techniques for organic explosives compounds and for propellants traces in firearms discharge residues 330(1985)121

α-Receptor blocking agents
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

β-Receptor stimulating agents
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Red dyes
Identification of natural red dyes in old Indian textiles. Evaluation of TLC systems 347(1985)447

Redox chemiluminescence detector
Selectivity of the redox chemiluminescence detector for complex sample analysis 349(1985)395

Re-injection method
The re-injection method for recovery of components adsorbed during syringe gas sampling: a statistical evaluation 329(1985)119

Relaxins
HPLC of amino acids, peptides and proteins. LXIII. Reversed-phase HPLC characterisation of several polypeptide and protein hormones 336(1985)25

Reserpine
Sub-nanogram analysis of yohimbine and related compounds by HPLC 342(1985)333

Resin acids
Improved solvent-extraction based procedure for the GC analysis of resin and fatty acids in pulp mill effluents 346(1985)205

Resol
Separation and identification of phenol-formaldehyde condensates by GC-MS. II. Base-catalysed condensation products 331(1985)91

Separation and identification of phenol-formaldehyde condensates by GC-MS. III. Hemiformal isomers in resol 333(1985)161

Resorcinol
Determination of metabolites of cytochrome P-450 model systems using HPLC 341(1985)373

Response factors
Quantitative responses of normal alkanes in GC-MS analysis 350(1985)27

Retention behaviour
Vizualization of protein retention and migration in reversed-phase LC 327(1985)67

HPLC of amino acids, peptides and proteins. LXV Studies on the optimisation of the reversed-phase gradient elution of polypeptides: evaluation of retention relationships with β-endorphin-related polypeptides 327(1985)115

Influence of the biological matrix on retention behaviour in TLC: evidence of systematic differences between pure and extracted drugs 342(1985)241

LC retention behavior of organometallic compounds and ligands with amine-, octadecylsilica- and β-cyclodextrin-bonded phase columns 347(1985)51

Effect of column temperature on HPLC behaviour of inorganic polyphosphates. I. Isocratic ion-exchange chromatography 348(1985)27

Ligand-exchange GC of dialkyl sulphides 350(1985)63

Retention gaps
On-column injection of large sample volumes using the retention gap technique in capillary GC (Review) 334(1985)129

Retention indices
Comparison of mathematical methods for the calculation of Kováts indices 328(1985)11

Temperature-programmed retention indices of 221 halogenated organic compounds with 1-bromoalkanes as references 328(1985)35

GLC analyses. XLII. Retention behaviour of C_1-C_{12} *n*-alkyl esters of benzoic and pentafluorobenzoic acids on SE-30 and OV-351 capillary columns 328(1985)325

GLC analyses. XLIII. Retention increments for 2-chloro-, 2,2-dichloro- and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns 329(1985)43

GLC analyses. XLIV. Benzoyl and pentafluorobenzoyl esters of lower saturated branched-chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 329(1985)359

Retention indices as a tool for the prediction of chromatographic conditions for achieving the separation of mixtures of organic compounds 330(1985)71

Retention indices of dimethylbenzamidines and benzylideneamines on a non-polar column 330(1985)79

Correlation of retention indices obtained with two temperature programmes 330(1985)167

Vapor pressure estimates of individual polychlorinated biphenyls and commercial fluids using GC retention data 330(1985)203

GC retention indices of chemical warfare agents and simulants 331(1985)47

Kováts retention indices of high-molecular-weight monomethyl-, cyclopentyl-, cyclohexyl- and phenylalkanes 331(1985)219

Calculation of retention indices in temperature-programmed capillary GC 331(1985)229

Capillary GC of chloro derivatives of 1,4-dimethylbenzene. Separation, identification and prediction of boiling points 331(1985)237

Retention indices of some ergopeptines in reversed-phase HPLC 331(1985)245

A problem with Kováts' retention index equation 331(1985)389

GC of bicyclo[*n*.4.0]alken-2-ones 331(1985)406

Indirect determination of specific retention volumes in GC. Use of the retention indices of McReynolds' samples 333(1985)1

Relationships between GLC behaviour and structure of arylaliphatic acids 333(1985)21

Correlation of the chromatographic retention of some phenylacetic and phenylpropionic acid derivatives with molecular structure 333(1985)309

Polarity of polyoxyethylene glycol dialkyl ethers and some sulphur analogues measured by GC 333(1985)319

Estimation of dead time and calculation of Kováts indices (Review) 334(1985)95

Use of dual-column fused-silica capillary GC in combination with detector response factors for analytical toxicology 341(1985)81

Quantum chemical parameters in correlation analysis of GLC retention indices of amines 346(1985)53

Unified retention index of hydrocarbons separated on squalane 347(1985)366

Chromatography of monomers. IV. GLC studies of C_1-C_6 *n*-alkyl and C_3-C_6 isoalkyl acrylates and their hydrogen halide and halogen addition derivatives 348(1985)141

Retention mechanisms
Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. II. Influence of organic co-solvents on retention at 25°C 333(1985)41

LC retention behavior of organometallic compounds and ligands with amine-, octadecylsilica- and β-cyclodextrin-bonded phase columns 347(1985)51

Retention models
Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Verification of band width relationships for acetonitrile-water mobile phases 327(1985)27

HPLC of amino acids, peptides and proteins. LXVI Investigations on the effects of chromatographic dwell in the reversed-phase HPLC separation of proteins 327(1985)47

Optimization model for the gradient elution separation of peptide mixtures by reversed-phase HPLC. Application to method development and the choice of column configuration 327(1985)93

Adsorption effects in GLC. Solute retention in the hydrocarbon solute-polar stationary phase (polypropylene glycol 2000) system 330(1985)193

Affinity chromatographic examination of a retention model for macromolecules 331(1985)1

Stoichiometric displacement of solvent by non-polar solutes in reversed-phase LC 332(1985)147

Correlations between retention on liquid crystalline stationary phases and chemical structure. I. Dimethylnaphthalenes 333(1985)11

Proposed model for peak splitting in reversed-phase ion-pair HPLC with computer prediction of eluted peak profiles 336(1985)15

Retention prediction
Energy effects in the retention of aromatic acids in LC 332(1985)189

Utilization of the functional group contribution concept in LC on chemically bonded reversed phases 348(1985)49

Effect of column temperature on HPLC behaviour of inorganic polyphosphates. II. Gradient ion-exchange chromatography 350(1985)119

Separation of some peptides and related isopeptides by HPLC: structure-retention time relationships 350(1985)279

Computer-assisted prediction of retention times for inorganic polyphosphates in gradient ion-exchange chromatography 350(1985)461

Retention time
HPLC theory for the practitioner (Review) 334(1985)197

Retention volume
Adsorption effects in GLC. Solute retention in the hydrocarbon solute-polar stationary phase (polypropylene glycol 2000) system 330(1985)193

HPLC theory for the practitioner (Review) 334(1985)197

Retention volumes
Indirect determination of specific retention volumes in GC. Use of the retention indices of McReynolds' samples 333(1985)1

Retinaloximes
HPLC and spectroscopic characterization of stereoisomeric retinaloximes. Improvements in resolution and implications of the method 331(1985)265

Retinoids
Simple HPLC method for the separation of retinoids including N-(4-hydroxyphenyl)-all-*trans*-retinamide 341(1985)187

Rapid analysis of the major classes of retinoids by step gradient reversed-phase HPLC using retinal (O-ethyl) oxime derivatives 348(1985)241

Retinol
Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 345(1985)275

Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285

Retinol binding protein
Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269

Rhein
HPLC separation and quantitative determination of 1,8-dihydroxyanthraquinones in plant cell cultures 329(1985)311

Ribalinium
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Riboflavin
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

Ribonucleosides
Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33

HPLC method for the analysis of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in biological fluids 339(1985)380

Preparation of trimethylsilyl derivatives of ribonucleosides for GC 348(1985)280

Rifampicin
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Antibiotic monitoring in body fluids (Review) 340(1985)361

R_M values
Comparison of relationships between the chemical structures and mobilities of hyaluronate oligosaccharides in TLC and HPLC 328(1985)73

Influence of the mobile phase composition on the reversed-phase TLC behaviour of a series of prostaglandins. Comparison of the extrapolated R_M values 347(1985)209

tRNA
HPLC of tRNAs on novel stationary phases 326(1985)79

Roquefortine
TLC of mycotoxins (Review) 334(1985)211

Rubeanic acid
Rubeanic acid: a general spray reagent for the detection of steroids on thin-layer plates 350(1985)468

Rubratoxins
TLC of mycotoxins (Review) 334(1985)211

Ryosidine
Determination of ryosidine in human plasma by GLC 343(1985)427

S

Saccharides
see also Sugars and carbohydrates

Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Characterization of lacto-N-hexaose and two fucosylated derivatives from human milk by HPLC and proton NMR spectroscopy 328(1985)259

Saccharin
Studies on the analysis of food additives by HPLC. V. Simultaneous determination of preservatives and saccharin in foods by ion-pair chromatography 346(1985)333

Saccharose
HPLC of carbohydrates, alcohols and diethylene glycol on ion-exchange resins 350(1985)381

Salbutamol
Determination of salbutamol in human plasma and urine by HPTLC 345(1985)111

Salicylamide
Analysis of antipyretics by semimicro LC 332(1985)269

Salicylates
HPLC determination of certain salicylates and their major metabolites in plasma following topical administration of a liniment to healthy subjects 342(1985)315

Separation and identification of non-steroidal antirheumatic drugs containing a free carboxyl function using HPLC 345(1985)77

Salicylic acid
Determination of paracetamol, salicylic acid and acetyl salicylic acid in serum by HPLC 329(1985)196

Analysis of antipyretics by semimicro LC 332(1985)269

Application of thin-layer stick chromatographic identification test methods to drugs contained in preparations in the Pharmacopoeia of Japan 333(1985)115

Salicylic acids
HPLC determination of contaminants in salicylic acids after computer-assisted optimization of the separation parameters (in German) 350(1985)471

Salicyluric acid
HPLC determination of certain salicylates and their major metabolites in plasma following topical administration of a liniment to healthy subjects 342(1985)315

Salsolinol
Comparison of HPLC with electrochemical detection and GC-mass fragmentography for the assay of salsolinol, dopamine and dopamine metabolites in food and beverage samples 336(1985)199

Identification and quantification of 1-carboxysalsolinol and salsolinol in biological samples by GC-MS 343(1985)1

Sample enrichment
Precolumn sample enrichment device for analysis of ambient volatile organics by GC-MS 329(1985)399

Sample plug
Rejection of the sample plug after on-column injection into capillaries at high temperatures 333(1985)156

Sample volumes
On-column injection of large sample volumes using the retention gap technique in capillary GC (Review) 334(1985)129

Sampling efficiencies
Comparison between theoretical and experimental sampling efficiencies on Tenax GC 331(1985)207

Sampling methods
The re-injection method for recovery of components adsorbed during syringe gas sampling: a statistical evaluation 329(1985)119

Use of bonded-phase silica sorbents for rapid sampling of impurities in illicit amphetamine for HPLC analyses 331(1985)339

Sauvagines
Reversed-phase HPLC of natural and synthetic sauvagines 346(1985)390

Schiff base chelates
Applications of fused-silica wall-coated open-tubular columns. I. Schiff base chelates 350(1985)353

Scintillator
Radioactivity detection system with a CaF_2 (Eu) scintillator for HPLC 333(1985)83

Scutellariae Radix
Simultaneous determination of baicalein, wogonin, oroxylin-A and their glucuronides in Scutellariae Radix by ion-pair HPLC 328(1985)289

Secalonic acids
TLC of mycotoxins (Review) 334(1985)211

Secoiridoid glycosides
Analytical and preparative reversed-phase LC of secoiridoid glycosides 346(1985)396

Secretin
Rapid, small-scale preparation of gastrointestinal hormones by HPLC on a C_{18} column 326(1985)121

Sedative hypnotics
Drug level monitoring: Sedative hypnotics (Review) 340(1985)139

Sedimentation field flow fractionation
Preparative isolation of plasmid DNA with sedimentation field flow fractionation 342(1985)67

Selectivity
Effects of test solutes on the characterization of stationary phases in GC using a Snyder triangle 329(1985)5

Selenate
Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Selenite
Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Sendai virus proteins
Combined size-exclusion and reversed-phase HPLC of a detergent extract of Sendai virus 326(1985)173

Detection of Sendai virus protein by reversed-phase HPLC combined with immuno-chromatography 327(1985)377

Sepharose CL-4B
Three independent methods for quantitative determination of octyl covalently coupled to Sepharose CL-4B 346(1985)255

Sepiapterin
Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

Use of reversed-phase C_{18} Sep-Pak cartridges for the purification and concentration of sepiapterin and other pteridines 350(1985)389

Serine
Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

Serine carboxypeptidase
Purification of an acid protease and a serine carboxypeptidase from *Aspergillus niger* using metal-chelate affinity chromatography 329(1985)165

Serotonin
3-Methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, and homovanillic acid in human cerebrospinal fluid. Storage and measurement by reversed-phase HPLC and coulometric detection using 3-methoxy-4-hydroxyphenyllactic acid as an internal standard 336(1985)259

Determination of serotonin in plasma by LC with electrochemical detection 341(1985)445

Alkyl boronates as catechol-specific mobile phase pairing agents. Application of HPLC analysis of amines, precursors and metabolites in brain tissue 342(1985)370

Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Determination of the biogenic amines and their major metabolites in single human brain tissue samples using a combined extraction procedure and HPLC with electrochemical detection 345(1985)33

Simplified determination of the brain catecholamines norepinephrine, 5-hydroxyindoleacetic acid, dopamine and 5-hydroxytryptamine by HPLC using electrochemical detection 345(1985)140

Serratia marcescens
Availability of 6-hydroxynicotinic acid for rapid identification of *Pseudomonas*

aeruginosa and *Serratia marcescens* 338(1985)51

Sesquiterpene alcohols
General method for the HPLC prefractionation of essential oils and flavor mixtures for GC-MS analysis. Identification of new constituents in cold pressed lime oil 330(1985)141

SEX
Reversed-phase gradient HPLC of nitramine munitions and characterization of munitions process samples by GC-MS 329(1985)285

Shikimate pathway intermediates
HPLC of shikimate pathway intermediates 329(1985)268

Shikimate 3-phosphate
HPLC of shikimate pathway intermediates 329(1985)268

Shikimic acid
HPLC of shikimate pathway intermediates 329(1985)268

Sialic acids
Structural analysis of underivatized sialic acids by combined HPLC-MS 337(1985)231

Sialyl-α(2-3)-lactose
Structural analysis of underivatized sialic acids by combined HPLC-MS 337(1985)231

Silane monomers
Use of silane monomers for molecular imprinting and enzyme entrapment in polysiloxane-coated porous silica 347(1985)11

Silica gel
Long-chain alkyl grafts and mixed alkyl-alkane layers at the surface of macroporous silicas. Their GC properties below and above the phase transition 329(1985)331

Quality criteria and structure of silica gel column packing material 329(1985)351

Adsorption chromatographic separations on buffered silica gel (Review) 334(1985)35

Simazine
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Simulants
GC retention indices of chemical warfare agents and simulants 331(1985)47

Sisomicin
Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Skimmianine
HPLC separation of furo- and dihydrofuroquinolinic alkaloids on octadecyl-silica. Determination in *in vitro* cultures of Rutaceae (in French) 331(1985)437

Snyder triangle
Effects of test solutes on the characterization of stationary phases in GC using a Snyder triangle 329(1985)5

Sodium
Determination of sodium, ephedrine and procaine in pharmaceuticals by capillary isotachophoresis 330(1985)436

Improvement of ion chromatography with UV photometric detection and comparison with conductivity detection for the determination of serum cations 341(1985)53

Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251

Sodium hypochlorite
High-speed LC analysis of amino acids by post-column sodium hypochlorite-*o*-phthalaldehyde reaction 327(1985)17

Solanidine glycosides
Determination of the total C_{27}-steroidal alkaloid composition of *Solanum* species by high resolution GC 331(1985)285

Solasodine
Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

Solubility parameter optimization
Solubility parameter and window diagram optimization methods for the reversed-phase chromatographic optimization of complex hydroxyl aromatic mixtures 328(1985)81

Solu-medrol
Efficient chromatographic fractionation of steroids in human serum through regulation of liquid-liquid distribution ratios 339(1985)253

Solution thermodynamics
Possibility and consequences of using different concentration scales in the study of solution thermodynamics by GLC 350(1985)339

Solvent displacement
Stoichiometric displacement of solvent by non-polar solutes in reversed-phase LC 332(1985)147

Solvent disturbance peaks
Theory of solvent disturbance peaks and experimental determination of thermodynamic dead-volume in column LC 349(1985)211

Solvent effects
Computer simulation of abnormal high-performance liquid chromatograms caused by solvents 329(1985)13

Examination of the effect of solvent composition on bonded phase LC packings by ^{13}C Fourier transform NMR spectroscopy 331(1985)23

Somatomedin inhibitors
Separation of somatomedins and somatomedin inhibitors by size exclusion HPLC 339(1985)388

Somatomedins
Separation of somatomedins and somatomedin inhibitors by size exclusion HPLC 339(1985)388

Sorbents
Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Sorbic acid
Studies on the analysis of food additives by HPLC. V. Simultaneous determination of preservatives and saccharin in foods by ion-pair chromatography 346(1985)333

Sorbinil
HPLC analysis of glucose, fructose, sorbitol and low levels of the aldose reductase inhibitor sorbinil in human lens and plasma 343(1985)402

Sorbitol
Elevation of certain polyols in the cerebrospinal fluid of patients with multiple sclerosis 336(1985)351

HPLC analysis of glucose, fructose, sorbitol and low levels of the aldose reductase inhibitor sorbinil in human lens and plasma 343(1985)402

Sotalol
HPLC determination of sotalol in biological fluids 336(1985)313

Specific surface area
Adsorption effects in GLC. Solute retention in the hydrocarbon solute-polar stationary phase (polypropylene glycol 2000) system 330(1985)193

Spectinomycin
Derivatization of secondary amines with 2-naphthalenesulfonyl chloride for HPLC analysis of spectinomycin 333(1985)365

Spermidine
Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Spheron Ara
Adsorption of SH-containing amino acids, peptides and proteins on the 2-nitrophenylhydrazonopropanedinitrile derivative of Spheron Ara 328(1985)402

Sphingomyelin
Estimation of amniotic fluid phospholipids by HPLC 336(1985)139

Sphondin
Reversed-phase HPLC separation of closely related furocoumarins 346(1985)456

Spiramycin
Automated HPLC determination of spiramycin by direct injection of plasma, using column-switching for sample clean-up 344(1985)275

Spironolactone
Simultaneous determination of althiazide and spironolactone in tablets by HPLC 329(1985)422

Determination of the serum concentration of spironolactone and its metabolites by HPLC 341(1985)279

Spiroperidol
Sensitive analysis of butyrophenone neuroleptics by HPLC with UV detection at 254 nm 341(1985)465

Spray reagents
Rubeanic acid: a general spray reagent for the detection of steroids on thin-layer plates 350(1985)468

SQ 27,519
Determination of SQ 27,519, the active phosphinic acid-carboxylic acid of the prodrug SQ 28,555, in human serum by capillary GC with nitrogen-phosphorus detection after a two-step derivatization 345(1985)299

Stachynene
PC of the essential oils occurring in the genus Stachys 333(1985)288

Stachynone
PC of the essential oils occurring in the genus Stachys 333(1985)288

Stationary phase effects
Role of polar groups in fractionating isomeric polystyrene oligomers 347(1985)39

Stationary phases
see also Chiral stationary phases

Stationary phases, GC
Effects of test solutes on the characterization of stationary phases in GC using a Snyder triangle 329(1985)5

Open-tubular columns support-coated *in situ* with cross-linked SE-54 stationary phase 329(1985)123

Improved method for the preparation of immobilised pyrimidonesilicones as polar stationary phases used for the capillary column GC separation of underivatised barbiturates 329(1985)213

Coordination polymers as stationary phases and adsorbents for GC. VIII. An attempt to bind poly[chromium(III) diphenylphosphinate] to silica 330(1985)370

New liquid crystalline stationary phases for GC of positional and geometrical isomers having similar volatilities 331(1985)69

Soviet polysiloxane stationary phases for GC (Review) 334(1985)1

Chromatographic properties of oxidised OV-17 stationary phase 346(1985)25

Preparation of glass and fused-silica capillary columns coated with immobilized XE-60 346(1985)341

Comparative GC studies of bonded and physically coated PEG 20M and its derivative phases with reference to some high-boiling isomeric solutes 346(1985)359

New approach to capillary columns for GC? Condensation of hydroxyl-terminated stationary phases 347(1985)351

Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

Catalytic hydrogenation and exchange reactions on GC columns, in the presence of supported $(\eta^5\text{-}C_5H_5)NiOs_3(\mu\text{-}H)_3(CO)_9$ 349(1985)173

Effect of the mesh-size distribution of porous polymer beads on efficiency and selectivity of stationary phases 349(1985)189

Variation in the GC stationary phase properties of tetra-*n*-butylammonium salts as a function of the anion type 349(1985)235

Evaluation of tetraalkylammonium tetrafluoroborate salts as high-temperature stationary phases for packed and open-tubular column GC 349(1985)249

Selectivity of complexing GC stationary phases containing lanthanide β-diketonates 349(1985)457

Contributions to hydrocarbon retention in gas-liquid-solid chromatography with polyethylene glycol 6000 as stationary phase 350(1985)15

Stationary phases, LC

High-performance hydrophobic-interaction chromatography on ether-bonded phases. Chromatographic characteristics and gradient optimization 326(1985)45

A new, stabilized, hydrophilic silica packing for the high-performance gel chromatography of macromolecules 326(1985)63

HPLC of tRNAs on novel stationary phases 326(1985)79

High-performance liquid affinity chromatography of β-lactamase 326(1985)163

Comparison of weak and strong high-performance anion-exchange chromatography 327(1985)139

Separation of peptides by strong cation-exchange HPLC 327(1985)147

Re-packing reversed-phase HPLC columns as means of regenerating column efficiency and prolonging packing life 328(1985)347

Adsorption of SH-containing amino acids, peptides and proteins on the 2-nitrophenylhydrazonopropanedinitrile derivative of Spheron Ara 328(1985)402

LC on triacetylcellulose. Preparative separation of enantiomers on an axially compressed column 329(1985)307

Long-chain alkyl grafts and mixed alkyl-alkane layers at the surface of macroporous silicas. Their GC properties below and above the phase transition 329(1985)331

Quality criteria and structure of silica gel column packing material 329(1985)351

Insertion of various long alkyl chain molecules in brush-type grafted monolayers. Chromatographic study of the resulting materials 330(1985)227

Properties of six octadecylsilane stationary phases for the separation of dansylamides 330(1985)243

Column lifetime of a new agarose medium for high-performance gel filtration chromatography at basic pH 330(1985)360

Examination of the effect of solvent composition on bonded phase LC packings by ^{13}C Fourier transform NMR spectroscopy 331(1985)23

Influence of the stationary phase on the retention characteristics of purine and pyrimidine bases, ribonucleosides and deoxyribonucleosides in reversed-phase HPLC 331(1985)33

Effects of solutions used for storage of size-exclusion columns on subsequent chromatography of peptides and proteins 331(1985)253

Polyallylamine-coated silica gel microbore column for LC 332(1985)15

New gel permeation column for the separation of water-soluble polymers 332(1985)39

On-column preparation of chemically bonded stationary phase with maximum surface coverage and high reproducibility, and its application to packed microcapillary columns 332(1985)57

Recent progress in gel packing materials and detectors for modern LC in Japan 332(1985)227

Use of Toyopearl as a support for the immobilization of urease 332(1985)237

Retention mechanism of nucleotides, nucleosides and their bases on polyvinyl alcohol 332(1985)255

Preparative high-performance ion-exchange chromatography 333(1985)29

Resin-based support for reversed-phase chromatography of proteins 333(1985)93

High-performance affinity chromatography of trypsins on Asahipak GS-gel coupled with *p*-aminobenzamidine 333(1985)107

Adsorption chromatographic separations on buffered silica gel (Review) 334(1985)35

LC retention behavior of large, fused polycyclic aromatics. Normal bonded phases 346(1985)81

Performance characteristics of some commercially available low-capacity anion-exchange columns suitable for non-suppressed ion chromatography 346(1985)139

Three independent methods for quantitative determination of octyl covalently coupled to Sepharose CL-4B 346(1985)255

LC retention behavior of organometallic compounds and ligands with amine-, octadecylsilica- and β-cyclodextrin-bonded phase columns 347(1985)51

Polyethylene oxide: a ligand for mild hydrophobic interaction chromatography? 347(1985)291

Novel GPC packing composed of cross-linked and porous spherical particles of poly-γ-methyl-L-glutamate 347(1985)357

Selectivity of a diol phase HPLC system in trace analysis of anabolic compounds 347(1985)375

Reversed-phase HPLC of insulin and insulin derivatives. A comparative study 348(1985)347

Reversed-phase LC on rigid vinyl alcohol copolymer gels 349(1985)323

Swelling propensity (SP factor) of semi-rigid chromatographic packing materials 350(1985)145

Comparative study of affinity chromatography of components of the hyaluronate-proteoglycan complex to immobilized hyaluronate 350(1985)227

Preparation of low-capacity anion-exchange resins for ion chromatography on a methacrylic copolymer matrix 350(1985)296

HPLC separation of aromatic carboxylic acids with β-cyclodextrin-bonded stationary phases 350(1985)371

Steady-state rheoelectrolysis
Isoelectric focusing in non-amphoteric buffers. Catastrophe and non-catastrophe theories (Review) 334(1985)71

Steroidal acids
Bile acids. LXXVI. Analyses of bile acids and sterols by HPLC with a microbore column 347(1985)155

Steroidal alkaloids
Determination of the total C_{27}-steroidal alkaloid composition of *Solanum* species by high resolution GC 331(1985)285

Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

Steroidal sulphates
Clinical analysis of steroids. XXXII. New method for the determination of hapten number of the antigen prepared for the radioimmunoassay of steroidal sulphates using ion chromatography 338(1985)404

Steroid receptor proteins
Isolation of unliganded steroid receptor proteins by high-performance size-exclusion chromatography. An investigation of steroid-dependent structural alterations 327(1985)221

High-performance hydrophobic-interaction chromatography of steroid hormone receptors 327(1985)237

High-performance chromatofocusing of steroid receptor proteins in the presence and absence of steroid. Investigation of steroid-dependent alterations in surface charge heterogeneity 327(1985)247

Rapid, high-resolution procedure for assessment of estrogen receptor heterogeneity in clinical samples 327(1985)369

Steroids
see also Hormones

Chromatography of bis-quaternary amino steroids. I. Separation on silica by TLC and HPLC 328(1985)279

Rapid extraction and purification of diethylstilboestrol in bovine urine hydrolysates using reversed-phase C_{18} columns before determination by radioimmunoassay 328(1985)413

Characterization of C_{19} steroids by two-dimensional TLC 329(1985)171

Analysis of 25-hydroxycholesterol in plasma and cholesterol-containing foods by HPLC 330(1985)339

Reversed-phase HPLC with isocratic elution of 6α-methyl-17α-acetoxyprogesterone (MAP) and its impurities 330(1985)415

Catalysed derivatisation of trimethylsilyl ethers of ecdysterone. A preliminary study 333(1985)186

Reversed-phase HPLC of C_{21} metabolites of progesterone 336(1985)361

Clinical analysis of steroids. XXXI. Assay of oestradiol 17-sulphate 4-hydroxylase activity by HPLC with electrochemical detection 337(1985)114

Determination of testosterone propionate in human plasma by GC-MS 337(1985)205

Total oestriol in maternal serum or plasma as measured by LC 337(1985)379

15-Hydroxycyproterone acetate and cyproterone acetate levels in plasma and urine 338(1985)179

Quantitative determination of methyltestosterone and methyltestosterone-d_3 in serum by GC-MS 338(1985)281

Clinical analysis of steroids. XXXII. New method for the determination of hapten number of the antigen prepared for the radioimmunoassay of steroidal sulphates using ion chromatography 338(1985)404

GC-MS of epimeric 19-norandrostan-3-ol-17-ones as the trimethylsilyl ether, methyloxime-trimethylsilyl ether and trimethylsilyl-enol trimethylsilyl ether derivatives 339(1985)11

Determination of plasma testosterone by mass fragmentography using [3,4-^{13}C]testosterone as an internal standard 339(1985)233

Efficient chromatographic fractionation of steroids in human serum through regulation of liquid-liquid distribution ratios 339(1985)253

Quantitative determination of betamethasone and its major metabolite in equine urine by micro-LC-MS 341(1985)261

Determination of oestriol in pregnancy urine by normal-phase HPLC with electrochemical detection 341(1985)271

Determination of the serum concentration of spironolactone and its metabolites by HPLC 341(1985)279

3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 341(1985)426

Simultaneous determination of cholestanol and cholesterol in human serum by HPLC with fluorescence detection 341(1985)432

Simple HPLC method for the determination of medroxyprogesterone acetate in human plasma 341(1985)437

Rapid method for the measurement of methylprednisolone and its hemisuccinate in plasma and urine following "pulse therapy" by HPLC 342(1985)251

Determination of progesterone by LC-MS using a moving-belt interface and isotope dilution 343(1985)397

HPLC assay for methylprednisolone and its soluble prodrug esters in dog plasma 344(1985)41

Specific and sensitive determination of medroxyprogesterone acetate in human serum by GC-MS 345(1985)1

Studies on steroids. CCXVII. Separation and characterization of bile acid 3-glucuronides in human urine by HPLC 345(1985)241

Simultaneous determination of free and conjugated ecdysteroids by LC 346(1985)301

Chromatography of bis-quaternary amino steroids. II. Separation on chemically bonded silica phases 347(1985)284

Selectivity of a diol phase HPLC system in trace analysis of anabolic compounds 347(1985)375

Studies on steroids. CCXVI. Separation of bile acid 3-glucuronides by HPLC 348(1985)151

Rubeanic acid: a general spray reagent for the detection of steroids on thin-layer plates 350(1985)468

Sterol acetates
Determination of sterol and triterpene alcohol acetates in natural products by reversed-phase LC and GC-MS 329(1985)231

Sterols
Bile acids. LXXVI. Analyses of bile acids and sterols by HPLC with a microbore column 347(1985)155

Stilboestrol
Separation and detection of enantiomers of stilbestrol analogues by combined HPLC-thermospray MS 344(1985)378

Stilphostrol
Application of a capillary GC-selected-ion recording MS technique to the analysis of diethylstilbestrol and its phosphorylated precursors in plasma and tissues 339(1985)87

Storage effects
Effects of solutions used for storage of size-exclusion columns on subsequent chromatography of peptides and proteins 331(1985)253

***Streptomyces* R61**
Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary GC-MS 341(1985)139

Streptomycin
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Determination of streptomycin in serum by HPLC 343(1985)379

Structure-activity relationships
TLC and HPLC in the evaluation of the lipophilicity of aryloxoalkanoic and arylhydroxyalkanoic acids 347(1985)335

Structure parameters
Structure parameters of molecules and media evaluated by chromatographic partition. I. Controlled-pore glasses 350(1985)1

Structure-retention relationships
Temperature-programmed retention indices of 221 halogenated organic compounds with 1-bromoalkanes as references 328(1985)35

Comparison of relationships between the chemical structures and mobilities of hyaluronate oligosaccharides in TLC and HPLC 328(1985)73

GLC analyses. XLII. Retention behaviour of C_1-C_{12} *n*-alkyl esters of benzoic and pentafluorobenzoic acids on SE-30 and OV-351 capillary columns 328(1985)325

GLC analyses. XLIII. Retention increments for 2-chloro-, 2,2-dichloro- and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns 329(1985)43

Relationships between the chromatographic behaviour and structure of some substituted dibenzo[*b,f*]thiepins and their analogues 329(1985)113

GLC analyses. XLIV. Benzoyl and pentafluorobenzoyl esters of lower saturated branched-chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 329(1985)359

Retention indices of dimethylbenzamidines and benzylideneamines on a non-polar column 330(1985)79

HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. II. Retention behavior, quantitative structure-retention relationships and octanol-water partition coefficients 330(1985)181

Kováts retention indices of high-molecular-weight monomethyl-, cyclopentyl-, cyclohexyl- and phenylalkanes 331(1985)219

Phenol effect in PC of metal complexes 331(1985)273

GC of bicyclo[*n*.4.0]alken-2-ones 331(1985)406

Energy effects in the retention of aromatic acids in LC 332(1985)189

Correlations between retention on liquid crystalline stationary phases and chemical structure. I. Dimethylnaphthalenes 333(1985)11

Relationships between GLC behaviour and structure of arylaliphatic acids 333(1985)21

Correlation of the chromatographic retention of some phenylacetic and phenylpropionic acid derivatives with molecular structure 333(1985)309

Chromatography of monomers. IV. GLC studies of C_1-C_6 *n*-alkyl and C_3-C_6 isoalkyl acrylates and their hydrogen halide and halogen addition derivatives 348(1985)141

Physico-chemical characterization of cyclic nucleotides by reversed-phase HPLC. II. Quantitative determination of hydrophobicity 350(1985)105

Separation of some peptides and related isopeptides by HPLC: structure-retention time relationships 350(1985)279

Styrene copolymers
Correlation between chemical and physical characteristics and adsorption properties of some styrene copolymers 330(1985)55

Pre-concentration of organic pollutants. Potential interference from the use of styrene copolymer adsorbents 349(1985)181

Substrate-selective polymers
Molecular imprinting of amino acid derivatives in macroporous polymers. Demonstration of substrate- and enantio-selectivity by chromatographic resolution of racemic mixtures of amino acid derivatives 347(1985)1

Use of silane monomers for molecular imprinting and enzyme entrapment in polysiloxane-coated porous silica 347(1985)11

Succinimides
Determination of antiepileptic drugs (Review) 340(1985)199

Succinyl-lysozyme-diaminooctyl-polycytidylic acid
Neonucleoproteins. Preparation and HPLC characterization of succinyl-lysozyme-diaminooctyl-polycytidylic acid and related polycytidylic acid conjugates 326(1985)311

Sufentanil
Quantitative analysis of common anaesthetic agents (Review) 340(1985)273

Sugar acids
High-speed LC of sugar acids on anion-exchange resins 329(1985)371

Sugar alcohols
Counter-ion effects in partition chromatography 332(1985)203

Sugars
see also Carbohydrates and saccharides

Counter-ion effects in partition chromatography 332(1985)203

Radioactivity detection system with a CaF_2 (Eu) scintillator for HPLC 333(1985)83

Measurement of neutral sugars in glycoproteins as dansyl derivatives by automated HPLC 333(1985)123

Identification of inositols and their mono-O-methyl ethers by GLC 333(1985)167

Identification of 6-deoxyallitol and 6-deoxygulitol in human urine. Electron-impact mass spectra of eight isomers of 6-deoxyhexitol 336(1985)345

Elevation of certain polyols in the cerebrospinal fluid of patients with multiple sclerosis 336(1985)351

Isolation and quantitation of carbohydrates in sheep plasma by HPLC 338(1985)303

Evidence for the degradation of sugars during the resin hydrolysis of glycoproteins 339(1985)182

Quantitative method for the detection of glucose in body fluids by HPLC with fluorescence detection 342(1985)179

HPLC analysis of glucose, fructose, sorbitol and low levels of the aldose reductase inhibitor sorbinil in human lens and plasma 343(1985)402

Sugar composition of lipopolysaccharide from *Haemophilus paraphrophilus* 345(1985)119

Modifications in the alditol acetate method for analysis of muramic acid and other neutral and amino sugars by capillary GC-MS with selected ion monitoring 347(1985)109

Capillary GC of partially methylated alditol acetates on a SP-2100 wall-coated open-tubular column 350(1985)304

Sulbactam
HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 341(1985)115

Sulphadiazine
Rapid determination of five sulphonamides in swine tissue by HPLC 333(1985)256

Sulphadoxine
Rapid determination of five sulphonamides in swine tissue by HPLC 333(1985)256

Sulphamethazine
Determination of oxytetracycline, sulphamethazine and sulphamethoxypyridazine in feed premixes 330(1985)425

Rapid determination of five sulphonamides in swine tissue by HPLC 333(1985)256

Sulphamethizole
HPLC determination of phenazopyridine hydrochloride, tetracycline hydrochloride and sulphamethizole in combination 333(1985)249

Sulphamethoxypyridazine
Determination of oxytetracycline, sulphamethazine and sulphamethoxypyridazine in feed premixes 330(1985)425

Sulphanilamide
Rapid determination of five sulphonamides in swine tissue by HPLC 333(1985)256

Sulphaquinoxaline
Rapid determination of five sulphonamides in swine tissue by HPLC 333(1985)256

Sulphate
Determination of inorganic sulphate in plasma by reversed-phase chromatography using UV detection and its application to plasma samples of patients receiving different types of haemodialysis 337(1985)259

Determination of inorganic sulfate in human saliva and sweat by controlled-flow anion chromatography. Normal values in adult humans 337(1985)267

Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Sulphate esters
Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

Sulphides
Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Ligand-exchange GC of dialkyl sulphides 350(1985)63

Ligand-exchange GC separation of aromatic sulphides 350(1985)75

Sulphidopeptide leukotrienes
Quantitation of sulfidopeptide leukotrienes by reversed-phase HPLC 343(1985)213

Sulphite
Rapid determination in water of chloride, sulphate, sulphite, selenite, selenate and arsenate among other inorganic and organic solutes by ion chromatography with UV detection below 195 nm 347(1985)419

Sulphonamides
Rapid determination of five sulphonamides in swine tissue by HPLC 333(1985)256

Sulphosalicylic acid
Effect of complexing agents on the chromatographic separation of polyvalent cations 347(1985)147

Sulphur
Elemental sulphur analysis using HPLC on 10-μm rigid polymer particles 348(1985)317

Sulphur gases
GC and MS identification of sulphur gases in kraft digestion plants 346(1985)405

Sulphydryls
Rapid and simple method for quantitative determination of non-protein sulphydryls in mouse liver by reversed-phase HPLC 338(1985)209

Sulpiride
Quantitative analysis of sulpiride and sultopride by HPLC for pharmacokinetic studies (in French) 341(1985)391

Sulpyrin
Analysis of antipyretics by semimicro LC 332(1985)269

Sultopride
Quantitative analysis of sulpiride and sultopride by HPLC for pharmacokinetic studies (in French) 341(1985)391

Supercritical fluid chromatography
Supercritical fluid chromatography-IR spectroscopy of oligomers: use of buffer-memory technique 332(1985)47

Separation of oligomers with UV-absorbing side groups by supercritical fluid chromatography using eluent gradients 346(1985)69

Effects of density and temperature on efficiency in capillary supercritical fluid chromatography 349(1985)305

Gradient programming and combined gradient-pressure programming in supercritical fluid chromatography 349(1985)317

Group separation of oil residues by supercritical fluid chromatography 349(1985)439

Supercritical fluid extraction-supercritical fluid chromatography
New double-stage separation analysis method. Directly coupled laboratory-scale supercritical fluid extraction-supercritical fluid chromatography, monitored with a multiwavelength-UV detector 332(1985)107

Superose
Agarose-based media for high-resolution gel filtration of biopolymers 326(1985)33

Column lifetime of a new agarose medium for high-performance gel filtration chromatography at basic pH 330(1985)360

Superoxide dismutase
Application of a new ion exchanger TSK-GEL DEAE-5PW, to the purification of Cu,Zn-superoxide dismutase of bovine erythrocytes 327(1985)301

Preparative high-performance ion-exchange chromatography 333(1985)29

Suramin
Determination of suramin in plasma by HPLC 343(1985)224

Surfactants
Increments of some polarity parameters for polyoxyethylene glycol dialkyl ethers and for some of their sulphur analogues 330(1985)61

Sweeteners
HPLC analysis of Aspartame 346(1985)417

Ion-pair HPLC analysis of aspartame and related products 349(1985)83

Swelling propensity
Swelling propensity (SP factor) of semi-rigid chromatographic packing materials 350(1985)145

Sweroside
Analytical and preparative reversed-phase LC of secoiridoid glycosides 346(1985)396

Swertiamarin
Analytical and preparative reversed-phase LC of secoiridoid glycosides 346(1985)396

Switching valve
Automated pre-column system for trace enrichment and clean-up of plasma samples in narrow-bore LC 330(1985)113

Synephrine
Detection of phenolalkylamines by capillary column GC negative chemical ionization MS 330(1985)315

System peaks
Occurrence and origin of system peaks in non-suppressed ion chromatography of inorganic anions with indirect UV absorption detection 346(1985)125

T

Tabun
Capillary column GC-MS of tabun 347(1985)257

Tamoxifen citrate
Quantitative GC determination of tamoxifen citrate in pharmaceuticals 331(1985)432

Taurine
Electron-capture GC of taurine as its N-pentafluorobenzoyl di-*n*-butylamide derivative 339(1985)370

Determination of cysteinesulfinate, hypotaurine and taurine in physiological samples by reversed-phase HPLC 343(1985)303

Taurocyamine
Quantitative overestimation of free taurocyamine using short liquid cation-exchange chromatography columns? 345(1985)215

Tear-gases
GC-MS identification of tear-gases in dilute solutions using large injection volumes 330(1985)379

Tecoflex
Chromatographic analysis of thermoplastic, medical-grade, polyether-based polyurethane elastomers 349(1985)347

Temperature effects
Denaturation and the effects of temperature on hydrophobic-interaction and reversed-phase HPLC of proteins. Bio-Gel TSK-Phenyl-5-PW column 327(1985)77

Effects of density and temperature on efficiency in capillary supercritical fluid chromatography 349(1985)305

Effect of column temperature on HPLC behaviour of inorganic polyphosphates. II. Gradient ion-exchange chromatography 350(1985)119

Temperature gradient
TLC separations using a temperature gradient 333(1985)69

Tenax GC
Comparison between theoretical and experimental sampling efficiencies on Tenax GC 331(1985)207

Teniposide
HPLC determination of etoposide in plasma using electrochemical detection 336(1985)434

Drug level monitoring: cytostatics (Review) 340(1985)31

Tensides
Interaction of some non-ionic tensides with dioleoyl phosphatidyl choline, studied by charge-transfer chromatography 349(1985)295

Terbutaline
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Terbutryn
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Terbutylazine
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Terephthalamidine
Analysis of candidate anticancer drugs by thermospray HPLC-MS and by high-resolution MS 347(1985)137

Terpene aldehydes
HPLC analysis of terpene aldehydes in cotton 329(1985)91

Terpenols
The aroma of grapes. I. Extraction and determination of free and glycosidically bound fractions of some grape aroma components 331(1985)83

Territrems
TLC of mycotoxins (Review) 334(1985)211

Tertatolol
Quantitative determination of tertatolol in biological fluids by GC-MS 339(1985)97

Testosterone
Determination of testosterone propionate in human plasma by GC-MS 337(1985)205

Determination of plasma testosterone by mass fragmentography using [3,4-^{13}C]testosterone as an internal standard 339(1985)233

Tetraalkylammonium tetrafluoroborate salts
Evaluation of tetraalkylammonium tetrafluoroborate salts as high-temperature stationary phases for packed and open-tubular column GC 349(1985)249

Tetrabutylammonium phosphate
Removal of counter-ion reagent after semi-preparative reversed-phase ion-pair HPLC 331(1985)445

Tetra-*n*-butylammonium salts
Variation in the GC stationary phase properties of tetra-*n*-butylammonium salts as a function of the anion type 349(1985)235

Tetrachloroethylene
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233

2,3,5,6-Tetrachloropyridine
Analysis of some chlorinated pyridine isomers in mixtures by reversed-phase HPLC 348(1985)309

Tetracyclines
HPLC determination of phenazopyridine hydrochloride, tetracycline hydrochloride and sulphamethizole in combination 333(1985)249

Antibiotic monitoring in body fluids (Review) 340(1985)361

Tetraethylenepentamine
Ion-pair HPLC determination of dansylated aliphatic polyamines with fluorescence and UV detection 333(1985)220

1,2,3,4-Tetrahydro-β-carbolines
Derivatization in aqueous solution, isolation and separation of tetrahydro-β-carbolines and their precursors by LC 341(1985)287

Tetrahydropterins
Estimation of tetrahydro, dihydro and fully oxidised pterins by HPLC using sequential electrochemical and fluorometric detection 343(1985)35

Tetrahydrothiophene
On-line analysis of sulphur compounds in natural gas with flame photometric detection 349(1985)465

2,3,4,6-Tetra-O-acetyl-β-D-glucopyranosyl isothiocyanate
Determination of the enantiomeric composition of chiral epoxides using chiral derivatization and LC 331(1985)349

Theobromine
Analysis of antipyretics by semimicro LC 332(1985)269

HPLC determination of dimethylxanthine metabolites of caffeine in human plasma 338(1985)369

Theophylline
Analysis of antipyretics by semimicro LC 332(1985)269

Automated theophylline assay using GC and a mass-selective detector 336(1985)285

HPLC determination of dimethylxanthine metabolites of caffeine in human plasma 338(1985)369

Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

Improved HPLC method for the simultaneous determination of caffeine and its N-demethylated metabolites in plasma using solid-phase extraction 342(1985)105

Two HPLC methods for quantitation of theophylline in plasma 343(1985)359

Thermionic detector
Nitrogen-sensitive thermionic detection in microcolumn LC 333(1985)291

Thiacetazone
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Thiamazole
Determination of thiamazole in serum by HPLC with electrochemical detection 339(1985)149

Thiamine
Determination of thiamine and its phosphate esters in human and rat blood by HPLC with post-column derivatization 332(1985)181

Evaluation of internal standards and extraction solvents in the GC determination of thiamine 347(1985)89

Thiamphenicol
Antibiotic monitoring in body fluids (Review) 340(1985)361

Thiazides
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Influence of the chain length of chemically bonded phases on the behaviour of several thiazide, potassium-sparing and loop diuretics in HPLC 349(1985)301

Thin-layer stick chromatography
Application of thin-layer stick chromatographic identification test methods to drugs contained in preparations in the Pharmacopoeia of Japan 333(1985)115

2,2'-Thiodiethanol
Analysis of 2,2'-thiodiethanol in aqueous matrices by LC with electrochemical detection 330(1985)388

Thiodiglycol
Analysis of 2,2'-thiodiethanol in aqueous matrices by LC with electrochemical detection 330(1985)388

Thioglycolic acid
Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193

6-Thioguanine
Sensitive HPLC determination of 6-mercaptopurine, 6-thioguanine, 6-mercaptopurine riboside and 6-thioguanosine in biological fluids 336(1985)422

HPLC assay of 6-methylthioguanine, a major metabolite of 6-thioguanine, in urine 337(1985)321

Drug level monitoring: cytostatics (Review) 340(1985)31

6-Thioguanosine
Sensitive HPLC determination of 6-mercaptopurine, 6-thioguanine, 6-mercaptopurine riboside and 6-thioguanosine in biological fluids 336(1985)422

Thiols
Adsorption of SH-containing amino acids, peptides and proteins on the 2-nitrophenylhydrazonopropanedinitrile derivative of Spheron Ara 328(1985)402

Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193

Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Thiomercurimetric detector
Electrophoretic separation and thiomercurimetric monitoring of ionogenic thiols 331(1985)193

7α-Thiomethylspirolactone
Determination of the serum concentration of spironolactone and its metabolites by HPLC 341(1985)279

Thiopental
Determination of thiopental and pentobarbital in plasma using HPLC 343(1985)437

Thiophene
Sorption properties of some modified molecular sieves 13X towards thiophene and benzene 329(1985)128

GC detection of reduced sulfur compounds using ozone chemiluminescence 347(1985)121

Thiophenes
Preconcentration of volatile organosulphur compounds from the atmosphere on selected porous polymers 333(1985)327

Thioridazine 5-sulphoxide
Determination of diastereoisomeric pairs of thioridazine 5-sulfoxide by HPLC 336(1985)452

Thiourea
Thin-layer polarographic detector for the HPLC detection of thiourea derivatives 328(1985)127

Threonine
Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

Thromboxanes
Simultaneous determination of prostanoids in plasma by GC-negative-ion chemical-ionization MS 338(1985)273

Qualitative and quantitative measurement of hydroxy fatty acids, thromboxanes and prostaglandins using stable isotope dilutions and detection by GC-MS 344(1985)11

Tiamenidine
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Ticarcillin
Clavulanate-potentiated ticarcillin: HPLC assays for clavulanic acid and ticarcillin isomers in serum and urine 337(1985)301

Timolol
Timolol determination in plasma and urine by HPLC with UV detection 338(1985)249

Tiodazosin
Determination of tiodazosin concentrations in human plasma with a fluorescence HPLC method 342(1985)221

Tobramycin
Quantitative determination and separation of analogues of aminoglycoside antibiotics by HPLC 348(1985)229

Tocainide
LC determination of mexiletine and tocainide in human plasma with fluorescence detection after reaction with a modified *o*-phthalaldehyde reagent 344(1985)221

α-Tocopherol
Simultaneous determination of serum vitamin A and E by LC with fluorescence detection 341(1985)198

Determination of α-tocopherol in rat liver by HPLC utilizing UV detection 344(1985)137

Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 345(1985)275

Separation of three dimers of α-tocopherol by HPLC 347(1985)434

Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285

Tolazoline
Quantitative determination of tolazoline in serum and urine 338(1985)123

Tolbutamide
Determination of glibenclamide, chlorpropamide and tolbutamide in plasma by HPLC with UV detection 339(1985)203

Toluene diisocyanate
Capillary GC method for the determination of complex mixture of isocyanates and amines 346(1985)191

Toxaphene
Separation of organochlorine residues by adsorption chromatography prior to capillary GC 328(1985)372

Toxins
Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Rapid purification of host-specific pathotoxins from *Alternaria alternata* f. sp. *lycopersici* by solid-phase adsorption on octadecylsilane 350(1985)255

Toyopearl
Use of Toyopearl as a support for the immobilization of urease 332(1985)237

Trace analysis
Theoretical and practical limitations of capillary isotachophoresis in trace analysis 328(1985)353

Studies on sample preconcentration in ion chromatography. II. An automated, single pump preconcentration system for non-suppressed ion chromatography with conductivity detection 330(1985)95

Automated pre-column system for trace enrichment and clean-up of plasma samples in narrow-bore LC 330(1985)113

Automated monitoring of organic trace components in water. I. Continuous flow extraction together with on-line capillary GC 330(1985)263

Trace organics
Optimization of the open stripping system for the analysis of trace organics in water 348(1985)67

Tragacanth gum
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Tralomethrin
Coupling a high-performance liquid chromatograph with a liquid scientillation

detector: example in the field of pyrethroids 349(1985)105

Tramadol
Quantitative determination of tramadol in human serum by GC-MS 341(1985)65

Tranexamic acid
Determination of plasma tranexamic acid using cation-exchange HPLC with fluorescence detection 343(1985)109

Tranilast
HPLC determination of tranilast in plasma 341(1985)228

Simultaneous determination of tranilast and metabolites in plasma and urine using HPLC 345(1985)345

Tranylcypromine
Determination of tranylcypromine in plasma using GC-chemical-ionization MS 344(1985)356

Trazodone
GC-MS method for trazodone and a deuterated analogue in plasma 339(1985)303

Determination of trazodone in human plasma by LC with fluorescence detection 342(1985)442

Measurement of trazodone in biological fluids by HPLC 345(1985)187

Tremorgens
TLC of mycotoxins (Review) 334(1985)211

Trenimon
Analysis of candidate anticancer drugs by thermospray HPLC-MS and by high-resolution MS 347(1985)137

Triacetylcellulose
LC on triacetylcellulose. Preparative separation of enantiomers on an axially compressed column 329(1985)307

Triacylglycerol
Rapid separation of plasma lipoproteins by GPC on agarose gel Superose 6B 341(1985)154

Triadimenol
Separation of enantiomers of fungicides and some analogues by capillary GC using Chirasil Val 329(1985)181

Trialkylammonium phosphates
Reversed-phase HPLC of insulin and insulin derivatives. A comparative study 348(1985)347

Triamcinolone
Drug level monitoring of antiasthmatic drugs (Review) 340(1985)243

***s*-Triazines**
Recovery of *s*-triazines from water and their analysis by GC with photoionization detection 329(1985)186

Triazolam
Capillary GLC determination of the benzodiazepine triazolam in plasma using a retention gap 339(1985)192

1,2,4-Triazole
HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 341(1985)115

Trichloroacetic acid
Simple determination of trichloroacetic acid in urine using head space GC: a suitable method for monitoring exposure to trichloroethylene 337(1985)126

1,1,1-Trichloroethane
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233

Trichloroethylene
Determination of volatile chlorinated hydrocarbons by reaction GC (in German) 328(1985)233

Simple determination of trichloroacetic acid in urine using head space GC: a suitable method for monitoring exposure to trichloroethylene 337(1985)126

2,2,2-Trichloroethyl esters
GLC analyses. XLIII. Retention increments for 2-chloro-, 2,2-dichloro- and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns 329(1985)43

Trichothecenes
Rapid and sensitive determination of zearalenone in cereals by HPLC with fluorescence detection 328(1985)271

TLC of mycotoxins (Review) 334(1985)211

Analysis of some trichothecene mycotoxins by LC 347(1985)429

Tricyclic nucleoside
Reversed-phase HPLC determination of tricyclic nucleoside and tricyclic nucleoside 5'-phosphate in biological specimens 337(1985)55

Triethylenephosphoramide
GC analysis of triethylenethiophosphoramide and triethylenephosphoramide in biological specimens 338(1985)335

GLC analysis of N,N',N"-triethylene thiophosphoramide and and N,N',N"-triethylene phosphoramide in biological samples 343(1985)196

Triethylenetetramine
Ion-pair HPLC determination of dansylated aliphatic polyamines with fluorescence and UV detection 333(1985)220

Triethylenethiophosphoramide
GC analysis of triethylenethiophosphoramide and triethylenephosphoramide in biological specimens 338(1985)335

GLC analysis of N,N',N"-triethylene thiophosphoramide and and N,N',N"-triethylene phosphoramide in biological samples 343(1985)196

GC assay of triethylenethiophosphoramide in serum and urine 345(1985)173

Trifluoroacetic acid
Spectral properties of trifluoroacetic acid-acetonitrile gradient systems for separation of picomole quantities of peptides by reversed-phase HPLC 347(1985)83

Trifluoroacetyl trisaccharides
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Trifluorothymidine
Determination of ophthalmic therapeutic trifluorothymidine and its degradation product by reversed-phase HPLC 348(1985)299

Trifluperidol
Sensitive analysis of butyrophenone neuroleptics by HPLC with UV detection at 254 nm 341(1985)465

Triglycerides
HPLC technique for the separation of lipid classes 329(1985)273

Assay of lipids in dog myocardium using capillary GC and derivatization with boron trifluoride and methanol 339(1985)25

HPLC of triglycerides of Flacourtiaceae seed oils containing cyclopentenyl fatty acids (chaulmoogric oils) 348(1985)441

Quantitative capillary GLC of triglycerides on a fused-silica column with a chemically bonded stationary phase 350(1985)87

Trihydroxy pentacyclic triterpene alcohols
TLC and GLC separation of trihydroxy pentacyclic triterpene alcohols from *Calendula officinalis* flowers 330(1985)177

Trilostane
HPLC assay for trilostane and its major metabolite, 17-ketotrilostane, in human plasma 339(1985)440

Trimecaine
Determination of trimecaine metabolites in blood plasma by capillary isotachophoresis 342(1985)167

Trimethylamine
GC-MS analysis of volatile amines produced by several strains of *Clostridium* 337(1985)213

2,2,4-Trimethyl-1,2-dihydroquinoline
Analysis technique for determining the levels of organic additives in an ethylene-propylene diene monomer rubber for use in underwater acoustic applications 350(1985)187

Trimethyllysine
HPLC determination of plasma free trimethyllysine in humans 344(1985)319

Trimethylolpropane trimethacrylate
Analysis technique for determining the levels of organic additives in an ethylene-propylene diene monomer rubber for use in underwater acoustic applications 350(1985)187

Trimoprostil
Determination of acetyltrimoprostil and its metabolite trimoprostil in human or dog plasma by GC-negative-ion chemical ionization MS 339(1985)313

2,4,6-Trinitrotoluene
Metabolic studies of explosives. II. HPLC-MS of metabolites of 2,4,6-trinitrotoluene 339(1985)127

Tripamide
Improved assay procedure for tripamide in blood 339(1985)210

Tripeptides
Separation of free radicals by HPLC with electron spin resonance detection 332(1985)71

Trisaccharides
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Tris(2,2,7-trimethyloctane-3,5-dionato)chromium(III)
GC and LC separation of geometrical isomers of tris(2,2,7-trimethyloctane-3,5-dionato)chromium(III) and cobalt(III) chelates and other Cr-β-diketonates 349(1985)469

Tris(2,2,7-trimethyloctane-3,5-dionato)cobalt(III)
GC and LC separation of geometrical isomers of tris(2,2,7-trimethyloctane-3,5-dionato)chromium(III) and cobalt(III) chelates and other Cr-β-diketonates 349(1985)469

Triterpene alcohol acetates
Determination of sterol and triterpene alcohol acetates in natural products by reversed-phase LC and GC-MS 329(1985)231

Triterpene alcohols
Preparation and GLC of trimethylsilyl derivatives of dammarane-type triterpene triols and tetrols 329(1985)189

TLC and GLC separation of trihydroxy pentacyclic triterpene alcohols from *Calendula officinalis* flowers 330(1985)177

Triterpenic glycosides
Separation of natural polar substances by reversed-phase HPLC, centrifugal TLC and droplet counter-current chromatography 349(1985)69

Tropic acid
Study of the separation of enantiomers of some aromatic carboxylic acids by HPLC on a β-cyclodextrin-bonded stationary phase 333(1985)59

Troponin
Ion-exchange HPLC purification of bovine cardiac and rabbit skeletal muscle troponin subunits 343(1985)315

Troponin I
General method for the separation of cyanogen bromide digests of proteins by HPLC. Rabbit skeletal troponin I 326(1985)349

Separation of peptides by strong cation-exchange HPLC 327(1985)147

Trospium
Fluorimetric determination of the quaternary compound trospium and its metabolite in biological material after derivatization with benoxaprofen chloride 345(1985)99

Troxerutin
Separation and determination of the various constituents of troxerutin by HPLC. Application to the pharmaceutical control (in French) 346(1985)382

Trypsin
High-performance affinity chromatography of trypsins on Asahipak GS-gel coupled with *p*-aminobenzamidine 333(1985)107

Tryptathionine
Investigation of the synthesis of tryptathionine using HPLC 348(1985)421

Tryptophan
Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

Direct chiral resolution in an aqueous two-phase system using the counter-current distribution principle 333(1985)211

Purification of synthetic lipid associating peptides and the monitoring of the deformylation of N^{in}-formyltryptophan by reversed-phase HPLC 347(1985)237

Tryptophyl-leucine
Large-scale preparative countercurrent chromatography for separation of polar compounds 348(1985)131

Tryptophyllin
Reversed-phase HPLC characterization of a novel proline-rich tryptophyllin 349(1985)117

TSKgel GMPW
New gel permeation column for the separation of water-soluble polymers 332(1985)39

TSKgel Phenyl-5PW
Resin-based support for reversed-phase chromatography of proteins 333(1985)93

Preparative high-performance hydrophobic interaction chromatography of proteins on TSKgel Phenyl-5PW 333(1985)202

Tunaxanthin
Direct diastereomeric resolution of carotenoids. II. All ten stereoisomers of tunaxanthin (ε,ε-carotene-3,3'-diol) 328(1985)387

Tyramine
Highly sensitive HPLC technique for the simultaneous measurement of histamine, 1-methylhistamine and other biogenic amines 343(1985)407

Tyrosine
Analysis of sulfate and phosphate esters of amino acids by ion-exchange chromatography on polymeric DEAE 327(1985)9

Separation of some aromatic amino acids by reversed-phase HPLC using α- or β-cyclodextrin as mobile phase component 329(1985)206

T-2 toxin
GC assay with pharmacokinetic applications for monitoring T-2 and HT-2 toxins in plasma 343(1985)67

U

Ubichromenol-9
HPLC of coenzyme Q-related compounds and its application to biological materials 342(1985)35

Ultrafiltration cones
Use and regeneration of Amicon ultrafiltration cones for deproteination of microsomal solutions before chromatographic analysis 342(1985)465

Ultraviolet detector
High-performance UV absorption detector for LC. I. Preliminary experiments 329(1985)25

Uracil
HPLC of the photoproducts of nucleic acid components. II. Photohydrates of pyrimidines 329(1985)33

Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Identification of *Escherichia coli* by detection of hydroquinone and uracil in the urine system 345(1985)11

Uraemic middle molecules
Removal of counter-ion reagent after semi-preparative reversed-phase ion-pair HPLC 331(1985)445

Urea
Determination of biuret and urea by HPLC 331(1985)161

Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Urease
Use of Toyopearl as a support for the immobilization of urease 332(1985)237

Uremic toxins
Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Uric acid
HPLC determination of uric acid in soil 331(1985)359

HPLC isolation of uric acid from soil for isotopic determination 331(1985)371

Simultaneous determination of serum cations, anions and uremic toxins by ion chromatography using an immobilized enzyme 344(1985)145

Chromatographic evaluation of perfusion on charcoal in uraemia 345(1985)251

Uridine
Reversed-phase LC investigation of nucleosides and bases in mucosa and modified nucleosides in urines from patients with gastrointestinal cancer 332(1985)127

Reversed-phase HPLC investigation of mucosal nucleosides and bases and urinary modified nucleosides of gastrointestinal cancer patients 343(1985)21

Urinary proteins
Fast protein LC scale-up procedures for the preparation of low-molecular-weight proteins from urine 327(1985)269

Urocanic acid
HPLC determination of urocanic acid in stripped human stratum corneum 342(1985)247

Uroporphyrin
Ion-pair HPLC separation of porphyrin isomers 349(1985)283

Ursodeoxycholic acid
Determination of bile acids in rat bile by HPLC 338(1985)195

Rapid and accurate reversed-phase HPLC determination of conjugated bile acids in human bile for routine clinical applications. Therapeutic control during gallstone dissolution therapy 339(1985)263

V

Valnoctamide
Rapid GC assay for monitoring valnoctamide in plasma 337(1985)408

Valproic acid
Rapid and selective derivatization method for the nitrogen-selective detection of carboxylic acids in biological fluids prior to GC analysis 336(1985)241

Concurrent determination of valproic acid with other antiepileptic drugs by HPLC 338(1985)131

Determination of antiepileptic drugs (Review) 340(1985)199

Improved direct injection method and extractive methylation method for determination of valproic acid in serum by GC 342(1985)89

Valyl-tyrosine
Large-scale preparative countercurrent chromatography for separation of polar compounds 348(1985)131

Vanadium chelates
HPLC of copper, cobalt, nickel, chromium and vanadium chelates with 2-(5-bromo-2-pyridylazo)-5-(N-propyl-N-sulphopropylamino)phenol 333(1985)133

Vancomycin
Antibiotic monitoring in body fluids (Review) 340(1985)361

Van der Waals volumes
Energy effects in the retention of aromatic acids in LC 332(1985)189

Vanillic acid
GC profiling of phenolic acids in urine of patients with cirrhosis of the liver 338(1985)25

Vanillylmandelic acid
HPLC assay of free norepinephrine, epinephrine, dopamine, vanillylmandelic acid and homovanillic acid 343(1985)9

Rapid and reliable estimation of urinary free catecholamines in patients with pheochromocytoma: comparison with plasma catecholamines and vanillylmandelic acid excretion 344(1985)313

Vanilmandelic acid
Measurement of urinary vanilmandelic acid and homovanillic acid by HPLC with electrochemical detection following extraction by ion-exchange and ion-moderated partition 336(1985)173

Vapor pressure
Vapor pressure estimates of individual polychlorinated biphenyls and commercial fluids using GC retention data 330(1985)203

Vapour-phase protein sequencer
Automated on-line identification of phenylthiohydantoin-amino acids from a vapor phase protein sequencer 350(1985)217

Vasoactive intenstinal polypeptide
Measurement and chromatographic characterization of vasoactive intestinal peptide from guinea-pig enteric nerves 336(1985)41

Vasoactive intestinal polypeptide
Rapid, small-scale preparation of gastrointestinal hormones by HPLC on a C_{18} column 326(1985)121

Vasodilating agents
Drug level monitoring: cardiovascular drugs (Review) 340(1985)73

Vasopressin
Quantitative affinity HPLC of neuroendocrine polypeptides using porous and non-porous glass derivatives 327(1985)193

HPLC and studies of neurophysin-neurohypophysial hormone pathways 336(1985)63

Veralipride
Quantitative analysis of veralipride in plasma and urine by GC-MS and GC with flame-ionization detection 338(1985)79

Verapamil
Simultaneous quantitation of verapamil, norverapamil, and N-dealkylated metabolites in human plasma following oral administration 337(1985)160

HPLC measurement of verapamil in plasma using a diol column 337(1985)412

Rapid GLC method for plasma verapamil level determination 342(1985)406

Veratramine
Isolation of teratogenic alkaloids by reversed-phase HPLC 336(1985)211

Verruculogen
TLC of mycotoxins (Review) 334(1985)211

Vesicants
GC retention indices of chemical warfare agents and simulants 331(1985)47

Vigabatrin
Quantitative analysis of vigabatrin in plasma and urine by reversed-phase HPLC 341(1985)232

Viloxazine
One-step extraction procedure for GC determination of viloxazine as its acetyl derivative in human plasma 337(1985)142

Vinblastine
HPLC determination of vinca-alkaloids in plasma and urine 345(1985)309

Vinca alkaloids
Drug level monitoring: cytostatics (Review) 340(1985)31

Vincamine
Capillary GC determination of vincamine in plasma 344(1985)367

Vincristine
HPLC determination of vinca-alkaloids in plasma and urine 345(1985)309

Vindesine
HPLC determination of vinca-alkaloids in plasma and urine 345(1985)309

Vinyl alcohol copolymer gels
Reversed-phase LC on rigid vinyl alcohol copolymer gels 349(1985)323

γ-Vinyl-4-aminobutyric acid
Determination of amino acids by separation of their ion pairs with dodecyl sulphate 341(1985)11

Vinyl arene compounds
Separation of oligomers with UV-absorbing side groups by supercritical fluid chromatography using eluent gradients 346(1985)69

N-Vinylcarbazole oligomers
Separation of oligomers with UV-absorbing side groups by supercritical fluid chromatography using eluent gradients 346(1985)69

Vinyl ether lysophospholipids
Reversed-phase HPLC separation of molecular species of alkyl ether, vinyl ether, and monoacyl lyosphospholipids 338(1985)61

Viomellein
TLC of mycotoxins (Review) 334(1985)211

Viomycin
Chromatographic analysis of antituberculosis drugs in biological samples (Review) 340(1985)321

Virus proteins
Characterization of a solvent system for separation of water-insoluble poliovirus proteins by reversed-phase HPLC 326(1985)91

Separation of flavivirus membrane and capsid proteins by multistep HPLC optimized by immunological monitoring 326(1985)113

Combined size-exclusion and reversed-phase HPLC of a detergent extract of Sendai virus 326(1985)173

Isolation of the subunits of the coronavirus envelope glycoprotein E2 by hydroxyapatite HPLC 326(1985)191

Detection of Sendai virus protein by reversed-phase HPLC combined with immuno-chromatography 327(1985)377

Vitamin A
Simultaneous determination of serum vitamin A and E by LC with fluorescence detection 341(1985)198

Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 345(1985)275

Rapid analysis of the major classes of retinoids by step gradient reversed-phase HPLC using retinal (O-ethyl) oxime derivatives 348(1985)241

Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285

Vitamin B_6
Quantities of B_6 vitamers in human milk by HPLC. Influence of maternal vitamin B_6 status 337(1985)249

Stability of pyridoxal-5-phosphate semicarbazone: applications in plasma vitamin B_6 analysis and population surveys of Vitamin B_6 nutritional status 342(1985)277

Vitamin D
Solid phase extraction system for vitamin D and its major metabolites in human plasma 343(1985)43

Vitamin D_3
Specific mass fragmentographic assay for 25,26-dihydroxyvitamin D in human plasma using a deuterated internal standard 338(1985)289

Vitamin E
Simultaneous determination of serum vitamin A and E by LC with fluorescence detection 341(1985)198

Determination of α-tocopherol in rat liver by HPLC utilizing UV detection 344(1985)137

Determination of vitamins A and E in serum and plasma using a simplified clarification method and HPLC 345(1985)275

Separation of three dimers of α-tocopherol by HPLC 347(1985)434

Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285

Vitamin K_1
Quantitative analysis of phylloquinone (Vitamin K_1) in soy bean oils by HPLC 329(1985)257

Vitamins
Application of thin-layer stick chromatographic identification test methods to drugs contained in preparations in the Pharmacopoeia of Japan 333(1985)115

Analysis of the fat-soluble vitamins using narrow-bore HPLC with multichannel UV-VIS detection 350(1985)285

Vitexin
Medicinal African plants. XIX. HPLC determination of vitexin in a crude extract of *Combretum micranthum* G. Don (in French) 346(1985)428

Void volume
Heterogeneities of blue dextran and differences among samples 350(1985)292

Volatile oils
see also Essential oils

Volatiles
Trace analysis of organic pollutants in air by polymer adsorption and dual flame ionization detection-electron-capture detection capillary GC 328(1985)367

Automated monitoring of organic trace components in water. I. Continuous flow extraction together with on-line capillary GC 330(1985)263

Simple method of concentrating volatiles in water for GC analysis by vacuum distillation 346(1985)408

Pre-concentration of organic pollutants. Potential interference from the use of styrene copolymer adsorbents 349(1985)181

Voltammetric detector
LC with voltammetric detection for quantitation of phenolic acids 348(1985)397

W

Washing
Elution and washing in fixed bed affinity chromatography 330(1985)356

Water
Automated monitoring of organic trace components in water. I. Continuous flow extraction together with on-line capillary GC 330(1985)263

Water adsorption
Adsorption from multi-component liquid mixtures on to heterogeneous solid surfaces: the role of adsorption of water on silica in liquid-solid adsorption chromatography 348(1985)1

Water pollutants
Automated monitoring of organic trace components in water. I. Continuous flow

extraction together with on-line capillary GC 330(1985)263

Water pollution
Stripping analysis and chromatographic sniffing for the source identification of odorous compounds in drinking water 328(1985)219

Weak acids
Effects of various strengths of bases on the electrophoretic behavior of weak acids in a non-aqueous solvent 330(1985)19

Weight correction factor
Quantitative capillary GLC of triglycerides on a fused-silica column with a chemically bonded stationary phase 350(1985)87

Wheat proteins
Detection of quality differences among wheats by HPLC 327(1985)333

Window diagram optimization
Solubility parameter and window diagram optimization methods for the reversed-phase chromatographic optimization of complex hydroxyl aromatic mixtures 328(1985)81

Wogonin
Simultaneous determination of baicalein, wogonin, oroxylin-A and their glucuronides in Scutellariae Radix by ion-pair HPLC 328(1985)289

X

Xanthan gum
GC determination of polysaccharide gums in foods after hydrolysis and derivatization 350(1985)237

Xanthines
Improved micro-method for the HPLC determination of caffeine and paraxanthine in biological fluids 336(1985)293

HPLC determination of dimethylxanthine metabolites of caffeine in human plasma 338(1985)369

Xanthomegnin
TLC of mycotoxins (Review) 334(1985)211

Xanthosine 5'-monophosphate
High-performance liquid chromatographic detection of XMP as a basis for an improved IMP dehydrogenase assay 343(1985)160

Xylene
New liquid crystalline stationary phases for GC of positional and geometrical isomers having similar volatilities 331(1985)69

Xylenol
Use of cyclodextrins in chromatography for selective separations, pre-concentration and preparation of defined mixtures 349(1985)167

Xylotriose
Capillary column GC of trifluoroacetyl trisaccharides 328(1985)253

Y

Yohimbine
Sub-nanogram analysis of yohimbine and related compounds by HPLC 342(1985)333

Yohimbinic acid
Sub-nanogram analysis of yohimbine and related compounds by HPLC 342(1985)333

Yutac
HPLC method for quantitative determination of Yutac, a new antiarrhythmic agent, in dog plasma 338(1985)428

Z

Zearalenone
Rapid and sensitive determination of zearalenone in cereals by HPLC with fluorescence detection 328(1985)271

TLC of mycotoxins (Review) 334(1985)211

Zeolites
On the possibility of increasing the effectiveness of chromatographic columns filled with zeolites 331(1985)402

Zirconium salt
A new, stabilized, hydrophilic silica packing for the high-performance gel chromatography of macromolecules 326(1985)63

Zona pellucida
Studies of fish zona pellucida by high-performance ion-exchange chromatography on agarose columns and free zone electrophoresis 341(1985)295

Zorbax Bio Series GF-250
A new, stabilized, hydrophilic silica packing for the high-performance gel chromatography of macromolecules 326(1985)63

Zygosporins
TLC of mycotoxins (Review) 334(1985)211

COLLOIDS AND SURFACES

NOW IN ITS SEVENTH YEAR!

AN INTERNATIONAL JOURNAL DEVOTED TO THE APPLICATIONS AND PRINCIPLES OF COLLOID AND INTERFACE SCIENCE.

Editor-in-Chief:
P. Somasundaran,
Henry Krumb School of Mines, Columbia University, New York, NY 10027, U.S.A.

Regional Editors:
E.D. Goddard,
Union Carbide Corp., Tarrytown Technical Center, Old Saw Mill River Road, Tarrytown, NY 10591, U.S.A.

R.J. Hunter,
Dept. of Chemistry, The University of Sydney, Sydney, N.S.W., Australia 2006

Th.F. Tadros,
Plant Protection Division, Imperial Chemical Industries Limited, Jealott's Hill Research Station, Bracknell, Berkshire, RG 12 6EY, U.K.

Scope:

COLLOIDS AND SURFACES is an international journal concerned with applications and principles of colloidal and interfacial phenomena.
It is designed to encourage publication of basic colloid and surface science and, in particular, its application in engineering and applied science. In addition to research papers, the journal contains notes, brief communications, book reviews and announcements.
Areas, topics and subjects covered include emulsions, foams, aerosols, detergency and wetting, flocculation and dispersion, rheology, cosmetics, paints, foods, paper and pulp, electrokinetic and electrode phenomena, friction and lubrication, thin films, liquid membranes and bilayers, biomaterials and biocolloids, polymer colloids, pharmaceutical sciences, health sciences, environmental and aquatic systems, water treatment and dewatering, agricultural and soil science, minerals extraction and metallurgy, precipitation, crystal growth and modification.

Subscription Information:
1986: Volumes 16-21 (in 24 issues)
US $ 575.50 / Dfl. 1554.00 including postage

Ask for a FREE SAMPLE COPY

ELSEVIER

P.O. Box 211, 1000 AE Amsterdam, The Netherlands.

52 Vanderbilt Ave. New York, N.Y. 10017.

The Dutch guilder price is definitive. US $ prices are subject to exchange rate fluctuations.

7351

Write today for your free copy of our catalogue...



Send your request to...

ELSEVIER SCIENCE PUBLISHERS

P.O. Box 211, 1000 AE Amsterdam, The Netherlands
52 Vanderbilt Avenue, New York, NY 10017, USA